Published in final edited form as:

Tetrahedron. 2006 December 11; 62(50): 11572–11579.

Syntheses of Highly Functionalized δ,γ -Unsaturated- α -Amino Acids

Siyuan Chen and Robert M. Williams*

Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523; University of Colorado Cancer Center, Aurora, CO 80045

Abstract

The synthesis of several γ,δ -unsaturated- α -amino acid derivatives is described. The method features regioselective stannylation of propargylglycine derivatives followed by Stille coupling reactions.

Keywords

syntheses; amino acids

1. Introduction

Non-proteinogenic α -amino acids are important nitrogenous building blocks that are useful for the synthesis of natural products and a multitude of biologically significant substances.¹ There has been a continuing interest in the development of new methods for the synthesis of enantiomerically pure amino acids with substituents strategically placed at side-chain positions including the capacity to install unsaturation. Many methods have been developed for the asymmetric synthesis of amino acids.^{2,3} Despite the plethora of methodologies extant, many reactive functionalities remain incompatible with existing templates.²

As part of a strategy to synthesize the natural metabolites spiroquinazoline (**1a**)⁴ and alantrypinone (**1b**),⁵ we required access to γ -aryl- γ,δ -unsaturated- α -amino acids such as **2**. Several reported methods to prepare this type of amino acid were attempted but we were unsuccessful in constructing the target substances **2**.⁶ Jackson, et al., have reported the synthesis of enantiomerically pure unsaturated α -amino acids *via* coupling to readily available serine-derived β -iodoalanine derivatives using zinc/copper reagents.⁷ We have adapted this approach to prepare a variety of γ -aryl- γ,δ -unsaturated- α -amino acids.

2. Results and Discussion

Our approach commenced with *N*-(*tert*-butoxycarbonyl)-L-iodoalanine methyl ester **3** which is commercially available or can be prepared on a large scale by using the reported procedure with serine as the starting material.⁸ Iodide **3** was converted into the corresponding Zn/Cu complex and then coupled with ethyl 3-bromopropiolate or *t*-butyl 3-bromopropiolate.⁷ The Zn/Cu complex **4** reacted with ethyl-3-bromopropiolate to produce the propargyl species **5a** and **5b** in 77% and 48% yields, respectively.

Corresponding author. Tel.: +1-970-491-6747; FAX: +1-970-491-5610; e-mail: rmw@chem.colostate.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Next, we examined the conjugate stannylation of organocopper (I) reagents to the γ,δ -alkynyl residue of **5**. The addition of compound **5a** to solution of $\text{Bu}_3\text{SnCu}\cdot\text{SMe}_2$ or $[\text{Bu}_3\text{SnCuCN}]\text{Li}^{9,10}$ in THF resulted in no reaction. When **5a** was added to $\text{Me}_3\text{SnCu}\cdot\text{SMe}_2$ in THF, compound **6** was obtained in 23% yield with 40% recovery of **5a**. Piers reported that (trimethylstannyl)copper(I) dimethylsulfide complex was relatively unreactive towards α,β -unsaturated carbonyl compounds.⁹ When a more reactive species $[\text{Me}_3\text{SnCuCN}]\text{Li}^{10}$ was used in place of $\text{Me}_3\text{SnCu}\cdot\text{SMe}_2$ for this reaction we observed that the yield was dramatically improved. When **5a** was added to $[\text{Me}_3\text{SnCuCN}]\text{Li}$ in dry THF at -78°C , compound **6** and the corresponding *Z*-isomer were produced in a nearly 1:1 ratio with no recovery of starting material. Separation of the two isomers by flash chromatography proved to be very difficult and synthetically intractable. When $[\text{Me}_3\text{SnCuCN}]\text{Li}$ was treated with EtOH prior to the addition of **5a**, the product obtained was exclusively the desired *E*-isomer. The geometric configuration of the product was determined by the coupling constant between the α -olefinic proton and the tin atom (^{117}Sn , ^{119}Sn) of the Me_3Sn group. It is well-known that when a trialkylstannyl group and a proton are vicinal on a $\text{C}=\text{C}$ bond, the $^3J_{\text{Sn-H}}$ values are much larger when these moieties are *trans*- as opposed to *cis*-configured.¹¹ The $^3J_{\text{Sn-H}}$ value in the present case of 69 Hz, falls into the expected range for the *E*-stereochemistry. The observation of a significant ^1H *nOe* between the vinyl proton and the Me_3Sn -protons provided corroborating evidence to support the *E*-stereochemistry assigned for **6** (Figure 2). The optical integrity of **6** was determined by ^1H NMR analysis of the derived Mosher's amide¹² which revealed that **6** was obtained in at least 99.5:0.5 er. Compound **6** proved to be a stable substance and can be kept as an oil, exposed to air at ambient temperature for several weeks without detectable decomposition.

Compound **6** was then coupled with a variety of aryl and heterocyclic halides under Stille cross-coupling conditions (Table 1).¹³ Our initial efforts examined $\text{Pd}(\text{PPh}_3)_4$ as the catalyst in the presence of CuI^{14} at room temperature. But under these conditions either no product or trace amount of product were obtained. Higher temperature only led to decomposition of compound **6**. Amino acid **2a** was successfully prepared by coupling of *o*-bromiodobenzene with compound **6** by using Pd_2dba_3 in the presence of AsPh_3 and CuI^{15} . Only the bromide product **2a** was observed as evidenced by ^{13}C NMR and MS spectroscopy; the corresponding iodide was not observed. The 3-cyanophenyl moiety can be introduced under similar conditions (entry c). Heterocyclic species such as 2-bromocyclopent-2-enone and 4-bromofuran-2(5*H*)-one were also successfully coupled with compound **6** to yield **2d** and **2e** in 77% and 57% yields, respectively (entries d and e). The coupling of **6** with 3-iodothiophene or 5-iodofuran-2-carboxaldehyde also provided the cross-coupling products in useful yields (entry f and g).

In all cases, the reaction was complete in just two hours and no loss of optical integrity was observed by chiral HPLC analysis.¹⁶ *o*-Iododinitrobenzene coupled with compound **6** to deliver **2b**, however in very poor yield. When $\text{PdCl}_2(\text{CH}_3\text{CN})_2$ was used as the catalyst, incomplete reaction was observed. We found that the yields could be improved by increasing catalyst loading, but this also caused more homocoupling. Eventually, it was found that treatment of $\text{PdCl}_2(\text{CH}_3\text{CN})_2$ with *tri-n*-butyltin hydride followed by addition of compound **6** and *o*-iodonitrobenzene portionwise resulted in a 79% yield of **2b** (based on a small amount of unreacted starting material).

It proved unnecessary to employ dry, deoxygenated solvent (DMF) and the reaction can be manipulated in air at ambient temperature. It should be noted that no reaction occurred when commercially available Pd-black was used. This system also proved effective for the coupling of *p*-iodobenzaldehyde with compound **6** (entry h). In this case the catalyst could be added in one portion and heating to 45°C was required. It is known that for aryl iodide substrates, the oxidative addition of Pd(0) in the Stille reaction cycle is accelerated by electron-withdrawing substituents.¹⁷ Strong electron-withdrawing groups, such as the aldehyde or nitro groups

consequently accelerate the oxidative addition step so that no ligand is needed (entries b and h). For the other halides examined, which were devoid of strong electron-withdrawing substituents, the ligand has to be added to facilitate the oxidative addition.¹⁸

In contrast to the copper (I) chemistry described above, compound **5a** underwent a Pd-catalyzed hydrostannylation reaction¹⁹ with *tri-n*-butyltin hydride providing **7a** as the major product in 71% yield along with regioisomer **7b** (10%) as shown in Scheme 2.

The two regioisomers **7a** and **7b** can be separated by flash chromatography and may be stored exposed to air for weeks without significant decomposition. The major isomer **7a** can couple with a variety of halides to give the amino acids listed in Table 2. The catalytic system Pd₂dba₃/CuI/AsPh₃ works quite well for the Stille coupling reactions of **7a** and the halides listed in Table 2. It doesn't appear to matter whether the substituent on the phenyl group is an electron-withdrawing group such as cyano, or an electron-donating group such methoxy. 2-Bromocyclopent-2-enone and 4-bromofuran-2(5H)-one can couple with compound **7a** to yield **8c** and **8d** respectively. The coupling of **7a** with 3-iodothiophene or 5-iodofuran-2-carboxaldehyde delivered **8e** and **8f** respectively in good yields.

Compound **5a** underwent hydrostannylation with good regioselectivity favoring the δ -stannane **7a**. We also examined reaction conditions by using a hindered *tert*-butyl ester that would favor the γ -stannane-type regioisomer **7b**. Compound **5b** underwent conjugate addition of [Me₃SnCuCN]Li to yield the γ -substituted stannane **9** in 53% yield (Scheme 3). As a preliminary demonstration of the utility of this species, compound **9** underwent Stille cross-coupling with *o*-iodonitrobenzene to produce the γ -aryl- γ,δ -unsaturated amino acid derivative **10** in which the two ester groups are differentiated. This amino acid, and related derivatives, are currently being examined for their utility in an intramolecular S_N2'-type strategy to access the core ring systems of spiroquinazoline and alantrypinone.

3. Conclusion

In summary, we have developed an efficient strategy to synthesize two types of γ,δ -unsaturated- α -amino acids bearing substitution at either the γ - or δ -positions. Further applications of this methodology towards the total synthesis of spiroquinazoline and other nitrogen-rich substances are currently underway and will be reported on in due course.

4. Experimental

4.1. General

All anhydrous reactions were conducted under an inert Argon atmosphere. Dry solvents were obtained from a glass contour solvent purification system. Flash column chromatography was carried out using silical gel 60. All chemical shifts (δ) are reported in parts per million. Ethyl 3-bromopropiolate and *t*-butyl 3-bromopropiolate were prepared by Leory's method.²⁰ 2-Bromocyclopent-2-enone²¹ and 4-bromofuran-2(5H)-one²² were prepared by literature procedures.

4.2. (S)-1-ethyl 6-methyl 5-(*tert*-butoxycarbonylamino)hex-2-ynedioate (**5a**).⁷

A suspension of zinc (9.83 g, 150.4 mmol) in 11.2 mL dry THF and 1,2-dibromoethane (0.65 mL, 7.53 mmol) was heated under Ar to 60 °C for 3 min. After cooling the mixture to 35 °C, trimethylsilyl chloride (0.194 mL, 1.53 mmol) was added and the mixture was vigorously stirred for 30 min. At this point the reaction vessel was warmed to 35 °C, compound **3** (8.25 g, 25.1 mmol) in 50 mL dry THF was slowly added, and the mixture was stirred for 15-40 min until no starting material remained. The solution of zinc reagent was then converted to the zinc-copper reagent **4** by the following procedure. The solution of zinc reagent was cooled to -10

°C, and a solution prepared from CuCN (2.27 g, 25.1 mmol) and LiCl (2.15 g, 50.2 mmol) in 50 mL dry THF was added. The mixture was stirred at 0 °C for 10 min and then cooled to -55 °C. A solution of ethyl 3-bromoprop-2-ynoate (5.91 g, 33.4 mmol) in 67 mL dry THF was introduced followed by stirring at this temperature for 20 h. After quenching with saturated aqueous NH₄Cl, the mixture was extracted with ethyl acetate (3×400 mL), the combined organic layers were washed with 400 mL water, dried (MgSO₄) and concentrated *in vacuo*. The residue was separated by silica gel flash chromatography (eluted with 20 % ethyl acetate/hexane) to give 6.2 g of **5a** as a yellow oil (yield 77%). $[\alpha]_D^{20} +55.7$ (c 0.670, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆): δ 0.79 (3H, t, *J*=7.2 Hz), 1.38 (9H, s), 2.39 (1H, dd, *J*=17.4 Hz, 5.4 Hz), 2.54 (1H, dd, *J*=17.4 Hz, 5.1 Hz), 3.18 (3H, s), 3.81 (2H, q, *J*=7.2 Hz), 4.34 (1H, ddd, *J*=7.8 Hz, 5.4 Hz, 5.1 Hz), 5.29 (1H, d, *J*=7.8 Hz). ¹³C NMR (75 MHz, C₆D₆): δ 14.3, 23.3, 28.7, 52.5, 52.6, 62.1, 76.6, 80.3, 83.6, 153.5, 155.4, 170.7. IR (NaCl, neat) 3366, 2980, 2240, 1748, 1712, 1507. HRMS (FAB+) calc. mass for C₁₄H₂₂NO₆ 300.1447 (M+1), found: 300.1453. *R_f* 0.33 (eluted with 20 % ethyl acetate/hexane).

4.3. (S)-1-*tert*-butyl 6-methyl 5-(*tert*-butoxycarbonylamino)hex-2-ynedioate (**5b**)

A procedure similar to the one described above for the preparation of **5a** was used for the synthesis of **5b** starting from 2.1 g (6.38 mmol) of compound **3** and 1.44 g (7.02 mmol) of *tert*-butyl 3-bromoprop-2-ynoate. Yield: 1.0 g (48%). $[\alpha]_D^{20} +52.5$ (c 1.46, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆): δ 1.24 (9H, s), 1.37 (9H, s), 2.39 (1H, dd, *J*=17.1 Hz, 5.4 Hz), 2.53 (1H, dd, *J*=17.1 Hz, 5.1 Hz), 3.15 (3H, s), 4.34 (1H, ddd, *J*=7.8 Hz, 5.4 Hz, 5.1 Hz), 5.26 (1H, d, *J*=7.8 Hz). ¹³C NMR (75 MHz, C₆D₆): δ 23.1, 28.1, 28.6, 52.3, 52.5, 77.8, 80.2, 81.3, 83.0, 152.7, 155.4, 170.8. IR (NaCl, neat) 3366, 2980, 2244, 1750, 1709, 1505. HRMS (FAB+) calc. mass for C₁₆H₂₆NO₆ 328.1760 (M+1), found 328.1746. *R_f* 0.33 (eluted with 20 % ethyl acetate/hexane).

4.4. (S,E)-1-ethyl 6-methyl 5-(*tert*-butoxycarbonylamino)-3-(trimethylstannyl)hex-2-enedioate (**6**)⁹

To a cold (-20 °C), stirred solution of 5.2 g of hexamethylditin (15.87 mmol) in 150 mL dry THF was added a solution of 10.7 mL (17.12 mmol) of a 1.6M solution of methyl lithium in ether. The mixture was stirred -20 °C for 15 min to afford a pale yellow solution of Me₃SnLi. This solution was then cooled to -78 °C and 1.66 g (18.52 mmol) of solid CuCN was added in one portion. The mixture was stirred at -78 °C for 5 min and at -48 °C for 15 min to afford a bright orange solution. This solution was then cooled to -78 °C and 0.95 mL (16.0 mmol) of dry ethanol was added. After 5 min, a solution of compound **5a** in 50 mL dry THF was added dropwise and the mixture was stirred at -78 °C for 4 h. 200 mL NH₄Cl-NH₄OH buffer (consisting of a 9:1 ratio of sat. aq. NH₄Cl: 28%~30% NH₄OH, pH=8) was added, the mixture was opened to the atmosphere, was allowed to warm to rt, and was stirred vigorously until the aqueous phase became a deep blue color. The organic phase was separated and the aqueous phase was extracted thoroughly with ether. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by silica gel flash column chromatography (eluted with 15 % ethyl acetate/hexane) to give 5.0 g of compound **6** as a yellow oil (81% yield). $[\alpha]_D^{20} +60.8$ (c 2.65, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆): δ 0.18 (9H, s), 0.95 (3H, t, *J*=7.2 Hz), 1.36 (9H, s, ²*J*_{Sn-H} = 54.3 Hz), 2.89 (1H, dd, *J*=12.0 Hz, 3.9 Hz), 3.28 (3H, s), 3.90 (1H, t, *J*=12.0 Hz), 3.95 (2H, q, *J*=7.2 Hz), 4.68 (1H, *J*=12.0 Hz, 8.1 Hz, 3.9 Hz), 6.09 (1H, d, *J*=8.1 Hz), 6.22 (1H, s, ³*J*_{Sn-H} = 69.0 Hz). ¹³C NMR (75 MHz, C₆D₆): δ -8.7, 28.7, 28.9, 37.2, 52.3, 54.5, 79.5, 81.0, 133.4, 156.2, 164.9, 165.5, 172.8. IR (NaCl, neat) 3371, 2977, 1751, 1717, 1598, 1507, 1446 cm⁻¹. HRMS (FAB+) calc. mass for C₁₉H₃₆NO₆Sn: 494.1565 (M+1), found: 494.1557. *R_f* 0.33 (eluted with 15 % ethyl acetate/hexane).

4.5. (S,E)-1-ethyl 6-methyl 3-(2-bromophenyl)-5-(tert-butoxycarbonylamino)hex-2-enedioate (2a)

A solution of *o*-iodobromobenzene (41.2 mg, 0.146 mmol), CuI (6.67 mg, 35 μ mol), AsPh₃ (10.7 mg, 35 μ mol), and Pd₂dba₃ (9.45 mg, 8.74 μ mol) in 1.2 mL dry DMF under Ar was treated with compound **6** (61.3 mg, 0.132 mmol). The reaction mixture was then stirred for 2 h. The solution was then diluted with EtOAc (5 mL) and washed with water. The combined aqueous layers were back-extracted with EtOAc (2 $\times$ 5 mL). The combined organic extracts were dried over Na₂SO₄, filtered, and evaporated to dryness. The resulting oil was purified by silica gel chromatography (eluted with 25% EtOAc in hexanes) to yield 41.0 mg of **2a** as a yellow oil (yield 68%). [α]_D²⁰ +53.2 (*c* 1.17, CH₂Cl₂). ¹H NMR (400 MHz, C₆D₆): δ 0.91 (3H, t, *J*=7.2 Hz), 1.41 (9H, s), 3.13 (3H, s), 3.33 (1H, dd, *J*=12.4 Hz, 4.0 Hz), 3.88 (2H, q, *J*=7.2 Hz), 4.02 (1H, t=12.4 Hz), 4.70 (1H, ddd, *J*=12.4 Hz, 6.6 Hz, 4.0 Hz), 5.89 (1H, s), 5.96 (1H, d, *J*=6.6 Hz), 6.58-7.25 (4H, m). ¹³C NMR (75 MHz, C₆D₆): δ 14.4, 28.7, 35.4, 52.1, 53.3, 60.9, 79.6, 121.8, 124.6, 127.9, 128.9, 130.1, 130.9, 133.6, 142.5, 156.2, 167.0, 172.5. IR (NaCl, neat) 3382, 2978, 1748, 1717, 1645, 1507, 1436 cm⁻¹. HRMS (FAB+) calc. mass for C₂₀H₂₇NO₆Br: 456.1021 (*M*+1, ⁸⁰Br), found: 456.1005. *R*_f 0.30 (eluted with 25 % ethyl acetate/ hexane).

4.6. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(2-nitrophenyl)hex-2-enedioate (2b)

A 25 mL flask, equipped with a magnetic stir bar was charged with 80 mg (0.308 mmol) of PdCl₂(CH₃CN)₂ and 10 mL of DMF. Tri-*n*-butyltin hydride (182 μ L, 0.339 mmol) was added dropwise and a black precipitate was formed immediately. The mixture was stirred for 10 minutes and was added to a solution of compound **6** (1.98 g, 4.27 mmol) and 1.12 g (4.49 mmol) of *o*-iodonitrobenzene in DMF (50 mL) in three portions (one portion every 5 h). The mixture was stirred overnight. The reaction was then quenched with water and the mixture was extracted with ether (3 $\times$ 100 mL). The combined organic extracts were washed with 300 mL water, dried over Na₂SO₄ and concentrated *in vacuo*. The residue was separated by silica gel flash column chromatography (eluted with 25 % ethyl acetate/ hexane) to give 850 mg (47 % conversion and 79% yield based on recovered **6**) of **2b** as a yellow oil along with 800 mg of unreacted **6**. [α]_D²⁰ +17.8 (*c* 0.835, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆): δ 0.89 (3H, t, *J*=7.2 Hz), 1.42 (9H, s), 3.11 (3H, s), 3.16 (1H, dd, *J*=12.9 Hz, 3.6 Hz), 3.85 (2H, q, *J*=7.2 Hz), 4.03 (1H, t, *J*=12.9 Hz), 4.64 (1H, ddd, *J*=12.9 Hz, 9.0 Hz, 3.6 Hz), 5.75 (1H, s), 5.9 (1H, d, *J*=9.0 Hz), 6.5-7.6 (4H, m). ¹³C NMR (75 MHz, C₆D₆): δ 14.8, 28.4, 35.6, 52.1, 53.3, 79.6, 81.4, 124.2, 125.3, 128.9, 131.7, 133.6, 137.3, 147.4, 153.4, 156.3, 166.3, 172.4. IR (NaCl, neat) 3389, 2979, 1747, 1716, 1640, 1608, 1528, 1367, 1347 cm⁻¹. HRMS (FAB+) calc. mass for C₂₂H₃₁N₂O₈: 451.2080 (*M*+1), found: 451.2086. *R*_f 0.30 (eluted with 25 % ethyl acetate/ hexane).

4.7. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(3-cyanophenyl)hex-2-enedioate (2c)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **2c** starting from 19.5 mg (0.0421 mmol) of compound **6** and 10.6 mg (0.0464 mmol) of 3-iodobenzonitrile. Yield: 12.0 mg (71%). [α]_D²⁰ +46.7 (*c* 0.405, CH₂Cl₂). ¹H NMR (400 MHz, C₆D₆): δ 0.99 (3H, t, *J*= 7.0 Hz), 1.36 (9H, s), 3.16 (3H, s), 3.20 (1H, dd, *J*=13.4 Hz, 10.0 Hz), 3.59 (1H, dd, *J*=13.4 Hz, 9.6 Hz), 3.96 (2H, q, *J*=7.0 Hz), 4.46 (1H, ddd, *J*=10.0 Hz, 9.6 Hz, 8.4 Hz), 5.71 (1H, d, *J*=8.4 Hz), 5.81 (1H, s), 6.40-7.20 (3H, m), 7.21 (1H, s). ¹³C NMR (100 MHz, C₆D₆): δ 14.5, 28.6, 33.4, 52.1, 53.5, 60.9, 79.8, 113.9, 118.7, 122.8, 129.7, 130.9, 131.1, 132.6, 141.5, 153.0, 155.9, 166.7, 172.2. IR (NaCl, neat) 3371, 2878, 2231, 1744, 1713, 1632, 1510 cm⁻¹. HRMS (FAB+) calc. mass for C₂₁H₂₇N₂O₆: 403.1869 (*M*+1), found 403.1869. *R*_f 0.30 (eluted with 25 % ethyl acetate/ hexane).

4.8. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(5-oxocyclopent-1-enyl)hex-2-enedioate (2d)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **2d** starting from 21.0 mg (0.0454 mmol) of compound **6** and 8.05 mg (0.05 mmol) of 4-bromocyclopent-2-enone. Yield: 13.3 mg (77%). $[\alpha]_D^{20} +46.5$ (c 0.550, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆) : δ 0.91 (3H, t, *J*=6.9 Hz), 1.39 (9H, s), 1.58 (2H, m), 1.78 (2H, t, *J*=4.5 Hz), 3.07 (1H, dd, *J*=13.5 Hz, 5.1 Hz), 3.33 (3H, s), 3.74 (1H, *J*=13.5 Hz, 10.8 Hz), 3.91 (2H, q, *J*=6.9 Hz), 4.70 (1H, ddd, *J*=10.8 Hz, 8.1 Hz, 6.9 Hz), 6.37 (1H, d, *J*=8.1 Hz), 7.41 (1H, t, *J*=3.0 Hz), 7.66 (1H, s). ¹³C NMR (100 MHz, C₆D₆) : δ 14.7, 26.3, 28.9, 32.6, 36.2, 52.4, 54.5, 61.0, 79.8, 122.5, 140.5, 144.0, 156.4, 163.0, 168.4, 172.8, 205.7. IR (NaCl, neat) 3371, 2978, 1746, 1708, 1625, 1510, 1367 cm⁻¹. HRMS (FAB+) calc. mass for C₁₉H₂₈NO₇ 382.1866 (M+1), found 382.1873. *R*_f 0.30 (eluted with 50 % ethyl acetate/ hexane).

4.9. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(5-oxo-2,5-dihydrofuran-3-yl)hex-2-enedioate (2e)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **2e** starting from 45.0 mg (0.097 mmol) of compound **6** and 17.4 mg (0.107 mmol) of 4-bromofuran-2(5H)-one. Yield: 21.1 mg (57%). $[\alpha]_D^{20} +36.0$ (c 2.19, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆) : δ 0.93 (3H, t, *J*=7.2 Hz), 1.35 (9H, s), 2.90 (1H, dd, *J*=13.2 Hz, 4.8 Hz), 3.21 (1H, dd, *J*=13.2 Hz, 9.6 Hz), 3.24 (3H, s), 3.87 (2H, q, *J*=7.2 Hz), 3.96 (2H, s), 4.69 (1H, ddd, *J*=9.6 Hz, 8.1 Hz, 4.8 Hz), 5.34 (1H, s), 5.62 (1H, d, *J*=8.1 Hz), 6.15 (1H, s). ¹³C NMR (75 MHz, C₆D₆) : δ 14.4, 28.6, 32.3, 52.4, 53.9, 61.3, 70.0, 80.0, 120.3, 123.2, 143.9, 155.9, 162.1, 166.0, 171.9, 172.2. IR (NaCl, neat) 3376, 2979, 1789, 1752, 1714, 1630, 1599, 1507 cm⁻¹. HRMS (FAB+) calc. mass for C₁₈H₂₆NO₈ 384.1658 (M+1), found 384.1673. *R*_f 0.30 (eluted with 50 % ethyl acetate/ hexane).

4.10. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(thiophen-3-yl)hex-2-enedioate (2f)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **2f** starting from 19.6 mg (0.0423 mmol) of compound **6** and 9.8 mg (0.0466 mmol) of 3-iodothiophene. Yield: 13.0 mg (80%). $[\alpha]_D^{20} +46.1$ (c 0.460, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆) : δ 0.99 (3H, t, *J*=7.1 Hz), 1.38 (9H, s), 3.27 (1H, dd, *J*=12.9 Hz, 5.1 Hz), 3.28 (3H, s), 3.76 (1H, dd, *J*=12.9 Hz, 9.9 Hz), 3.96 (2H, q, *J*=6.9 Hz), 4.77 (1H, ddd, *J*=7.8 Hz, 6.9 Hz, 5.1 Hz), 6.22 (1H, d, *J*=7.8 Hz), 6.26 (1H, s), 6.65 (1H, dd, *J*=5.1 Hz, 3.0 Hz), 6.88 (1H, dd, *J*=5.1 Hz, 1.5 Hz), 7.35 (1H, dd, *J*=3.0 Hz, 1.5 Hz). ¹³C NMR (100 MHz, C₆D₆) : δ 14.8, 28.9, 33.4, 52.3, 54.9, 60.8, 79.7, 118.5, 125.8, 125.9, 126.9, 141.3, 148.7, 156.1, 167.8, 172.6. IR (NaCl, neat) 3366, 2978, 1745, 1711, 1698, 1619, 1503, 1450 cm⁻¹. HRMS (FAB+) calc. mass for C₁₈H₂₆NO₆S 384.1481 (M+1), found 384.1489. *R*_f 0.40 (eluted with 20% ethyl acetate/ hexane).

4.11. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(5-formylfuran-2-yl)hex-2-enedioate (2g)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **2g** starting from 53.5 mg (0.116 mmol) of compound **6** and 28.2 mg (0.127 mmol) of 5-iodofuran-2-carbaldehyde. Yield: 38.4 mg (84%). $[\alpha]_D^{20} +27.2$ (c 1.13, CH₂Cl₂). ¹H NMR (300 MHz, C₆D₆) : δ 0.94 (3H, t, *J*=7.1 Hz), 1.36 (9H, s), 3.17 (1H, dd, *J*=13.2 Hz, 5.7 Hz), 3.29 (3H, s), 3.50 (1H, dd, *J*=13.2 Hz, 9.9 Hz), 3.90 (2H, q, *J*=7.1 Hz), 4.69 (1H, ddd, *J*=9.9 Hz, 8.1 Hz, 5.7 Hz), 6.02 (1H, d, *J*=8.1 Hz), 6.37 (1H, d, 4.2 Hz), 6.55 (1H, d, 4.2 Hz), 6.67 (1H, s), 9.13 (1H, s). ¹³C NMR (75 MHz, C₆D₆) : δ 14.4, 28.6, 31.1, 52.3, 54.7, 61.0, 79.9, 113.9, 119.6, 121.1, 140.6, 153.4, 156.2, 156.7, 167.2, 172.2, 177.7. IR (NaCl, neat) 3366,

2978, 1746, 1713, 1683, 1623, 1500, 1447 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{19}\text{H}_{26}\text{NO}_8$ 396.1658 (M+1), found 396.1649. R_f 0.25 (eluted with 25 % ethyl acetate/ hexane).

4.12. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(4-formylphenyl)hex-2-enedioate (2h)

A procedure similar to the one described above for the preparation of **2b** was used for the synthesis of **2h** starting from 45.0 mg (0.0972 mmol) of compound **6** and 21.6 mg (0.117 mmol) of 4-iodobenzaldehyde. Yield: 26.4 mg (67%). $[\alpha]^{20}_{\text{D}} +66.8$ (c 0.467, CH_2Cl_2). ^1H NMR (300 MHz, C_6D_6): δ 0.99 (3H, t, $J=6.9$ Hz), 1.37 (9H, s), 3.19 (3H, s), 3.28 (1H, dd, $J=13.2$ Hz, 5.1 Hz), 3.75 (1H, dd, $J=13.2$ Hz, 9.9 Hz), 3.87 (2H, q, $J=6.9$ Hz), 4.55 (1H, ddd, $J=9.9$ Hz, 8.4 Hz, 5.1 Hz), 5.80 (1H, d, $J=8.4$ Hz), 6.03 (1H, s), 7.12 (2H, d, $J=4.8$ Hz), 7.39 (2H, d, $J=4.8$ Hz), 9.59 (1H, s). ^{13}C NMR (400 MHz, C_6D_6): δ 14.5, 28.7, 33.4, 52.1, 53.7, 60.9, 79.7, 122.7, 130.0, 130.2, 137.4, 145.7, 154.2, 155.9, 166.9, 172.4, 191.0. IR (NaCl, neat) 3371, 2979, 1745, 1704, 1604, 1511 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{21}\text{H}_{28}\text{NO}_7$ 406.1866 (M+1), found 406.1874. R_f 0.30 (eluted with 33 % ethyl acetate/ hexane).

4.13. (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-2-(tri-*n*-butylstannyl)hex-2-enedioate (7a) and (S,E)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-3-(tri-*n*-butylstannyl)hex-2-enedioate (7b).¹⁸

A solution of compound **5a** (347.8 mg, 0.22 mmol), and $\text{Pd}(\text{PPh}_3)_4$ (5 mg, 4.3 mmol) in dry THF (10 mL) under Ar was treated with tri-*n*-butyltin hydride (343 mmol, 0.24 mmol). The reaction mixture was then stirred for 2 h and the solvent was evaporated to dryness. The resulting oil was purified by silica gel chromatography (eluted with 5% ethyl acetate, 95% hexane). The fast eluting compound was **7b** (66 mg, 10%) and the slower eluting substance was **7a** (487 mg, yield 71%). Data for **7a**: $[\alpha]^{20}_{\text{D}} +33.5$ (c 1.10, CH_2Cl_2). ^1H NMR (400 MHz, C_6D_6): δ 0.96 (9H, t, $J=7.6$ Hz), 1.01 (3H, t, $J=7.2$ Hz), 1.02 (6H, t, $J=8.8$ Hz), 1.37 (6H, m), 1.42 (9H, s), 1.59 (6H, m), 2.90 (2H, m), 3.28 (3H, s), 3.95 (2H, q, $J=7.2$ Hz), 4.58 (1H, ddd, $J=13.2$ Hz, 7.2 Hz, 0.4 Hz), 5.76 (1H, d, $J=7.2$ Hz), 6.09 (1H, t, $J=7.4$ Hz, $^3J_{\text{Sn-H}}=45.0$ Hz). ^{13}C NMR (100 MHz, C_6D_6): δ 11.0, 14.3, 14.7, 28.0, 28.7, 29.6, 35.2, 52.1, 54.0, 60.8, 79.6, 141.1, 146.9, 156.1, 171.1, 172.8. IR (NaCl, neat) 3364, 2957, 2928, 2872, 2854, 1754, 1719, 1607, 1500 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{26}\text{H}_{50}\text{NO}_6\text{Sn}$: 591.2671 (M+1), found: 591.2647. R_f 0.33 (eluted with 10 % ethyl acetate/ hexane). Data for **7b**: $[\alpha]^{20}_{\text{D}} +54.3$ (c 3.30, CH_2Cl_2). ^1H NMR (400 MHz, C_6D_6): δ 0.93 (9H, t, $J=7.4$ Hz), 0.94 (3H, t, $J=7.0$ Hz), 1.02 (6H, t, $J=8.2$ Hz), 1.36 (6H, m), 1.41 (9H, s), 1.58 (6H, m), 2.97 (1H, dd, $J=12.4$ Hz, 3.6 Hz), 3.31 (3H, s), 3.92 (2H, q, $J=7.0$ Hz), 4.02 (2H, t, $J=12.4$ Hz), 4.71 (1H, ddd, $J=12.4$ Hz, 12.4 Hz, 8.4 Hz), 6.23 (1H, d, $J=8.4$ Hz), 6.34 (1H, s, $^3J_{\text{Sn-H}}=60.0$ Hz). ^{13}C NMR (100 MHz, C_6D_6): δ 10.7, 14.2, 14.5, 28.1, 28.8, 29.7, 37.6, 52.2, 54.6, 60.6, 79.5, 132.1, 156.5, 165.3, 168.1, 173.0. IR (NaCl, neat) 3364, 2957, 2929, 2872, 2854, 1754, 1719, 1607, 1503 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{26}\text{H}_{50}\text{NO}_6\text{Sn}$: 591.2671 (M+1), found: 591.2671. R_f 0.30 (eluted with 15 % ethyl acetate / hexanes).

4.14. (S,Z)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-2-(4-methoxyphenyl)hex-2-enedioate (8a)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **8a** starting from 40.0 mg (0.0678 mmol) of compound **7a** and 17.4 mg (0.0745 mmol) of 1-iodo-4-methoxybenzene. Yield: 20.4 mg (74%). $[\alpha]^{20}_{\text{D}} +20.9$ (c 0.675, CH_2Cl_2). ^1H NMR (300 MHz, C_6D_6): δ 0.94 (3H, t, $J=7.1$ Hz), 1.42 (9H, s), 2.81 (2H, t, $J=7.2$ Hz), 3.25 (3H, s), 3.28 (3H, s), 4.01 (2H, q, $J=7.1$ Hz), 4.61 (1H, q, $J=7.5$ Hz), 5.71 (1H, d, $J=7.5$ Hz), 5.89 (1H, t, $J=7.5$ Hz), 6.72 (2H, d, $J=8.7$ Hz), 7.23 (2H, d, $J=8.7$ Hz). ^{13}C NMR (75 MHz, C_6D_6): δ 14.4, 28.7, 33.5, 52.1, 54.1, 55.1, 61.2, 79.8, 114.4, 129.3, 130.6, 131.6, 138.5, 156.1, 160.4, 168.6, 172.8. IR (NaCl, neat) 3371, 2978, 1745, 1715, 1608, 1513 cm^{-1} .

HRMS (FAB+) calc. mass for $C_{21}H_{30}NO_7$ 408.2022 (M+1), found 408.2003. R_f 0.33 (eluted with 33 % ethyl acetate/ hexane).

4.15. (S,Z)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-2-(3-cyanophenyl)hex-2-enedioate (8b)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **8b** starting from 41.3 mg (0.070 mmol) of compound **7a** and 17.6 mg (0.077 mmol) of 3-iodobenzonitrile. Yield: 22.0 mg (79%). $[\alpha]^{20}_D +27.4$ (c 0.625, CH_2Cl_2). 1H NMR (300 MHz, C_6D_6): δ 0.84 (3H, t, $J=6.9$ Hz), 1.41 (9H, s), 2.82 (1H, t, $J=7.5$ Hz), 3.25 (3H, s), 3.88 (2H, q, $J=6.9$ Hz), 4.57 (1H, q, $J=7.5$ Hz), 5.53 (1H, d, $J=7.2$ Hz), 5.75 (1H, t, $J=7.8$ Hz), 6.61 (1H, t, $J=7.5$ Hz), 6.8-7.4 (4H, m). ^{13}C NMR (100 MHz, C_6D_6): δ 14.3, 28.7, 33.7, 52.3, 53.8, 61.5, 80.1, 113.5, 118.9, 129.2, 131.5, 131.8, 132.1, 136.1, 138.1, 139.5, 156.0, 166.8, 172.5. IR (NaCl, neat) 3373, 2979, 2231, 1745, 1716, 1511, 1438 cm^{-1} . HRMS (FAB+) calc. mass for $C_{21}H_{27}N_2O_6$ 403.1869 (M+1), found 403.1864. R_f 0.30 (eluted with 25 % ethyl acetate/ hexane).

4.16. (S,Z)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-2-(5-oxocyclopent-1-enyl)hex-2-enedioate (8c)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **8c** starting from 42.4 mg (0.0718 mmol) of compound **7a** and 12.7 mg (0.079 mmol) of 4-bromocyclopent-2-enone. Yield: 19.7 mg (72%). $[\alpha]^{20}_D +20.2$ (c 0.130, CH_2Cl_2). 1H NMR (300 MHz, C_6D_6): δ 1.01 (3H, t, $J=7.2$ Hz), 1.66 (9H, s), 1.68 (2H, m), 1.85 (2H, m), 2.83 (2H, m), 3.30 (3H, s), 4.05 (2H, q, $J=7.2$ Hz), 4.59 (1H, m), 5.64 (1H, d, $J=6.9$ Hz), 6.82 (1H, t, $J=7.8$ Hz), 6.96 (1H, t, $J=2.9$ Hz). ^{13}C NMR (75 MHz, C_6D_6): δ 14.5, 26.3, 28.7, 33.2, 35.3, 52.2, 53.9, 61.4, 79.7, 130.0, 135.6, 140.5, 156.0, 160.0, 167.0, 172.6, 205.6. IR (NaCl, neat) 3367, 2978, 1745, 1709, 1509, 1367 cm^{-1} . HRMS (FAB+) calc. mass for $C_{19}H_{28}NO_7$ 382.1866 (M+1), found 382.1874. R_f 0.33 (eluted with 50 % ethyl acetate/ hexane).

4.17. (S,Z)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-2-(5-oxo-2,5-dihydrofuran-3-yl)hex-2-enedioate (8d)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **8e** starting from 36.0 mg (0.061 mmol) of compound **7a** and 10.93 mg (0.0671 mmol) of 4-bromofuran-2(5H)-one. Yield: 16.0 mg (69%). $[\alpha]^{20}_D +20.8$ (c 1.00, CH_2Cl_2). 1H NMR (300 MHz, C_6D_6): δ 0.82 (3H, t, $J=7.2$ Hz), 1.37 (9H, s), 2.65 (1H, m), 2.66 (1H, t, $J=8.1$ Hz), 3.21 (3H, s), 3.81 (2H, q, $J=7.2$ Hz), 4.11 (2H, s), 4.40 (1H, dd, $J=14.1$ Hz, 8.1 Hz), 5.36 (1H, d, $J=8.1$ Hz), 5.54 (1H, t, $J=7.8$ Hz), 5.98 (1H, s). ^{13}C NMR (75 MHz, C_6D_6): δ 14.2, 28.6, 33.7, 52.4, 53.5, 61.8, 70.5, 80.3, 117.5, 129.5, 140.4, 155.9, 158.7, 164.9, 172.1, 172.9. IR (NaCl, neat) 3364, 2979, 1785, 1750, 1716, 1634, 1516 cm^{-1} . HRMS (FAB+) calc. mass for $C_{18}H_{26}NO_8$ 384.1658 (M+1), found 384.1649. R_f 0.30 (eluted with 50 % ethyl acetate/ hexane).

4.18. (S,Z)-1-ethyl 6-methyl 5-(tert-butoxycarbonylamino)-2-(thiophen-3-yl)hex-2-enedioate (8e)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **8e** starting from 35.6 mg (0.0603 mmol) of compound **7a** and 13.9 mg (0.0633 mmol) of 3-iodothiophene. Yield: 19.4 mg (84%). $[\alpha]^{20}_D +17.0$ (c 0.635, CH_2Cl_2). 1H NMR (300 MHz, C_6D_6): δ 0.91 (3H, t, $J=7.1$ Hz), 1.40 (9H, s), 2.76 (2H, t, $J=7.5$ Hz), 3.25 (3H, s), 3.96 (2H, q, $J=7.1$ Hz), 4.57 (1H, q, $J=7.5$ Hz), 5.64 (1H, d, $J=7.5$ Hz), 5.95 (1H, t, $J=8.1$ Hz), 6.77 (1H, dd, $J=5.1$ Hz, 3.3 Hz), 6.94 (1H, dd, $J=5.1$ Hz, 1.2 Hz), 7.08 (1H, dd, $J=3.3$ Hz, 1.2 Hz). ^{13}C NMR (75 MHz, C_6D_6): δ 14.6, 28.9, 33.5, 52.3, 54.2, 61.5, 79.9, 123.2, 125.9, 127.2,

132.3, 133.4, 138.6, 156.0, 168.5, 172.6. IR (NaCl, neat) 3372, 2978, 1743, 1717, 1506, 1436, 1366 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{18}\text{H}_{26}\text{NO}_6\text{S}$ 384.1481 (M+1), found 384.1486. R_f 0.30 (eluted with 25 % ethyl acetate/ hexane).

4.19. (S,Z)-1-ethyl 6-methyl 5-(*tert*-butoxycarbonylamino)-2-(5-formylfuran-2-yl)hex-2-enedioate (8f)

A procedure similar to the one described above for the preparation of **2a** was used for the synthesis of **8f** starting from 40.3 mg (0.0683 mmol) of compound **7a** and 16.7 mg (0.0751 mmol) of 5-iodofuran-2-carbaldehyde. Yield: 20.5 mg (76%). $[\alpha]_D^{20} -1.3$ (c 0.555, CH_2Cl_2). ^1H NMR (300 MHz, C_6D_6): δ 0.92 (3H, t, $J=7.2$ Hz), 1.39 (9H, s), 2.78 (2H, t, m), 3.23 (3H, s), 3.93 (2H, q, $J=7.2$ Hz), 4.50 (1H, m), 5.42 (1H, d, $J=7.8$ Hz), 6.36 (1H, d, $J=3.6$ Hz), 6.46 (1H, d, $J=3.6$ Hz), 6.66 (1H, t, $J=7.95$ Hz), 9.22 (1H, s). ^{13}C NMR (75 MHz, C_6D_6): δ 14.3, 28.6, 33.2, 52.2, 53.7, 61.8, 80.0, 111.8, 121.8, 127.0, 137.5, 152.8, 154.9, 155.9, 165.0, 172.4, 177.0. IR (NaCl, neat) 3365, 2979, 1742, 1716, 1678, 1500, 1439 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{19}\text{H}_{26}\text{NO}_8$ 396.1658 (M+1), found 396.1640. R_f 0.30 (eluted with 33 % ethyl acetate/ hexane).

4.20. (S,E)-1-*tert*-butyl 6-methyl 5-(*tert*-butoxycarbonylamino)-3-(trimethylstannyl)hex-2-enedioate (9)

A procedure similar to the one described above for the preparation of **6** was used for the synthesis of **9** starting from 0.91 mg (2.78 mmol) of compound **5b** and 1.09 g (3.34 mmol) of hexamethylditin. Yield: 729 mg (53%). $[\alpha]_D^{20} +60.8$ (c 2.65, CH_2Cl_2). ^1H NMR (300 MHz, C_6D_6): δ 0.13 (9H, s, $^2J_{\text{Sn-H}} = 54.3$ Hz), 1.36 (9H, s), 1.42 (9H, s), 2.89 (1H, dd, $J=12.0$ Hz, 3.9 Hz), 3.28 (3H, s), 3.95 (1H, t, $J=12.0$ Hz), 4.68 (1H, $J=12.0$ Hz, 8.1 Hz, 3.9 Hz), 6.09 (1H, d, $J=8.1$ Hz), 6.22 (1H, s, $^3J_{\text{Sn-H}} = 69.0$ Hz). ^{13}C NMR (75 MHz, C_6D_6): δ -8.7, 28.7, 28.9, 37.2, 52.3, 54.5, 79.5, 81.0, 133.4, 156.2, 164.9, 165.5, 172.8. IR (NaCl, neat) 3371, 2977, 1751, 1717, 1598, 1507, 1446 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{19}\text{H}_{36}\text{NO}_6\text{Sn}$ 494.1565 (M+1), found 494.1557. R_f 0.33 (eluted with 15 % ethyl acetate/ hexane).

4.21. (S,E)-1-*tert*-butyl 6-methyl 5-(*tert*-butoxycarbonylamino)-3-(2-nitrophenyl)hex-2-enedioate (10)

A procedure similar to the one described above for the preparation of **2b** was used for the synthesis of **10** starting from 404 mg (1.03 mmol) of compound **9** and 284 mg (1.14 mmol) of *o*-iododinitrobenzene. 94.9 mg product was obtained along with the recovery of 246 mg compound **9** (25% conversion and 66% yield based on recovered **9**). $[\alpha]_D^{20} +17.8$ (c 0.835, CH_2Cl_2). ^1H NMR (300 MHz, C_6D_6): δ 1.36 (9H, s), 1.42 (9H, s), 3.11 (3H, s), 3.16 (1H, dd, $J=12.9$ Hz, 3.6 Hz), 4.03 (1H, t, $J=12.9$ Hz), 4.64 (1H, ddd, $J=12.9$ Hz, 9.0 Hz, 3.6 Hz), 5.75 (1H, s), 6.02 (1H, d, $J=9.0$ Hz), 6.5-7.6 (4H, m). ^{13}C NMR (75 MHz, C_6D_6): δ 28.4, 28.8, 35.6, 52.1, 53.3, 79.6, 81.4, 124.2, 125.3, 128.9, 131.7, 133.6, 137.3, 147.4, 153.4, 156.3, 166.3, 172.4. IR (NaCl, neat) 3389, 2979, 1747, 1716, 1640, 1608, 1528, 1367, 1347 cm^{-1} . HRMS (FAB+) calc. mass for $\text{C}_{22}\text{H}_{31}\text{N}_2\text{O}_8$ 451.2080 (M+1), found 451.2086. R_f 0.30 (eluted with 25% ethyl acetate/ hexane).

Acknowledgements

We thank the National Institutes of Health for financial support (GM068011).

References and Notes

1. For a few representative examples of amino acid-containing natural products see: (a) Kaneko I, Kamoshida K, Takahashi S. *J Antibiotics* 1989;42:236–241. [PubMed: 2925515] (b) Matasunaga S, Fusetani N, Hashimoto K, Walchi M. *J Am Chem Soc* 1989;111:2582–2588. (c) Helms HL, Moore

- RE, Niemczura WP, Patterson GML, Tomer KB, Gross ML. *J Org Chem* 1988;53:1298–1307. (d) McMillan JA, Paul IC, Goo YM, Rinehart KL. *Tetrahedron Lett* 1981;22:39–42.
2. For reviews see :Williams, RM. *Synthesis of Optically Active Amino Acids*. 1. Pergamon Press; Oxford [England], New York: 1989. O'donnell, MJ., editor. *Tetrahedron*. 44. 1988. p. 5253-5614. (c) Izumi Y, Chibata I, Itoh T. *Angew Chem Int Ed Engl* 1978;17:176–183. [PubMed: 417646]
3. For examples of amino acids natural products that have been synthesized see(a) Lane JW, Chen Y, Williams RM. *J Am Chem Soc* 2005;127:12684–12690. [PubMed: 16144418] (b) White JD, Amedeo JC. *J Org Chem* 1989;54:738–743. (c) Nishiyama S, Suzuki Y, Yamamura S. *Tetrahedron Lett* 1988;29:559–562.
4. Barrow CJ, Sun HH. *J Nat Prod* 1994;57:471–476. [PubMed: 7517439]
5. (a) Barrow CJ, Sun HH. *J Nat Prod* 1998;61:1154–1157. [PubMed: 9748389] For total syntheses of alantropinone see: (b) Hart DJ, Magomedov N. *J Am Chem Soc* 2001;123:5892–5899. [PubMed: 11414821] (c) Hart DJ, Magomedov N. *Tetrahedron Lett* 1999;40:5429–5432. (d) Chen Z, Fan J, Kende AS. *J Org Chem* 2004;69:79–85. [PubMed: 14703381]
6. (a) Schollkopf U, Hartwig W, Groth U. *Angew Chem Int Ed Engl* 1979;18:863–864. (b) Sinclair PJ, Zhai D, Reibenspies J, Williams RM. *J Am Chem Soc* 1986;108:1103–1104. (c) Orena M, Porzi G, Sandri S. *J Org Chem* 1992;57:6532–6536. (d) Genet JP, Feroud D, Judge S, Montes JR. *Tetrahedron Lett* 1986;27:4573–4576.
7. (a) Dunn MJ, Jackson RFW, Pietruszka J, Turner D. *J Org Chem* 1995;60:2210–2215. (b) Reißer M, Maas G. *Synthesis* 1998:1129–1132. Related work on propargylglycine: (c) van Esseveldt BCJ, van Delft FL, de Gelder R, Rutjes FPJT. *Org Lett* 2003;5:1717–1720. [PubMed: 12735760]
8. Jackson RFW, Perez-Gonzalez M. *Org Synth* 2005;81:77–88.
9. Piers E, Wong T, Howard EM, Chong JM. *Can J Chem* 1987;65:78–87.
10. Piers E, Wong T, Ellis KA. *Can J Chem* 1992;70:2058–2064.
11. Leusink AJ, Budding HA, Marsam JW. *J Organometal Chem* 1967;9:285–294.
12. The N-*t*-Boc residue was first removed from compound **6** by treatment with Et₂O·BF₃ in EtOAc. The free amine thus obtained was then coupled with Mosher's acid chloride to yield the corresponding Mosher's amide. See Mosher HS, Dale JA, Dull DL. *J Org Chem* 1969;34:2543–2549.
13. Farina, V.; Krishnamurthy, V.; Scott, WJ. *The Stille Reaction*. John Wiley & Sons; New York: 1998.
14. Scott W, Crisp GGT, Stille JK. *Org Synth* 1989;68:116–129.
15. Farina V, Krishna B, Marshall DR, Roth GPJ. *J Org Chem* 1993;58:5434.
16. Racemic products were prepared and separated on chiral HPLC. Conditions: Chiralcel OD (Column No. OD00CE-DL010), Chiral Technologies, Inc. Eluent: Hexanes/IPA (90/10); Flow Rate: 1.0 mL/min; Detection: UV270. For all products listed in Table 1, >99:1 *er* values were obtained.
17. Fauvarque JF, Pflüger f, Troupel M. *J Organometal Chem* 1981;208:419–427.
18. Ugo R, Pasini A, Cenini S. *J Am Chem Soc* 1972;94:7364–7370.
19. Zhang HX, Guibe F, Balavoine G. *J Org Chem* 1990;55:1857–1867.
20. Leroy J. *Synth Comm* 1992;22:567–572.
21. Shih C, Fritzen EL, Swenton JS. *J Org Chem* 1980;45:4462–4471.
22. Boukouvalas J, Lachance N. *Synlett* 1998;1:31–32.

Figure 1. Structures of spiroquinazoline, alantrypinone and γ,δ -unsaturated amino acid derivatives **2**.

Figure 2.
E-Stereochemical assignment for **6**.

Scheme 1.
Preparation of stannane **6**.

Scheme 2.

Scheme 3.
Synthesis of **10** via **9**.

Table 1

Stille coupling reactions of stannane **6**.

Entry	R-X	conditions	Yield %	R
a		Pd_2dba_3 , CuI , AsPh_3 , DMF	68	
b		Pd_2Cl_2 $(\text{CH}_3\text{CN})_2$, Bu_3SnH	79 (brsm)	
c		Pd_2dba_3 , CuI , AsPh_3 , DMF	71	
d		Pd_2dba_3 , CuI , AsPh_3 , DMF	77	
e		Pd_2dba_3 , CuI , AsPh_3 , DMF	57	
f		Pd_2dba_3 , CuI , AsPh_3 , DMF	80	
g		Pd_2dba_3 , CuI , AsPh_3 , DMF	84	
h		Pd_2Cl_2 $(\text{CH}_3\text{CN})_2$, Bu_3SnH , 45°C	67	

Table 2Stille coupling reactions of stannane **7a**.

Entry	R-X	conditions	Yield %	R
a		Pd ₂ dba ₃ , CuI, AsPh ₃ , DMF	74	
b		Pd ₂ dba ₃ , CuI, AsPh ₃ , DMF	79	
c		Pd ₂ dba ₃ , CuI, AsPh ₃ , DMF	72	
d		Pd ₂ dba ₃ , CuI, AsPh ₃ , DMF	69	
e		Pd ₂ dba ₃ , CuI, AsPh ₃ , DMF	84	
f		Pd ₂ dba ₃ , CuI, AsPh ₃ , DMF	76	