THE SOLDIER'S HEART.*

BY

SIR JAMES MACKENZIE, M.D., F.R.S.,

LECTURER ON CARDIAC RESEARCH AT THE LONDON HOSPITAL.

THAT the trying life of soldiers, exposed to many and varied vicissitudes, should put a strain upon the heart is what we all recognize. This strain inevitably finds out the impaired hearts, whether the impairment gives rise to physical signs of lesions or not. This is well understood, and, in treatment, presents no feature which has not been recognized in civil life, and about this class of case I do not intend to say anything.

These cases, however, form but a small percentage of the cases that are actually invalidated on account of their hearts. It has long been recognized, particularly since the instructive account published by physicians during and after the American Civil War, that there is a form of heart trouble to which soldiers are particularly liable, and which has gone under the name of the "soldier's heart" or the "irritable heart of soldiers."

Shortly after the outbreak of this war soldiers invalidated on account of heart trouble began to appear in my consulting-room. These were few at first, but as time went on they came in increasing numbers, and I took great pains to find out the circumstances and signs connected with their complaints. These observations brought to light what seemed to me the main cause of the illness which affected the soldiers, so that when I was asked to take charge of the inquiry into disorders of the heart for the medical history of the war, I had before me an object which would give our researches a very profitable aim.

In this work I have been associated with Dr. Wilson, and we have examined about 400 cases. In at least 90 per cent. of the cases that have been certified and treated as heart affections we find that they are not primarily heart cases, and that the principles of treatment applicable to these cases are widely different from those who suffer from heart failure.

SIGNS AND SYMPTOMS.

When we see the soldiers in this country invalidated because of heart trouble we find a good deal of variation in their appearance and symptoms. The face is often lined and drawn; many are spare and thin with a great vasomotor instability, as shown by the manner in which the peripheral circulation varies, the hands and fingers at times going pale and cold, at other times the fingers are thick and red, and the nose likewise becomes red and even blue with slight exposure to cold. If they have been treated for some months by rest and feeding, some become pale, fat, and scant of breath.

The chief complaint is an absence of the feeling of being well; they often feel out of sorts—"rotten" is a term frequently employed. A sense of fatigue or exhaustion easily induced is common to all. Breathlessness on moderate exertion is frequent, and more infrequent is pain over the region of the heart. The physical signs are variable. The heart's rate is often not increased; in some it is persistently increased, as frequent as 120 per minute. More frequently, at rest the rate may be quite moderate, but exertion, sometimes slight, may produce an undue rapidity, and it is in consequence of this excitability that the term "irritable heart" has been used.

Murmurs, systolic in time and heard in different regions, are frequent, while an increase in size, usually slight, is not uncommon. In a few cases there is a slight oedema of the legs.

The mental condition is somewhat varied. Periods of depression are not infrequent, and the patients are often very irritable. They accept the view that they have something wrong with their hearts and readily yield to all restrictions, and are often content to lie in bed and brood over their woes.

These are the salient features in a great majority of cases, and you will recognize them as common in civil life. We find identical conditions, for instance, in people recovering from an exhausting illness, as typhoid fever or influenza, or after a severe surgical operation. We

see all these phenomena well marked in people who are suffering from some infective organism, and we also find them in people who have suffered a long mental and physical strain, particularly with insufficient sleep, as in a daughter who has for long periods nursed an ailing mother.

The recognition of the character of these complaints gives us a guide in seeking for the cause of the exhaustion in these invalidated soldiers, and in a great majority of cases we get a history pointing to an illness which preceded their breakdown.

Onset of Symptoms.

The account given of the onset of symptoms is peculiarly instructive. They will say they were in the trenches and felt well and fit until one day they felt seedy and ill, and this continued until they were compelled to seek medical advice, when they were found to have a raised temperature. A few days' relief was obtained, and they returned to their work still feeling far from well, and, after a few weeks of the strenuous life in the trenches, they collapsed, sometimes with loss of consciousness, breathlessness, or even pain.

The story of the onset varies. Sometimes it is an attack of diarrhoea, which persists for a time; sometimes it is after a definite illness, as measles, but most give a history which we can safely surmise as being due to an infection. In a few cases one cannot get such a definite account of the starting of the illness, but many do recognize the gradual onset of the trouble. That is the story of the majority of cases, but there are a number from whom we can get no suspicious history of infection, but where there is an account of a very strenuous life.

To grasp fully how this exhaustion is brought about, we must understand the life in the trenches. The story of some of these soldiers is illuminating. One in December, 1914, suffered from appendicitis, and was operated on. He returned to duty three months after, and, in June, went to the front. Immediately on arrival he went into the trenches and was there a fortnight. Every night was spent in repairing the damaged parapets. They were constantly being shelled. He never slept at night, and occasionally got a few hours' sleep in the day, often being twenty-four hours without sleep. One day a shell exploded in the trench, knocking him partly unconscious. On regaining consciousness he stuck to his work for twenty-four hours, but had to give in, feeling weak and ill with pain over the region of the heart.

Effects of Strain during Infection.

So great is the mental strain and bodily exertion, with sleepless days and nights, while the trenches are frequently bombarded, that one might be disposed to consider that these factors would be sufficient to account for all the symptoms. But we find identical symptoms present in many who have never been to the front, and who have had no excessive bodily or mental strain, but who have suffered from some febrile infection. The majority also, as I have said, give a history of some previous illness, and it is necessary we should recognize the effects of strain or effort complicated by febrile illness. For that purpose I recite to you the following experience:

A young officer was training for a foot race, and every day spent some time running round a track. One day, while doing this, he collapsed, fell, and was picked up partly unconscious. A doctor was summoned, who said he had strained and dilated his heart, and that was the cause of his collapse. I was asked to see him, and, after getting this account, I recognized that a man who had taken a considerable amount of exercise the day before could not possibly suffer from heart failure, so I asked him if he felt quite well before he began to run, and he admitted that he felt rather seedy, but hoped to shake off the feeling with the exercise. The temperature was taken and found to be raised a few degrees, so I had no hesitation in stating that the cause of collapse was the toxic influence of some microbe invasion and that, with the subsidence of the fever and a few days' rest, he would be all right. That is what happened, and he was able to resume his training, and ultimately ran his race in perfect health.

The action of toxins produced in the body by bacterial invasions has not yet been clearly worked out, but so far as I have gone into the matter a clear and distinct group

* A paper read in opening a discussion in the Section of Therapeutics of the Royal Society of Medicine on January 18th.

of illnesses can be attributed to the toxins even when the organisms themselves do little damage. The phenomena I have described as characteristic of the soldier's heart will give you an idea of the point I want to establish—that the condition is one of general exhaustion, and the circulatory symptoms are but parts of a general manifestation. To appreciate this more clearly so that we may separate them from those in which there is an actual lesion of the heart, I will call attention to some striking differences between the symptoms of the poisoned heart and the symptoms of the infected heart (that is, where the organisms have invaded the heart) and of chronic disease of the heart.

1. *From Infected Heart.*—In some cases there may be doubt as to whether the condition may not be due to an infection of the heart itself. It may not be possible to settle the question while there is a rise of temperature, so judgement will have to be deferred until the fever subsides. If the temperature remains raised for a period of weeks, and no source of infection can be found, then the suspicion that there is an affection of the heart becomes strengthened. With the permanent fall of temperature, the heart can usually be assumed to have escaped infection and the abnormal manifestations (increased rate, systolic murmur, and increase in size) may be taken as the reaction of the heart to a toxin, or to the factors that produce a general exhaustion.

2. *From Chronic Heart Disease.*—The first sign of a failing heart affected by some old-standing lesion is invariably a subjective one—due to a distressful sensation arising when some effort is made which the individual used to perform with no distress. By far the most common sensation occasioned in this way is breathlessness, the next most frequent, pain. Both these may be associated with palpitation. These are sensations which everyone recognizes as indications of inefficiency of the heart. In the poisoned heart these are usually not the first signs, though they also appear—the first sign being the sense of exhaustion.

The Sense of Exhaustion.

This sensation is often taken as a sign of cardiac inefficiency, but it owes its causation to a different mechanism.

When people suffering from weakness consequent upon an infection, or after an exhausting illness, attempt to walk on the level until they are conscious of their weakness, the sensation by which they perceive their weakness is one of exhaustion or sometimes of giddiness. If they stop, sit, or lie down the sensation speedily disappears. If they attempt a more violent form of effort, as walking quickly or up a hill, or running upstairs, they may be pulled up by breathlessness or palpitation—sensations provoked by heart weakness. It is to the sense of exhaustion and its allied sensations, giddiness and faintness, that I wish to call attention, because the mechanism of production is different from that producing the sensation in patients with primary heart failure.

In people suffering from some toxic influence the central nervous system is always affected. This is shown by the sense of feeling ill, by depression, and irritability of temper. There is also evidence of vasomotor disturbances, as shown by susceptibility of the peripheral circulation—the hands and feet persistently cold, or made so by excitement. There is a persistent over-action of vasomotor influences, as the sense of chilliness that persists after a cold bath; at other times flushes of heat pass over the body, and warmth may tend to over-filling of the peripheral vessels. This is seen in people who faint when standing in a warm room, or who speedily become exhausted or even faint when exertion is made. In these the blood tends to accumulate in the peripheral veins of the limbs and in the large abdominal veins, with consequent anaemia of the brain. It is because of this anaemia of the brain that the sense of exhaustion and syncope is provoked.

I do not enter here into all the facts which support this explanation of the sense of exhaustion, as many observers beside myself have fully dealt with the subject. My purpose here in drawing attention to it is to emphasize the mechanism of its production and to show that it is produced in a different manner from the signs of cardiac exhaustion—breathlessness, palpitation, and pain.

The Character and Meaning of Cardiac Pain.

Another distinctive feature which is occasionally present is a sense of discomfort, amounting sometimes to pain even of a severe kind, felt over the region of the heart. The pain may be occasionally provoked by exertion, but more frequently it comes on when the individual is at rest. Its occurrence, not in response to effort, is characteristic of pain due to a poisoned heart, for angina pectoris due to disease in the young occurs when effort is made. While not absolutely diagnostic, the occurrence of pain, when this type of patient is at rest, is strongly suggestive of poisoning.

Heart Abnormalities are but a Feature in a General Condition.

Taking into consideration all the facts, it will be found that the condition from which certain of these soldiers suffer, who are usually understood to have acquired a heart affection, is not, properly speaking, cardiac in origin, but is the outcome of an injury to other systems as well as the heart, such as the central nervous system. Even when we find such marked abnormalities as increased rate, systolic murmur, and an increase in the size of the heart, the cause of these signs ought not to be looked upon as disease, but merely as part of the manifestation of general illness. The importance of this point of view will be realized when we consider the treatment.

THE CAUSE OF ILLNESS.

While the reason for attributing the majority of these cases to bacterial and toxic influences is so far purely clinical, good grounds can also be found from bacteriological evidence. As, however, bacteriology is outside my province, I will not touch on that evidence further than to point out that I have frequently met with these phenomena in civil life in people in whom the causative organism has been detected.

Here, however, I should like to point out that while the active agent producing these phenomena may be an organism or its toxin, or the products of physical exhaustion, some changes are produced in the economy—in the blood, central nervous system, or heart muscle—which impair the functional activity of these structures. I do not pretend to be able to decide which of these conditions is present, and in any case they do not materially affect the principles of treatment which have been found of value. It is possible, however, that where the illness is due to a microbe, vaccine therapy may be of value; but I am not qualified to speak of this.

NOMENCLATURE.

It is difficult to get a term which will convey clearly what the nature of the trouble is, so I use here the simple one of "general exhaustion."

It is a great misfortune that a custom has arisen that each sick soldier has to be certified as suffering from a definite complaint. Naturally, in the hurry and bustle of the earliest examination, the examining doctor has not the time to make a careful study of each individual. On their immediate breakdown the patients often present features of exhaustion in which the cardiac phenomena stand out most prominently, and the doctor who sees them naturally labels them by these phenomena. It is for this reason, no doubt, that so many of them are invalided on account of "valvular disease of the heart," "disordered action of the heart," or "dilatation."

My description, like my nomenclature, is very general, and as we inquire deeper into the subject we may be able to distinguish separate groups, each with a definite cause.

TREATMENT.

I have dealt at some length with the diagnosis of these cases, because if we recognize the nature of the illness, then we can readily see on what principles treatment should be based. The lines are those we employ in civil practice. If the patient be the host of some microbe, the treatment should be devoted to increasing his power of resistance. If he be poisoned, treatment should be devoted to the elimination of the poison. Should certain tissues be injured by the invading microbe, or by the toxin or by physical strain, then treatment should be devoted to their healthy renewal.

These indications are practically those which should

guide us in treating individuals after an exhausting illness. We have, however, in addition, to consider the mental condition of the soldier whose experiences have depressed his mental faculties, and who has been told he has some heart trouble. This knowledge always tends to depress, and to make them extremely amenable to that form of treatment that is the worst for them—rest in bed or the avoidance of exertion. If we recognize that such cardiac phenomena as murmurs, increase in size, and variability of rate are but the experiences of an irritated heart, we shall see that too much rest is not beneficial. Every organ benefits by the judicious exercise of its functions, and the heart can also be beneficially stimulated.

The principles of treatment, then, should be devoted to increasing the health of the body as a whole in such a way as to increase the natural resistance to infection, to eliminate toxic influences, and brace up the whole man bodily and mentally. The best way to achieve this is by fresh air and judicious exercise in the fresh air. I need not labour the fresh air idea as it is evident to all, but I want to emphasize the question of exercise.

There is a notion deeply rooted that if there is anything wrong with the heart the patient must be restricted in the amount of effort, and medical men who are themselves afflicted are quite content to lie in bed or move gingerly about.

The line I have adopted with those soldiers who have consulted me is that which I have adopted in civil life. All those I have seen have already been told that their hearts were affected, and this knowledge tended to depress. They often feel miserable, so that there is a mental side to the case, which is aggravated by the supposition that there is something amiss with the heart. This aspect of the case has to be met, and consequently in our treatment this view has to be kept in mind.

When all fever has subsided, or when there is but an occasional rise of temperature, I encourage the patient to get up and go about. I find out the form of outdoor exercise which gives him most pleasure. In my private practice my patients have been officers, so I recommend them to start as soon as possible any form of sport or game which they can get—fishing, riding, shooting, golf, etc.

In recommending such exercise I have two objects in view. The first is the mental condition of the patient, and I need not insist upon the beneficial effects of occupation. The more interesting that is, the more likely is the patient to be taken out of himself. I have referred to the fact that the sense of exhaustion is the predominant symptom in these cases, and that it is probably vasomotor in origin, due to the irritation of the central nervous system. We know that the mental state may play a part in producing exhaustion, for we find that a boring form of exercise will readily produce it, while a form of exercise that is full of interest can be borne for long periods without any exhaustion. Therefore, this out-of-door occupation should be in an attractive form.

In recommending effort, I know it will be said that there is danger in this, just as there was when the soldier was employed in his military duties, particularly when such physical signs as murmurs and dilatation are present. If the true nature of these signs be recognized—that they are but the expression of a temporary enfeeblement of the heart muscle, and that the same measure for restoring a flabby leg muscle applies to a flabby heart muscle—we shall see the importance of exercise judiciously employed. No doubt injudicious exercise may do harm, but there is one precaution of a very simple kind which suffices to prevent any danger, and that is to instruct the patient to indulge in his exercise so long as it gives him pleasure and causes no distress nor discomfort, but to stop or slow down as soon as he experiences the sense of exhaustion, breathlessness, or pain.

This simple and easily comprehended dictum contains in it the whole philosophy of the therapeutics of exercise as applied to the heart, and may be used with safety in all obscure and doubtful conditions of heart weakness.

It will, however, be said that it is all well and good for officers who can afford such means of pleasurable occupation, but what about the private soldier cooped up in the wards? It is for him I wish to prescribe. I used the officer to illustrate the principle so that I may strongly urge that the same principles should be applied to the

treatment of the private. If the officer is depressed and makes but slow progress in his comfortable quarters, what of the private soldier in the gloomy wards, meditating on his troubles, or grouching with his fellows, with an occasional joy-ride in a motor-car? I want this argument for congenial exercise to be fully considered, not only for the pleasure of the sick man, but so that the soldier may speedily regain his health and return the sooner to his duties.

I know that already in convalescent camps the aim of interesting occupation with fresh air is being carried out, and I recommend the same principle to be applied in the so-called heart cases. But in these exhaustion may arise too soon for the participation in mechanical occupation, and I would recommend that the same sort of mental stimulus which has been found beneficial in the officer should be employed—that is to say, games. Games absorb the attention and infuse a different spirit from any other kind of employment, and a selection of some games, as bowling, or quoits, or skittles, would add materially to our therapeutic resources.

As a practical outcome of these suggestions, I would recommend that the Government should provide a large hospital for the reception of soldiers who are supposed to be suffering from a heart affection; that these principles of treatment I have detailed should be employed for the recognition of the nature of the complaint, and the principles of treatment here recommended should be carried out.

As soon as signs of improvement are shown, the soldier should begin drilling—at first for a short time each day; and, as improvement goes on, the time increased until he was able to undertake a normal amount. In this way his preparation for fitness would begin early, and he would be rendered fit much sooner than under the present method of treatment.

I would also suggest that this view of the nature of the soldier's heart should be communicated to the medical officers, in order that the diagnosis should be in conformity with the actual condition, and in this way prevent the grievous injury that is done to the sick soldier by implanting in his mind the depressing idea that he suffers from an affection of the heart.

THE IRRITABLE HEART OF SOLDIERS.*

BY

R. McN. WILSON, M.B., CH.B. GLASG.

IN the course of my work as assistant to Sir James Mackenzie, in the inquiry conducted for the Medical Research Committee, I have examined upwards of 200 cases of the condition known as "the irritable heart of soldiers." Careful study of these cases has brought me to the same conclusions as those enunciated by Sir James Mackenzie. I have met what I can only regard as a definite clinical picture, sharply defined and differentiated. The cardinal symptoms he has indicated—great exhaustion after effort; breathlessness on slight exertion; a rapid pulse, which becomes much more rapid upon the slightest attempt at action; pain over the precordial region or along the left costal margin; and a vasomotor condition of greater or less stability—have been present in all my series. I have observed also the tendency to attacks of giddiness to which he drew your attention, the nervous symptoms, the generally high blood pressure, and the tendency to persistence of the condition so long as the soldier is kept confined to his bed.

I propose, therefore, to confine myself to a consideration of the history of the condition, and shall include under this heading the various theories regarding causation that have at one time or another been advanced.

In 1864 the British Government appointed a committee to inquire into the heart conditions prevailing in the army. The committee was assembled by Earl De Grey, and consisted of three generals and two doctors. It sat during the years 1864, 1865, 1866, 1867 and 1868, and issued four reports. In the last the question of infantry accoutrements was considered, and a recommendation was made that the

* Read at a meeting of the Section of Therapeutics of the Royal Society of Medicine.