AN ACCOUNT, WITH COMMENTARY,
OF
A CASE OF SPLENECTOMY

IN ADDENBROOKE'S HOSPITAL, CAMBRIDGE,

UNDER THE CARE OF

SIR CLIFFORD ALLBUTT, K.C.B., F.R.S.,

L. HUMPHRY, M.D.,
F.R.C.P.,

F. DEIGHTON, M.B.,
M.R.C.S.,

AND

DOROTHY C. HARE, M.D., B.S.LOND.

SPLENIC ANAEMIA, that is, a progressive "secondary" anaemia with enlargement of the spleen, is probably more common than was supposed before the comparatively recent investigations of Sir W. Osler and others drew the attention of the clinician more definitely to its characters. During the autumn of 1915 three cases of this malady came under the care of one of us (C. A.) in the Addenbrooke's Hospital. (Malaria has been unknown in Cambridgeshire for more than forty years.) One of these cases was seen once or twice only and lost sight of; the second was in a little boy (aged 6 or 7) who presented all the now well-known features of the disease. After long consideration it was decided to remove the spleen, an operation to which the mother gave her consent; unluckily however the boy got to hear of our intention and howled until the mother's resolution, not unnaturally, gave way, and he was removed. The case was a severe one; haemorrhages, intestinal and other, were frequent, and probably the boy is no longer living. In this case Dr. Dorothy Hare and Dr. Humphry took much pains to perfect the method of testing for corpuscular fragility, and Dr. Hare, on repeated examinations, found the fragility normal. There was no other point in it of sufficient importance to detain us. The third case was as follows:

Mrs. K., aged 45, was admitted under the care of Professor Allbutt on October 25th, 1915. She was extremely prostrate, blanched, and short of breath. The colour of the face—save some venous injection on the malar bones—and of the whole body was that of an old tallow candle. No discoloration could be found on the skin; indeed the axillary and other folds of the body were, like the skin elsewhere, unusually pallid. No notable emaciation. The mucous membranes were almost as pale as the skin. The teeth were ill kept, two of them loose and protruding. These teeth were drawn without any excessive bleeding; the rest, in consideration of the woman's state, were not extracted, but the mouth was thoroughly cleansed and the gums dressed daily with iodine. There was no definite pyrexia. The tongue was pallid, but quite normal in appearance and soft in its substance. The

spleen was moderately enlarged, fairly soft, and a little tender; the notch distinct. No sign anywhere of enlarged lymphatic glands. Excepting haemic murmurs and a very feeble pulse nothing of importance was noted on the side of the heart and blood vessels. The ankles were a little puffy. The liver was apparently normal, but the edge just perceptible below the ribs. No signs and no history of haemorrhages; but occasional passing attacks of diarrhoea or vomiting. From the time of admission onwards the charts showed a continuous pyrexia, as recorded in the annexed reprints of them.

As the woman grew steadily worse, and the occurrence of haemorrhages was feared, it was decided, after careful consultation with colleagues, and especially with Mr. Deighton, to remove the spleen. The patient at last gave her consent, and on December 3rd the operation, to be described by Mr. Deighton, was performed.

Operation.—The abdomen was opened by a vertical incision through the upper part of the left rectus; a few adhesions were met with. The spleen was then gently brought into the wound, the lienorenal ligaments and gastro-splenic omentum were tied and divided. The pedicle was then transfixed and ligatured in three portions with interlocking ligatures; there was hardly any haemorrhage. The wound was then closed.—(F. D.)

The spleen weighed 1½ lb. Histological examination showed nothing abnormal.

The patient then came under Dr. Humphry's charge, and for a few days after the operation suffered a good deal. So severe an operation was a great trial to a woman disheartened, anxious, and very feeble. She suffered also for some days rather severely from the after-effects of the anaesthetic; the stomach was upset, and food ill tolerated. Not only so, but, as recorded on the charts, the pyrexia, after a promising fall towards normal, became still sharper. Evidently, as the appended blood counts will show, a great commotion was set up in the blood factories. However, on December 28th—as seen in the charts—the pyrexia began to fall, and on January 5th was settling down when another sudden rise took place, coincident with an accumulation in the colon. Notwithstanding, during these weeks the patient was obviously rallying; a little bloom of colour and a little ray of hope brightened her face, and she began in small ways to help herself. And happily this rise of temperature proved to be the last; thenceforward the fever ceased, and on January 24th she was discharged convalescent. For a week she had been sitting up and walking about the ward. Her appearance, if still somewhat anaemic, had greatly improved, and she felt able to return to her home duties.

On the blood counts taken at intervals during her stay in the hospital we need not dwell, as the appended records—all taken by Dr. Dorothy Hare—speak for themselves. The variations are closely in accordance with those displayed in the other published cases. There is an excellent tabulated collection of blood counts in splenic anaemia

Table showing Results of Blood Examinations.

Date.	Haemoglobin.	Red Cell Count per c.mm.	Nucleated Reds. Counted per 100 Leucocytes	Total Leucocytes per c.mm.	Differential Leucocyte Counts.						Remarks.
					Poly-morpho-nuclears.	Lymphocytes, Small and Medium.	Mono-nuclears, Large.	Hyalines.	Coarsely Granular Eosinophiles.	Basophiles.	
1915.	Per cent.				Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	
Oct. 22	33	2,980,000	0	4,500	72	23	—	2	—	3	Slight poikilocytosis.
Nov. 27	35	2,110,000	0	33,000	—	—	—	—	—	—	Period of irregular pyrexia from Nov. 25 to Dec. 3.
Nov. 30	—	2,630,000	—	11,000	—	—	—	—	—	—	
Dec. 13	36	4,003,000	2	36,000	77	—	23	—	—	—	Dec. 3 operation. Count 10 days after operation. The lymphocytes were chiefly large cells, not distinguishable from atypical hyalines.
Dec. 18	42	3,850,000	3.5	22,600	69	21	6	3.5	—	0.5	15 days after operation.
Dec. 29	40	4,270,000	1.5	17,200	57	26	9	5.5	2	0.5	26 days after operation.
1916.											
Jan. 8	42	3,480,000	2	27,600	68	27	3	1.5	0.5	—	33 days after operation. Pyrexia during three previous days.
Jan. 19	44	4,010,000	0	10,000	28.5	55	9	4	2	1.5	47 days after operation.
Jan. 23	—	—	—	—	33	47	13	7	—	—	51 days after operation.

in Thursfield and Gow's paper in *St. Bartholomew's Hospital Reports*, Vol. I, part i.

As our paper is entirely clinical we need not enter into the obscure problems of splenic physiology. Suffice it to say—strange as it may seem—that this large organ, presumably with a function, or functions, so ample, may nevertheless be excised without ultimate, or indeed proximate, impairment of life. After some temporary commotion the body seems to settle down again to work without sensible shortening of its vital capacity. We have not yet data, it is true, to assure us that after splenectomy life may be extended to fullness of years, though some such patients are now under observation whose lives have been prolonged in apparent health for such periods as eight and ten years after the operation; two of Osler's patients have survived for nine and fourteen years (Ede). How this remarkable compensation comes about is variously explained; it is enough for the present to say that our experience is now sufficient to justify us in excising a diseased spleen if symptoms seem to require it.

Dr. Cutlbert Ede (in his M.D. thesis, Cambridge, December 10th, 1915) formulated the following conclusions:

1. No ill results follow removal of the organ; that is, of course, assuming that the operation is performed successfully and under favourable conditions.
2. The first and best marked change in the peripheral blood is an increase in the total number of the polymorphonuclear cells, and a large increase in the number of the red cells. This is instantaneous.
3. During the first year after the operation there is an increase in the absolute number of the lymphocytes.
4. Later these become normal in number, but eosinophiles are increased.
5. Finally the blood picture resumes a perfectly normal state.

Noguchi, quoted by Dr. Ede, comes to the same conclusions, and they are confirmed by our patient's case, and by Captain Jones's case in this JOURNAL, February 5th, 1916. Dr. Ede and others have suggested that in these cases of splenic anaemia the fragility of the reds is increased; to this point Dr. Humphry and Dr. Dorothy Hare gave close attention and care, but reported that in both the cases here mentioned (the boy and the woman) the fragility was not increased—was, in fact, normal.

Another curious point in recovery from splenectomy is that thereby the protective virtues of the body against infection are in no way impaired. (See Fuhs,⁶ and our own case, January 4th to 7th.)

Osler and Rolleston have done much to elucidate and differentiate diseases of the spleen. Splenic anaemia they describe as follows (the quotation is a little abbreviated):

1. Diminution of the number of reds with a falling haemoglobin value per corpuscle.
2. No leucocytosis, usually leucopenia.
3. Enlargement of the spleen, not correlated with any other known splenic disease.
4. The tendency to gastro-intestinal haemorrhage.
5. The termination, if sufficiently prolonged, in hepatic cirrhosis with ascites (Banti's disease).

We are indebted to Dr. Ede for references to published results of operation (excision)—namely, Johnston,⁷ Redman and Willard,⁸ and Sidney Boyd.⁹ The issue of the present series of operations is to show that the advantage of operating in an early and uncomplicated stage of the disease is enormous; the mortality is thus reduced in cases of this class from about 50 to about 10 per cent. Banti's figures are to the same effect. The most perilous event of the operation is that of intestinal haemorrhage, due perhaps to thrombosis of the splenic and other larger veins; but in cases operated upon betimes and under favourable conditions this event also is far less frequent.

It will be seen that our case, both in respect of its issue and of the behaviour of the corpuscles, was conformable with all the chief points of Dr. Ede's experience and that of the authors quoted by him.

NOTES ON TABLE (D. C. H.).

The most interesting feature of the chart is the record of the behaviour of the polymorphonuclears and lymphocytes. Even before the operation their numbers varied widely. The first count showed the leucopenia usually associated with the disease, but later there was a high leucocytosis, coincident with a period of increased irregularity of temperature.

After the operation the variations in the chart followed in the main the usual course. There was an early increase in the polymorphonuclears, succeeded in about six weeks by a sharp rise in the lymphocytes. A high polymorphonuclear count in the fifth week followed a few days of high temperature (January 4th to 7th), which supports the observation of Fuhs that removal of the spleen does not abate the capacity for reaction against infection.

The classification of the lymphocytes and hyalines in the early days after the operation presented some difficulty, and in the first count the large mononuclears are grouped together. Later the types became more distinct, but there still remained at the last examination a large number of these cells intermediate in type.

Before the operation the test for the fragility of the red corpuscles was carried out by Thursfield's method with unwashed corpuscles and gave a normal result. The urine was also tested for the presence of haemolysins and

a positive reaction obtained. The method used was that recently published by McKee (in the JOURNAL, October 23rd, 1915).

P.S.—On February 28th Mrs. K. presented herself again at the hospital for examination. She seemed quite well, and said she felt well. She had done all her housework almost from the date of leaving the hospital. Her tongue, pulse, temperature were all normal; appetite good, bowels regular. Complexion now well coloured, and all the tallowy tint had disappeared. She was strongly urged now to have her mouth attended to and all bad teeth removed from unhealthy gums. Dr. Hare made a fresh blood examination, which is here appended.—C. A.

Blood Examination, February 28th, 1916.

Haemoglobin	...	...	48 per cent.
Red blood cells	...	...	3,950,000 per c.mm.
White blood cells	...	...	41,000 per c.mm.
Differential leucocyte count:			
Polymorphonuclears	...	...	48 per cent.
Lymphocytes small and medium...	...	...	44 per cent.
Large mononuclears	...	...	5 per cent.
Hyalines...	...	...	1 per cent.
Basophiles	...	...	2 per cent.
Nucleated red cells,	1 per cent.	of leucocytes counted.	

REFERENCES (collected by Dr. Ede).

- ¹ Hutchinson, *Proc. Roy. Soc. Med.*, 1913, vi, Surg. Sec., 241. ² Faltin, *Deutsch. Zeit. Chir.*, cx, 160. ³ Noguchi, *Berl. klin. Woch.*, 1912, ii, 1839. ⁴ Thursfield and Gow, *St. Bartholomew's Hosp. Reports*, Vol. I, Part 1. ⁵ Fuhs, *Amer. Journ. Med. Sci.*, 1911 cxlii, 713. ⁶ Rolleston, *Clin. Journ.*, xix, 403. ⁷ Johnston, *Ann. Surg.*, 1908, xlviii, 50. ⁸ Rodmand and Willard, *Ann. Surg.*, 1913, lviii, 601. ⁹ Boyd, *Clin. Journ.*, April 1915. ¹⁰ Banti, *Fol. Haemat.*, 1910, x, 33. ¹¹ Otto, *Wien. klin. Woch.*, 1910, 54. ¹² Brill and Mandelbaum, *Amer. Journ. Med. Sci.*, 1913, cxlvi, 863. ¹³ Roth and Bernstein, *Arch. of Ped.*, 1914, xxxi, 340. ¹⁴ Chaurfard, *Presse med.*, 1907, xv, 345. ¹⁵ Thursfield, *Proc. Roy. Soc. Med.*, 1914, vii, Clin. Sec., 84. ¹⁶ Muhsam, *Deutsch. med. Woch.*, 1914, 377. ¹⁷ Ziegler, *Deutsch. med. Woch.*, 1910, ii, 2410. ¹⁸ Seefisch, *Sem. med.*, 1914, 35. ¹⁹ Weber, *Quart. Journ. Med. Sci.*, 1908, iii, 85. ²⁰ Kidd, *Proc. Roy. Soc. Med.*, 1913, vi, Surg. Sec., 231.

THE PRINCIPLES OF TREATMENT AND THEIR APPLICATION TO WOUNDS.

BY

CHARLES F. M. SAINT, M.D., M.S., F.R.C.S.,

CAPTAIN, R.A.M.C.(T.); SURGICAL SPECIALIST, CASUALTY CLEARING STATION, ON ACTIVE SERVICE; ASSISTANT SURGEON, HOSPITAL FOR SICK CHILDREN, AND SURGICAL REGISTRAR, ROYAL VICTORIA INFIRMARY, NEWCASTLE-ON-TYNE.

A WOUND is the local effect produced by an irritant on the tissues, and the consideration of irritation should include (1) its cause, (2) its effects, (3) the reaction to it, and (4) its complications.

1. *The cause of irritation* is described as an irritant. Irritants may be classified as (a) non-infective, and (b) infective; in the former class are included mechanical, chemical, electrical, and thermal agents, x rays, etc.; the infective are micro-organisms.

2. *The effects of irritation* are essentially destructive in nature, and they are general (shock) and local (tissue destruction). Whereas one has been accustomed to regard a wound as entailing a breach of surface, ordinarily produced by a mechanical agent, it is more in accordance with reality to include under the term all destruction of tissue, of whatever degree, in whatever situation, and by whatever variety of irritant brought about—for example, contusions, ruptures of viscera, fractures, burns, and injuries of all kinds.

3. *The reaction of the tissues to irritation* consists of vascular changes and cell proliferation (Rutherford Morison), vascular changes predominating in the acute and cell proliferation in the chronic form.

4. *The complications of irritation* result from (a) extension of the process of irritation; (b) superadded irritation; or (c) healing.

PRINCIPLES OF TREATMENT.

The principles of treatment are: (1) Remove the cause; (2) combat the effects; (3) assist the reaction; (4) prevent complications, or deal with them if they have arisen.

1. Removal of the Cause.

A sterile metallic body gives rise to a minimum of irritation sufficient for the production of a fibrous capsule, whereby the body is completely isolated. In a large proportion of cases missiles have come to be regarded as

sterile, particularly rifle bullets presenting a small wound of entry. Under special circumstances, even where the bullet is apparently sterile, it may require removal—for example, when from its position it gives rise to irritation of nerves, interferes with the movements of a joint, etc. Operative measures for the removal of metallic bodies ought to be restricted where x rays are not available. It is unjustifiable, then, to do any extensive dissection, and difficult operations should be postponed till the base is reached. The commonest and most important indication for the removal of bullets, etc., is sepsis in the track. Should operative measures be determined upon, it is essential to bear in mind that too small an incision increases shock from the excessive retraction demanded, and the longer time necessary for the operation.

2. Combating the Effects.

There is general agreement that in the present campaign the degree of shock is greater than might have been anticipated from the nature of the injury, and there seems to be little doubt that it is due especially to excessive mental and nervous strain; in prolonged engagements, also, insufficient food, arising from the difficulty of its transfer, contributes in no small measure. Rough transport following the receipt of the wound, especially in motor ambulances over paved roads, is responsible for a considerable increase in the degree of its severity.

The requirements of the milder cases are met by a good meal, with some tea or cocoa, immediately followed by quiet and sleep. In the more severe cases, rest, with the foot of the stretcher elevated, is essential. Should pain or restlessness be present, morphine must be administered hypodermically. It is remarkable what tolerance to the drug these patients exhibit, half a grain being a regular and not infrequently inadequate dose; it is of advantage as a prophylactic before transporting the patient from one place to another. Stimulants, whether by the mouth or hypodermically, must be given with caution, the rule being "repeated small doses." For example, one tenth of a grain of strychnine will increase the shock (Crile), while one-fifth of that amount given perhaps each hour will benefit the condition. Should the temperature, as is usual, be subnormal, no time must be lost in raising it by the careful application of heat, internally and externally. Lastly, in the worst cases, the chief stand-by is transfusion in one or other of its various forms. The use of saline is the commonest method, and it may be given per rectum, subcutaneously, or intravenously. Of these, the first two are preferable for ordinary use, since absorption takes place according to demand, while the rectal method has this commendation—that sepsis is almost impossible. With most people the intermittent administration of saline per rectum is more successful than the continuous, and it should be given, according to the tolerance of the individual, six to ten ounces every three or four hours. Half an ounce of brandy may be added to the injections, making an average of three ounces for an adult male in twenty-four hours. Auto-transfusion, though useful, is much less in vogue; rectal injection of one pint of hot coffee and half an ounce of sugar is an excellent substitute.

In the treatment of the local condition wounded blood vessels must be dealt with first, as haemorrhage predisposes to both shock and sepsis. Temporarily any of the first aid means may be utilized, while for permanent arrest the method must be chosen according to the case—for example, pad and pressure, forcipressure, ligature, suture or vessel suture, the last being reserved for vessels of some size. The effects of haemorrhage are counteracted ideally by blood transfusion, but, more commonly, saline transfusion, in one or other way already mentioned, is the means resorted to.

The importance of ascertaining that no tourniquet is left on a limb too long (anything above two hours is dangerous) cannot be too strongly emphasized.

3. Assisting the Reaction.

The reaction of the tissues, consisting of vascular changes and cell proliferation, is usually directed against both cause and effects. When dealing with the cause it is called "inflammation," and when dealing with the effects, "repair," but it is essentially similar.

Where open wounds are aseptic the first essential is to close them, but in war wounds do not often fulfil the