Notes on Military Orthopaedics.

III. THE SOLDIER'S FOOT AND THE TREATMENT OF COMMON DEFORMITIES OF THE FOOT.

BY

LIEUTENANT-COLONEL ROBERT JONES, CH.M.,

F.R.C.S.I. (HON.),

INSPECTOR OF MILITARY ORTHOPAEDICS, ARMY MEDICAL SERVICE.

PART I.—FLAT-FOOT.

THE foot must be regarded for our present purpose as consisting of two parts: The hinder part, behind the mid-tarsal joint formed by the os calcis and astragalus; and the fore part, in front of that joint, consisting of the rest of the tarsus, the metatarsus and the toes.

The Three Arches of the Foot.

The hinder part of the foot is designed to carry the body weight when standing. The os calcis and astragalus are strongly bound together by ligaments, and the astragalus is securely set between the two malleoli at the ankle-joint.

The fore part of the foot is much more mobile, and is designed to give spring to the foot when walking. This spring depends on the efficiency of the small muscles of the foot, on the integrity of the longitudinal and transverse arches, and on the freedom of movement of the toes themselves.

Regarded as a base of support, the foot is a tripod resting on the heel, the head of the first metatarsal bone, and the heads of the fourth and fifth metatarsal bones. Each side of the triangle formed by these three points is arched. The outer side formed by the os calcis, cuboid, and the fourth and fifth metatarsal bones is so slightly arched that practically the whole length of it rests on the ground when standing. This element of the foot is therefore not very susceptible of strain, and need concern us no more.

The inner side is strongly arched; the arch runs from the heel upwards and forwards to the neck of the astragalus, then downwards, forwards, and *inwards* through the scaphoid, internal cuneiform, and first metatarsal bones to the ball of the great toe. This arch, which varies in height in different individuals, is composed of a number of bones, is very susceptible to strain, and is in consequence the seat of a number of disabilities which we shall group together as "flat-foot."

The third side of the triangle runs from the head of the first metatarsal bone to the heads of the fourth and fifth metatarsal bones, and forms the front part of the transverse arch of the foot. It must be remembered that this transverse arch extends backward throughout the whole of the forepart of the foot. The cuneiform bones and cuboid bones are arranged in a transverse arch as well as the metatarsal bones.

Deformities and Disabilities of the Foot.

In considering deformities and disabilities of the foot due value must be given to its various mechanical elements—that is to say, not only to its bony structure but also to the muscles and ligaments which help to maintain its shape.

Looked at in this way we find among the deformities and disabilities of the toes such conditions as hallux rigidus, hallux valgus, and hammer-toe, and we note that the two former in particular are frequently associated with some degree of flat-foot.

Impairment of the transverse arch is specially apt to be associated with the painful condition known as metatarsalgia.

Impairments of the longitudinal arch are associated with a train of symptoms which are found in the various degrees of weak foot, everted foot, and flat-foot.

All these types of derangement of the mechanical elements of the foot and the resulting disabilities depend on strains about the complicated series of joints in the fore part of the foot.

In the hind part of the foot the disabilities are fewer. They are mainly connected with painful conditions of the heel, due to such conditions as strain of the insertion of the tendo Achillis, or periostitis of the os calcis, due to trauma or sepsis.

When a clear idea of these several disabilities and of the measures suitable for the prevention or treatment of each has been obtained, it will be more easy to grasp and understand the various disabilities of the foot to which the soldier is liable, always remembering that two or more types often occur simultaneously, and that all must be appropriately treated.

Flat-foot.

The term "flat-foot" may be taken as a generic term to include all degrees of strain of the longitudinal arch of the foot. To these various conditions different names have been given, which need not detain us here. The point to be understood is that all degrees occur, from slight strain of the ligaments and tendons by which the arch is maintained to complete descent of the arch with osseous deformity, obvious to the most casual observer.

The essential fact in the deformity is abduction and eversion of the fore part of the foot (Fig. 1). If it commences as a sudden acute condition, it is associated with strain of the inferior calcaneo-scapoid ligament; to this is due the tenderness elicited on pressure on this ligament just below the tubercle of the scaphoid. This tenderness is characteristic. Pain in this situation is often most acute in patients who have naturally a very high arch. When such an arch begins to give way the strain on the "spring" ligament is great and the pain correspondingly severe, but the foot is not physically flat, for the arch may still be higher than is normal in most people.

FIG. 1. — Flat everted foot.

As the foot becomes more strained under continued exercise, the bones of the tarsus begin to descend and pain is complained of across the dorsum of the foot.

Finally, when the deformity becomes still greater, there may be pain on the outer side of the os calcis below the tip of the external malleolus. This is probably to be attributed to bruising of the periosteum due to the os calcis impinging on the malleolus; in some cases an adventitious bursa is ultimately produced in this situation.

Acute Flat-foot.

All the above symptoms, and also actual descent of the arch with pronounced abduction and eversion of the foot at the mid-tarsal joint, may come on with great rapidity and then be associated with very severe pain, so that the patient cannot walk and can hardly bear to have the foot touched. Such a condition may fairly be described as acute flat-foot. It occurs in its typical form in people who are in poor physical condition, as, for instance, after an attack of influenza, and return too soon to work which involves much standing and lifting of weights—for example, hospital nurses. It occurs in exactly the same way in the recruit. Take, for example, the clerk who has taken little exercise and who has been in the habit of wearing boots in which there is no room for movement of the toes and bones of the fore part of the foot; all the muscles and ligaments controlling and supporting the arches of the foot are weak and out of condition from want of use; under the strain of hard drill they give way.

Treatment of Acute Flat-foot.

The treatment of such a case falls into three stages.

First Stage.—If the foot is acutely painful the patient should be put to bed and the feet gently massaged for two or three days till the acute tenderness passes off. During this time he must never be allowed to set foot to the ground, for the weight of the body would again strain the ligaments.

Second Stage.—As soon as the acute tenderness has passed off, the foot should be adducted and inverted so as completely to restore the arch, and the foot kept continuously in this position to allow stretched ligaments to shorten and become adapted to the correct position. Mere rest in bed is not enough, for then nothing is done to restore the proper shape of the arches, adaptation takes place in the flat-foot position, and the patient is left with a real flat-foot. It is often best to mould the foot into correct position and fix it in plaster-of-Paris for about ten days to allow the strained ligaments to recover completely.

Third Stage.—When the plaster is removed, after, say, ten days, the patient's foot is exercised—first of all by massage and exercise, particularly movements of the toes and inversion movements, never eversion movements. These inversion movements are intended to strengthen the muscles controlling the arch.

Boots.

The patient is allowed up in specially altered boots. The boots should grip comfortably round the ankle and heel, but the whole fore part of the boot should be roomy to allow free play of the small muscles inside the boot.

FIG. 2.—Boot with straight inner border, crooked and lengthened heel, patch to thicken sole.

Pointed toes are very harmful, as they abduct the fore part of the foot and help to produce the condition of flat-foot. The inner side of the boot should be straight, so that when the two boots are placed side by side the inner sides are parallel right forward to the great toe. This is not the conventional boot of the shoemaker, but it is the correct boot for a strong foot.

The present army boot is not perfect, but it is much better than it was some years ago.

Having seen that the boots are of suitable shape, they should be sent to the shoemaker to have the heels "crooked" on the inner side. By this is meant that the heel of the boot is made a third of an inch higher on the inner side, and tapering gradually to its outer side, so that the inner side of the heel is a third of an inch higher than the outer side. In addition, the length of the heel should be extended forwards by three-quarters of an inch on the inner side, and the sole should be raised one-third of an inch by a patch on the inner side of the sole just below the toe-joint (Fig. 2).

The patient must never put foot to the ground without an altered boot or shoe on, otherwise he will unwittingly let his arch descend and so inflict a fresh strain on the ligaments.

The man should then be instructed to walk with feet parallel, not with the toes turned out. The altered heel helps the patient to keep his toe turned in. Exercises should be graduated till the man is fit for full duty.

A bad case of *acute flat-foot* should be fit for light duty in three weeks, and for full duty with the heels of his boots raised on the inner side in six weeks to two months.

It is really a question whether the army authorities ought not to issue all boots with heels raised on the inner

FIG. 3A.—Thomas's wrench applied to invert everted foot.

FIG. 3B.—Showing the position of the foot and direction of force at end of correction to inversion.

side, as this is a great relief to the foot on a long march and increases the man's weight-carrying and lasting power.

Certainly all recruits with weak feet should be started off with crooked heels on the boots until the muscles of their feet get into training. They should also be taught to walk with their feet parallel. If this were done the feet

would grow stronger rather than weaker, and considerably fewer men would require to be taken off duty.

Traumatic Flat-foot.

Flat-foot in every respect similar to that just described as acute flat-foot may arise from injury. For example, when the wheel of a trap runs over a man's foot and strains all the ligaments.

Flat-foot from Periarthritis.

Similar mechanical conditions arise in gonorrhoeal periarthritis when all the ligaments are sodden with exudate, become soft, and stretch. The same is true when the infection is of an ordinary septic character, for if the man puts weight on his foot before the arch is restored to its strength he will get a flat-foot. He can, however, walk with safety in a boot with the heel well crooked almost as soon as the disappearance of pain allows him to put his foot to the ground.

Rigid Flat-foot.

These cases lead us naturally to the rigid form of flat-foot which results from neglect in the acuter stages, whether the cause of the sudden yielding of the ligaments be overstrain due to unaccustomed exertion in walking, to injury, or to weakening of the ligaments by the products of infective agents.

The foot, therefore, should be maintained in the inverted and adducted position during recovery from the acuter conditions, and when the patient begins to walk the body weight should be deviated from the inner to the outer side (Fig. 2). If this important routine should be neglected the recovery of ligaments will occur in a flat-footed position with stiffness in all the joints. The patient in that case cannot stand on the outer edge of the foot or turn it into an arched shape.

Treatment is very simple, but must be thorough. The foot should be wrenched so as to break down all adhesions and make it thoroughly pliable (Fig. 3). The foot is by this means converted into an acute traumatic flat-foot, and it must undergo the course of treatment already described for that condition. It should be moulded into the correct position, and plaster-of-Paris applied. After two or three weeks of such rest, to allow recovery from the trauma, graduated exercise may be taken in boots so altered as to deviate body weight from the inner to outer side of tarsus. An outside iron is often found a great assistance in maintaining the correct position of the foot during walking (Fig. 4).

Osséous Flat-foot.

A real stiff flat-foot which has been neglected for years becomes an "osseous flat-foot"—that is to say, changes occur in the shapes of the bones to adapt them to the erroneous position. Though something can be done to improve such cases, they cannot always be made fit to be soldiers; the surgeon must, however, be sure that he is dealing with real osseous change and not merely with a flat-foot which is rigid owing to ligamentous contractures and adhesions.

Flat-foot due to Spasm of the Peronei.

I described this condition many years ago (Fig. 5), and stated that it was by no means uncommon. It may be found in any out-patient department if looked for, and I have operated upon as many as 150 cases in one year. It occurs usually after puberty and in early adolescence; it is common among the robust, not only among the weakly, and is not associated with the temperament known as neurotic. I will describe a typical case. A young man

FIG. 4.—Outside iron in addition to crooked long heel and piece to sole.

FIG. 5.—Peroneal spasm producing a flat everted foot.

of 18 limps with a springless gait into the out-patient room. He walks with feet practically rigid and with toes pointing out. They are both everted, and the inner border over the region of the scaphoid appears thickened and even angular. Both the character of the walk and the appearance of the feet might lead to the diagnosis that osseous changes were advanced. On being questioned the patient may give a history of injury such as a fall on the feet; usually he cannot account for the origin of his trouble. The pain is often acute, the feet may perspire abnormally, and the patient may not be able to walk more than a short distance. Relief is experienced on removal of his boots; if he is asked to invert his foot he cannot do it, and when he attempts to do so the peronei become rigid. The surgeon when he examines the foot manually finds there is pain on pressure over the deltoid ligament, over the scaphoid, and over the tip of the external malleolus, where it impinges upon the os calcis, and tenderness over the peronei along the outer border of the foot. If the surgeon gently attempts to invert the ankle much pain is experienced and the peronei, so to speak, immediately place themselves on guard and strongly resist the surgeon's efforts. Now while the surgeon has firm hold of the foot, let him engage the patient in conversation, at the same time gently press in the direction of inversion. Then at the psychological moment, while the peronei are quiescent, very suddenly and very forcibly let the foot be inverted and held there. It is a very painful movement, but the character and contour of the foot is completely changed. It is no longer rigid, osseous changes are obviously absent, and the general appearance of the foot is almost normal. The moment the surgeon releases his hold, the old rigidity and deformity return.

For this condition mechanical measures are of no avail.

FIG. 6.—Exposure of tendons of peronei preparatory to excision of about $\frac{1}{2}$ in.

The patient must be anaesthetized to relax all spasm, and about three-quarters of an inch of each peroneus should be removed about an inch and a half above the malleolus (Fig. 6). The foot must then be fixed for about three weeks, well inverted and adducted at the mid-tarsal joint, and later the ordinary treatment for flat-foot. My old house-surgeon, Mr. Naughten Dunn, secured equally good results by pinching with a pair of forceps the nerve as it enters to supply the peronei. Simple

tenotomy of the peronei is not sufficiently drastic to prevent recurrence of deformity.

Diagnosis.

It is not an easy matter to lay down rules for a hard and fast differential diagnosis between these types of rigid feet, but the following considerations will help.

An osseous flat-foot which has become adapted to its new position is usually strong and painless.

A flat-foot rigid from adhesions and shortened ligaments is still susceptible to strain, and is liable to become painful after unusual exercise.

The type due to a spasmodic contraction of the peronei is characterized by pain, and the spasm may be overcome in the manner I have just indicated.

In the two last varieties, after the deformity has been over-corrected for a time, energetic massage and a correct deviation of body weight are essential elements in bringing about a cure. The osseous type, once the bones have completely dropped, may not only be painless, but may be strong enough to bear the strain of long marches. Wounded soldiers may constantly be met with whose feet are very flat, whose mid-tarsal joints are fixed, but who have undergone several months of hard training without a complaint. If in addition to these osseous changes the power of inversion of the foot is lost or the peronei are in spasm surgical attention is urgently called for.

The flat-foot for clinical purposes may be divided into two classes.

- (a) The foot which the patient can invert.
- (b) The foot which he cannot invert.

The first class will not respond to any mechanical treatment, such as a plate or alteration of the boot. It will not even suffer correction by means of a bandage to

an outside iron from knee to heel. An operation, or forcible manipulation and fixation under an anaesthetic, are essential preliminaries to the simpler mechanical methods.

The second class will never require an operation.

This simple clinical division should be helpful to the military surgeon.

Subacute Flat-foot (Ordinary Flat-foot).

The stage of flat-foot most commonly seen is neither an acutely painful foot too tender to be handled, nor a rigid flat-foot, but an intermediate stage, which may be called subacute.

The patient's feet give him little trouble in ordinary life, but a long walk makes them ache round the instep. When he goes to bed after a hard day's exercise his feet are apt to get stiff. Next morning they are stiff and painful when he gets up, but as he moves about this passes off; but later in the day, especially if he has to do a lot of standing and walking, the feet again become very painful.

Diagnosis.—This is made by considering the patient's history and by finding distinct tenderness on pressure below the tuberosity of the scaphoid; there may or may not be pain in other localities. Finally, the patient can voluntarily stand on the outer edge of his foot and can crook his foot into the arched position. This last point is very important, for it means that the patient will recover if the heels of his boots are crooked so as to deviate body weight on to the outer side of the foot.

The case should therefore at once be put through the last stage of treatment described under acute flat-foot—namely, graduated exercises—and he need not be taken off duty for more than a day or two while his boots are being altered. "Contrast baths" of hot and cold water alternately, to stimulate the circulation of the foot, are a great relief to the patient, especially at night, when his feet may be a little sore and swollen after having been walking about all day.

Conclusion.

I have purposely avoided all mention of tiptoe exercise. These exercises are excellent, but they will not cure a flat-foot if the patient is allowed to come down in the flat-footed position between the period of exercises.

If the principle of deviation of body weight on to the outer edge be loyally adhered to every ordinary case will recover, even without special tiptoe exercise. Loyal adhesion to the principle stated means that the patient never sets foot to the ground except in a properly crooked boot, shoe, or slipper. Even in his bath he must be careful to stand on the outer edge of the foot. The whole idea of treatment is to ensure that uninterrupted adaptive shortening of stretched structures shall take place.

Massage, tiptoe exercises, electric stimulation of muscles all help to hasten the recovery, but a few minutes' careless walking in stocking soles will undo all benefit from other treatment by again stretching recovering structures.

Frequently the high arched foot is the most troublesome type when the tendons are strained and the ligaments begin to give. The strain upon the astragalo-scaphoid joint in this type is, for obvious mechanical reasons, great, and the progress of the affection is often marked by a prominence about the scaphoid due to pressure osteitis.

THE census bureau of the United States Department of Commerce recently issued a summary of its statistics of causes of death in 1914 in the registration area, which comprises 435 cities and towns with populations ranging from 10,000 to 100,000, and represents about two-thirds of the total population of the United States. The total number of deaths was 898,059. Of these, 99,534 were due to diseases of the heart; 96,903 to tuberculosis (of disease of the lung in 84,366 cases); 83,804 to pneumonia; 67,545 to Bright's disease and nephritis; 52,420 to cancer and other malignant tumours; 52,407 to diarrhoea and enteritis; 51,272 to apoplexy; 15,044 to arterial diseases of various kinds; 11,786 to diphtheria; 10,666 to diabetes; 10,185 to typhoid fever; 15,617 to whooping-cough, measles, and scarlet fever. The statistics show that from 1904 there has been a continuous decrease in the mortality from tuberculosis and typhoid fever, while since 1900 there has been an almost continuous rise in that from cancer and diabetes.