Death from sepsis practically never occurs after the end of the second week following the wound. There is no long drawn-out fight with an ultimate and inevitable surrender, such as may be seen at times when the abdomen, the brain, or shattered bones are infected. Out of the last fifteen cases of death among examples that we have seen of haemothorax infected by anaërobe bacilli, fourteen died simply from the sepsis. One, the case related as No. III, died from a late secondary haemorrhage from a deep lung abscess round a retained piece of shell. Pericarditis was proved to be present in eight of the other fourteen fatally septic cases. Jaundice was noticed in seven, of whom two died without pericarditis. There were no instances of death from purulent bronchitis in this series.

None the less purulent capillary bronchitis with thick greenish-yellow airless sputum is a dangerous complication; and it appears to develop with greater severity in infected cases that have been left for many days without drainage. It may be the exacerbation of a cold that was present before the man was wounded; and a history of exposure on the battlefield after the wound is often given by patients suffering from this complication. The bronchi of the uninjured lung are more extensively infected than those on the wounded side, where collapse and immobility tend to restrict the spread of infection along the air tubes. Autopsy reveals thick plugs of green pus rising up from the orifices of every cut bronchiole, and there is sometimes a bronchopneumonic spread of the infection into surrounding lung tissue.

So soon as this character of the sputum is recognized stimulating expectorant medicines should be given freely, for this condition demands a treatment contrary to that usually practised in order to lessen cough in chest wounds. Its presence requires that the haemothorax shall be aspirated or drained away as soon as possible, so as to throw more lung tissue into active respiration. There is no evidence that such treatment introduces further danger by extending the bronchial infection along tubes that were previously collapsed.

Fever that continues for more than a week after rib resection, if not due to purulent bronchitis, is usually caused by inadequate drainage of the empyema. The pleural cavity should be explored by the finger to ascertain whether pus is accumulating in any recess below the level of the drainage tube. If this is found to be the case, a piece should be resected from a lower rib to improve the drainage, or a large catheter should be inserted and suction applied to it by some apparatus, such as the Sprengel pump. Drainage is improved by keeping the patient recumbent in bed.

Sooner or later the lung expands fully in practically all the cases, and the sinus closes without the need for any elaborate surgery. We have learnt the after-histories in England of thirty of our cases, whose chests were drained for infection of a haemothorax, some for anaërobe infections, and others for streptococci, etc. The sinus became closed in the majority during the third or fourth month after the original wound, and some of the men were at once classified as fit for duty. In only eight of the cases was the sinus open as late as the sixth month. There was only one case of chronic empyema cavity in which at the fourth month a minor Estlander operation was performed for its obliteration; and there were two cases in which small operations were needed for the drainage of a loculated collection of pus.

SUMMARY.

1. Infection by anaërobe bacilli occurs in about 10 per cent. of all cases of haemothorax from gunshot wounds of the chest that are seen in base hospitals in France.

2. The infection leads to the development of malodorous gas which may occasionally be so abundant as to simulate a large air pneumo-haemothorax, while the toxic features of the infection for the time being remain in the background.

3. In the majority of cases the septic features are much more prominent than gas formation. The most serious of these features are jaundice, a rapid pulse (which may be the index of commencing pericarditis), and semi-delirious restlessness. Less dangerous is the febrile type with a brilliant hectic flush on the cheeks.

4. Diagnosis depends on exploratory puncture. In cases with the clinical aspect of sepsis it may be necessary to

repeat the exploration on successive days, because it is a characteristic of anaërobe infection that the bacilli frequently fail to spread at once throughout the haemothorax, so that their presence is often missed at the first searching.

5. Life can be saved in at least four-fifths of the cases if the infected blood is promptly drained away. A day's delay in putting such treatment into force, when once the infection is well established, may mean death to the patient.

6. The clinical pictures caused by the sepsis vary widely in the individual cases of infection, but these differences cannot yet be correlated with the various forms of anaërobie bacilli that are proved to be present in each particular case.

REFERENCE.

¹ E. Foisy: *La pratique de la chirurgie de guerre*, Fascicule II, 1916, p. 185.

A CASE OF SPIROCHAETOSIS ICTERHAEMORRHAGICA.

By CAPTAIN R. RIMMER, R.A.M.C.,

M.O. LAB. LABORATORY, NO. — GENERAL HOSPITAL.

LITTLE evidence of epidemiological importance could be obtained from the patient whose case is here described. No previous case of jaundice had occurred on the boat, to his knowledge, and, since the boat does not call regularly at this port, no later information is available. Questioned as to the presence of rats, he was not able to state that they had been more noticeable than is usual on a ship.

A steward on H.M.T. —, aged 27, complained of vomiting for four days, vague epigastric pain and tenderness, and headache. The tongue was coated, and his mental condition was not clear. His replies to questions as to onset and present condition were vague and rambling. The temperature was 99.8° and the pulse 100, with no special characters. There was no jaundice.

On the morning of December 1st, 1916, he was markedly jaundiced, and his mental aberration was more noticeable. There was no vomiting; his tongue was brown and very furred (centrally disposed). Temperature 97.8° and pulse 86 in the morning; temperature 98° and pulse 90 in the evening. A guinea-pig inoculated intraperitoneally with 2 c.cm. of blood on this day remained healthy for the ten days it was under observation, having no pyrexia, and increasing steadily in weight.

On December 2nd the symptoms were more marked, the patient being delirious and of a deep orange colour, which apparently varied in depth from one part of the day to another. Small cutaneous haemorrhages were present on the limbs and trunk. There was also a small amount of haemorrhage from the mucous membrane of the mouth. He had incontinence of urine and faeces. There was no gross change in the size of the liver. *Tache cérébrale* was present. Herpes was not present. The faeces showed no special characteristics. The urine contained albumin, bile, and casts, mostly granular, but no leucin or tyrosin. He did not complain of any irritation of the skin.

A guinea-pig inoculated intraperitoneally with 2 c.cm. of urine showed fever (temperature 104°) on the sixth day, jaundice on the seventh day, and died with subnormal temperature and signs of collapse on the ninth day of observation. At the *post-mortem* examination haemorrhages of various sizes were found throughout the lung tissue on the right side; at the base was a considerable area of consolidation. The liver showed small patches (2 to 3 mm. in diameter) of a brick-red colour throughout its substance. Spirochaetes were present in the liver and blood, demonstrated by prolonged staining of smears with Giemsa, by silver nitrate, and Indian ink methods. In sections of the liver and kidney treated by Levaditi's method the organism was plentiful.

A second guinea-pig was inoculated with an emulsion of liver from the first animal, and the disease ran a very similar course. On the tenth day it was markedly jaundiced (with bile in the urine), and very dull, with loss of appetite, and a subnormal temperature. It was killed on this day. The *post-mortem* examination showed the usual haemorrhage into the lungs, and the liver was enlarged, very congested, and soft in consistency. The spirochaete was present in the liver and blood as before.

A third guinea-pig, inoculated from the second, died on the eighth day, and the liver showed the same bright red areas as were observed in that of the first animal.

The patient, on the fifth day after admission (probably the ninth day of the disease), showed marked improvement, the tongue cleaning, the degree of jaundice lessening, and his mental condition becoming normal, with return of control over both sphincters. No treatment was attempted. He took nourishment well throughout, the bowels moved freely, and the pulse was always good in quality. He subsequently developed some bronchitis, and had an evening temperature for some days

before evacuation. His condition when evacuated was quite satisfactory. He was still jaundiced, wasted, and anaemic. He stated that he felt quite well, and was apparently convalescing rapidly. When last heard from, on December 27th, 1916, he was up and expecting to be discharged from hospital in a few days.

CEREBRO-SPINAL MENINGITIS.

PRELIMINARY NOTE ON CERTAIN PATHOLOGICAL FINDINGS
AND THEIR BEARING ON TREATMENT.

BY

W. E. CARNEGIE DICKSON, M.D., B.Sc., F.R.C.P.E.,
PATHOLOGIST TO FULHAM MILITARY HOSPITAL, ETC.

OVER a year ago the results of *post-mortem* examinations on our fatal cases of cerebro-spinal meningitis gradually impressed upon my colleague, Captain A. Charles E. Gray, R.A.M.C., and myself the importance not merely of the mechanical blocking of the outflow of cerebro-spinal fluid from the cerebral ventricles by fibrino-purulent exudate in the region of the foramen of Magendie, but also of the actual, and in many cases widespread, involvement of the ventricles themselves in the inflammatory process. In practically every fatal case so examined the cerebro-spinal fluid in the whole chain of ventricles was found to be turbid—often considerably more so than the fluid outside the brain and in the spinal canal. Flakes of fibrino-purulent material were often present in the fourth and lateral ventricles, even in cases where little dilatation had yet occurred, whilst in cases in which hydrocephalus was supervening the inflammatory process was frequently more marked than on the external meninges. Microscopical examination of the fluid and fibrino-purulent exudate from the various ventricles usually showed large numbers of meningococci, sometimes in such numbers as to suggest the appearance of a "culture" of the organism. In many of these cases meningococci had become scanty or had disappeared from the fluid obtained by lumbar puncture during the clinical course of the case, and the important bearing of this finding upon treatment was early impressed upon us. After careful discussion with us, Major Lee, surgeon to the hospital, a year ago performed ventricular puncture upon several of the most serious cases, with a view to relieving the ventricular condition, and giving the patient a last chance of recovery. In the ventricular fluid so obtained meningococci were found in large numbers.

A second point of interest in connexion with the blocking of the outflow of fluid from the ventricles is that congestion and swelling of the brain substance, as well as blocking of the foramina in the roof of the fourth ventricle, may play a part in its production by forcing down the cerebellum into the foramen magnum around and in close apposition to the medulla, so that the foramen is sometimes blocked up as by a cork. The more the swelling of the brain increases, and with hydrocephalus also supervening, the more firmly is this "cork" forced down into the foramen, and the less possibility is there of relieving the pressure by lumbar puncture or any form of spinal drainage. In such cases I would strongly urge the advisability of ventricular puncture and drainage, as well as the introduction of antimeningococcal serum directly into the lateral ventricle, a procedure which Major Lee contemplates carrying out on a more extensive scale than heretofore. This should be done not merely in moribund cases as a last resort, but be thought of in cases in which, often after a preliminary improvement, cerebral symptoms increase. In such cases the examination of the fluid obtained by lumbar puncture alone may lead to an erroneous impression of improvement. Inflammatory cells may become less numerous and more mononuclear in character, meningococci may disappear, and yet the cases do not improve in a corresponding degree. *Post-mortem* examination may show in such cases a "clean" spinal cord but the ventricular infection above described. Nor need there be actual complete blocking by exudate of the foramina in the roof of the fourth ventricle to produce this result, though of course it is a common factor. Captain Gray has occasionally observed that relief of the cerebral pressure, probably by amelioration of the condition and subsidence of the swelling of the brain tissue, may

allow of fluid being drawn off by lumbar puncture where previously little fluid could be obtained—possibly by releasing the cerebellar plugging of the foramen magnum. It appears possible that in certain cases of this nature, while the pressure within the skull remains high, drawing off fluid from below may actually aggravate the condition by pulling the cerebellum still further down against the medulla.

Another point which we investigated over a year ago was the facility with which fluid introduced by lumbar puncture gained access to the meninges and ventricles. In several cases of meningitis and also in non-meningitic cases, as soon after death as possible, we injected a strong solution of methylene-blue by lumbar puncture, under conditions as regards amount, pressure and position as closely as possible approximating to the clinical injection of antimeningococcal serum. We found that, in non-meningitic cases and in meningitic cases without blocking of the openings in the epithelial and pial roof of the fourth ventricle, the 30 c.cm. of fluid injected at 15 in. of water pressure ran readily up the spinal canal and over the surface of the brain to the vertex and inner aspects of the cerebral hemispheres, and also into the chain of cerebral ventricles, reaching to all the tips of the horns of the lateral ventricles. On the other hand, where the foramina were blocked, the methylene-blue either penetrated only very slightly into the fourth ventricle or not at all, even though the surface of the brain was reached by the fluid. In one case, where there was extensive basal fibrino-purulent exudate and "corking" of the foramen magnum, scarcely any staining of the cerebellar and cerebral meninges occurred.

In a large number of cases examined *post mortem*, extensive softening of the spinal cord was found—most frequently in the upper dorsal, or upper dorsal and lower cervical region.

Several cases of the "serum reaction" which occurs in the cerebro-spinal fluid in non-meningococcal cases (tuberculous and syphilitic) in which antimeningococcal serum had been administered have come under my notice. These were punctured outside the hospital, and the serum administered before admission. On the day following such injection of serum the cerebro-spinal fluid withdrawn by lumbar puncture was turbid (in the tuberculous cases markedly so), and on microscopical examination was indistinguishable cytologically from the exudate of cerebro spinal meningitis, a well marked polymorphonuclear reaction occurring early, followed by progressive increase of mononuclears and phagocytic phenomena, etc., completely masking the real pathological condition. It was usually a matter of difficulty to obtain a sample of the first lumbar puncture fluid removed outside, but when this was possible the comparison was interesting and instructive, and often cleared up the doubtful diagnosis. The serum reaction of the cerebro-spinal fluid in meningococcal cases is also very interesting and important, but is beyond the scope of the present note, and I merely mention one of the most outstanding phenomena—namely, the bringing out into the fluid of a fresh "crop" of cells, mainly polymorphs, so that one can distinguish what I have termed double or even triple "broods" of cells—one fresh and undegenerated, staining well, another degenerated and undergoing "phagolysis" in various ways, and perhaps the third almost disintegrated—these crops corresponding to the reactions following the various injections of serum. This reaction appears to vary somewhat with the special brand of serum used.

In addition to the injection of antimeningococcal serum, washing out with saline, etc., we have on various occasions experimented with the injection of sodium citrate with a view to the prevention of the formation of fibrin, and of ensol as an antiseptic, but our results have not so far been sufficiently encouraging to allow of our publishing details. We hope to try the effect of flavine injections as soon as a sample of that substance is at our disposal.

NOTE.—Since the above was written Captain Gray has injected several cases with flavine. The results will be published later.

ON March 7th the American Codes Committee threw out bills intended to legalize the dissemination of birth control literature throughout New York State.