millimetres—seldom, if ever, more. You will obtain much better results by going slowly and by tiring out the stricture gradually.

In the bulbous urethra you can generally screw up at the final sitting to 45 Charrière. In the penile portion in some narrow urethras you may never get beyond 30 Charrière, in average urethras 35 Charrière. Never screw beyond 35 Charrière in the penile unless you are dealing with a very large urethra.

Never use Kollmann's dilators in the acute stage (five to eight weeks). Reserve them for the chronic stage, and never use them unless you have previously examined with the urethroscope and have also passed a nickel bougie on a previous occasion (No. 24 Charrière). As a general rule, do not use the Kollmann until the urine is clear of purulent haze. I advise you never to use Kollmann's dilators in the deep urethra. Only under the most exceptional circumstances does it do good in that situation, and you are likely to do much mischief with it. I know this is in direct contradiction with the teaching of many foreign authorities. All the same, I advise you to make it a general rule to which there are few exceptions.

Care of the Kollmann Dilators.

Irrigating Kollmann's dilators can be boiled after use and kept in a jar of lysol and distilled water, which prevents rusting. At the end of a session they ought to be boiled, dried out of methylated spirit, then dried in an oven, or by means of hot air, or with a "Fon." Finally, they should have sterile vaseline or paroline worked into each of their joints before being put away.

NERVE GRAFTING AS A MEANS OF RESTORING FUNCTION IN LIMBS PARALYSED BY GUNSHOT OR OTHER INJURIES.

BY

A. W. MAYO-ROBSON, C.V.O., C.B., D.Sc., F.R.C.S.,
MAJOR R.A.M.C.T. (LATE TEMPORARY COLONEL).

At the present time among the wounded returning from the battlefields are large numbers of cases of paralysed limbs due to more or less serious injuries to nerves.

In many of these cases the nerves are simply contused, and recovery is brought about by Nature's efforts in course of time, but in others, where there is complete destruction of a considerable section of the chief nerve trunks, recovery is impossible without operation. I hope it will not be considered out of place if I draw attention to work which I performed on this subject so far back as 1888.

In February, 1889, I read a paper before the Clinical Society of London on a case of nerve grafting, which was, I believe, the first in which such an operation had been performed in this country on the human subject.

The patient was a girl, aged 14, whose hand was paralysed after the removal of a tumour from the forearm which had involved taking away over two inches of the median nerve. Repair was effected by grafting 2½ in. of the posterior tibial nerve from an amputated leg of another patient, the transfer being effected within a few minutes of the removal of the nerve from the amputated limb. The patient was shown at the Clinical Society, Mr. Christopher Heath being in the chair. A discussion followed, in which Mr. Anthony Bowlby (now Sir Anthony Bowlby, K.C.M.G.) and Mr. Bland-Sutton (now Sir John Bland-Sutton), Mr. Bryant, and the President took part.

Sensation returned within thirty-six hours of the operation, and motion in the paralysed muscles returned gradually. When the patient was shown at the Clinical Society Mr. Bowlby remarked that it was the first recorded case of nerve grafting in this country. I had the opportunity of seeing this patient three years subsequently, when recovery was perfect in every respect.

In July, 1890, my late colleague, Mr. Atkinson, described two cases of his own, two cases of mine, and one case of Mr. Ward's, all of which were, at the time of being reported, progressing satisfactorily or had completely recovered.

The following case is still more important, as it shows that in the absence of recently removed nerve, such a structure as the spinal cord of a rabbit may be successfully used as a graft.

W. R., aged 29, a gardener, was admitted to the Leeds Infirmary under my care on January 18th, 1890, having been sent to me by Dr. Mackay of Hornby. He gave the history of having, seven months before, fallen on a scythe, which had produced a deep incision on the lower and inner part of the upper arm, the brachial artery being divided. At the time of injury a surgeon ligatured the artery, and the two ends of a divided nerve were sutured. The wound healed slowly by granulation.

Condition Before Operation.

On admission to the infirmary the patient was in robust health. There was a scar on the inner aspect of the right arm, extending from a point 2 in. above and to the inner side of the olecranon process, obliquely upwards and forwards for about 3 in. to the inner side of the prominence of the biceps. The circumference of the right forearm was two-thirds of an inch less than that of the left, that of the wrist half an inch less. The hypothenar eminence was absent, the thenar eminence was much atrophied, and the interossei muscles were very much wasted. He had entirely lost the power of grasping, though he could flex the metacarpo-phalangeal articulations to a right angle. Extension of the wrist was possible, apparently with considerable exertion, but flexion of the wrist was impossible against gravity. The thumb could not be adducted nor the fingers separated or approximated. All the muscles acting on the wrist and hand which had their nerve supply from the median and ulnar nerves were paralysed, while those supplied by the musculospiral were capable of acting.

Sensation was absent along a narrow line extending over the inner and anterior aspect of the ulnar border of the forearm from the elbow to the wrist. In the palm, sensation was absent except over the thenar eminence and part of the proximal phalanx of the thumb. On the back of the hand it was absent internal to a line drawn backward from the cleft of the middle and ring fingers, also over the whole of the back of the little and ring fingers, as well as over the distal one and a half phalanges of the index and middle fingers. The extensors reacted to faradism; the flexors showed no reaction. To a current of 10 milliamperes the C.C.C. was greater than the A.C.C. both in the flexors and extensors.

It was manifest that there had been a complete division of the median, ulnar, and internal cutaneous nerves, and it was clearly useless to attempt anything short of operative interference.

Operation.

On January 30th, 1890, the limb having been previously aseptically ether was administered, and an incision made along the line of the cicatrix; it was prolonged some distance upwards and downwards, and supplemented by a transverse incision about an inch above the elbow. The ulnar nerve was soon exposed above the internal condyle; it was firmly fixed in fibrous tissue. The lower end of the upper segment, which was bulbous, was connected by fibrous tissue with the upper end of the lower segment. A long search was then made for the median nerve, as the large cicatrix had altered the normal anatomy of the parts. A small nerve was discovered at the upper part of the wound, which proved to be the internal cutaneous. The lower end of this nerve was discovered later, and united to the upper by a catgut suture. The internal intermuscular septum was divided 1½ in. above the internal condyle, and the atrophied remains of the brachial artery defined. At last, about the middle of the upper arm, the bulbous lower end of the upper segment of the median nerve was discovered well under, and concealed by, the belly of the biceps. Later the upper end of the lower segment was found just above the bend of the elbow. This extremity was expanded, and sent ramifications into the cicatrix. The fibrous tissue between the ends of the ulnar nerve was excised, and the two healthy portions were united by grafting strands of the sciatic nerve of a rabbit so as to fill up the gap and produce continuity. It was quite impossible to bring the divided ends of the median nerve nearer than 2½ in. All the nerve tissue at my disposal having been exhausted in uniting the ulnar nerve, Mr. Littlewood, who was assisting me, dissected out the spinal cord of a rabbit just killed, and this was used as a graft to connect the ends of the median nerve, the inserted cord lying loose and quite free from tension when finally placed in position. Fine catgut sutures were used throughout. The edges of the wound were brought together and the usual dressings applied, the arm being fixed on a rectangular splint. Healing occurred by first intention, and there was an entire absence of fever or pain.

Early After-History.

On February 10th the patient could feel the scratch of a pin on the flexor aspect of the first phalanx of the thumb, as well as at the root of the index finger. He could tell when the hairs on the back of the first phalanges of the ring and little fingers were pulled, but could not feel the scratch of a pin in that position.

On February 17th sensation had returned over the whole of the palmar surface of the thumb and the proximate phalanx of the index finger.

On March 4th sensation seemed to be creeping slowly along the first finger, and to be present over the whole of the palmar area supplied by the median nerve, and extending as far down as the web of the fingers, and a short distance along the middle finger. As yet there was no marked improvement in the ulnar distribution.

On March 6th electrical reaction to faradism was absent in the muscles. C.C. 15 milliamperes, C.C.C. equal to A.C.C. in flexors; C.C.C. greater than A.C.C. in extensors. The muscles were

gradually developing, and the general nutrition of the hand showed improvement. There was slight power of grasp and some power of flexion in the wrist, as well as slight power of adduction of the thumb and flexion of the fingers. Sensation was felt all over the thumb and index finger, in the second finger up to the first phalanx on the palmar aspect, also in the third finger, though less distinct. No sensation could be elicited over the distribution of the ulnar nerve.

On March 30th he began to have sharp shooting pains along the distribution of the ulnar nerve. On April 1st the flexors of the forearm began to react visibly to C.C.C., though the muscles of the thenar eminence did not, and there was no faradic reaction. On April 30th sensation was returning at the back of the ring finger, and the flexors now reacted to A.C.C. of 5 milliamperes.

On June 13th the nutrition of the hand was much improved, and there was a greater feeling of warmth in it. He could pick up small objects, and although the grasp was not powerful, it was much improved. The powers of flexion of the wrist and adduction of the thumb were considerably increased, but there was no sensation as yet over the ulnar distribution.

Condition of the Patient Six Years after Operation.

Being interested in the case, I made inquiries from time to time after he left Leeds, but I never saw him again until February, 1896, six years after the operation, when I asked him if he would let me demonstrate his case at the Leeds and West Riding Medico-Chirurgical Society, where my colleagues and other members of the society had the opportunity of seeing him. Dr. Mackay had applied galvanism to the muscles of the forearm and hand from time to time, from the time of his leaving Leeds until the middle of November of the same year, when all treatment was stopped. He did not resume work till March, 1891, not because he was unable to undertake light work, but, as the patient put it, "He wanted to give the arm a fair chance of complete recovery." Since March, 1891, he had not missed a single day's employment, and was able to do all his duties completely, from wheeling a well-laden barrow to using a scythe.

There was a firm round scar along the inside of the arm. The circumference of the right forearm was 11 in., left forearm 11½ in.; right wrist 7 in., left wrist 7½ in.

From these measurements it will be seen that the muscles had almost completely returned to their former volume, and in the hand all the muscles except the abductor of the thumb were as well developed as in the left; this was quite as well marked in the interossei as in the other muscles. The movements of the arm were completely restored and almost as perfect as in the left. Flexion of the fingers and the grasp had completely returned, and the only weak muscle was the abductor of the thumb, which had no perceptible power. Sensation was completely restored, as was easily demonstrated by touching the different portions of the arm and hand with a pin.

All the muscles of the arm and hand reacted to faradism except the abductor of the thumb.

It will thus be seen that, though the recovery had been tardy, it was complete in almost every respect, the only exception being that of the abductor pollicis, which for some unexplained reason did not recover its function. The case is very encouraging, since it clearly demonstrates the possibility of restoring continuity of nerves by grafting. Whether the spinal cord in this case simply served as a basis on which the nerve tissue was built up I am unable to say, but that it answered its purpose is clearly shown in the report of the case. Why restoration of function in the ulnar nerve should have taken longer than in the median I am unable to say, as the ends of the nerve were not separated more than ¾ in., whereas the ends of the median nerve were between 2 in. and 3 in. apart. Could it be that the spinal cord offered a better medium for establishing continuity? The patient was seen years subsequently, completely recovered in every respect.

Since operating on the above cases I have had considerable experience in the surgery of nerves, both with regard to suture of divided ends and in nerve grafting.

During the present war I have had experience both in the Dardanelles, in Egypt, and in the base hospitals at home in the treatment of nerve injuries, and my previous experience in civil practice has stood me in good stead in dealing with this class of cases.

Nerve grafting cannot, of course, be successfully performed in septic wounds, but where the paralysis remains after healing of the wound and it is known that it is not merely due to division of nerve or of some portion of the nerve trunk, as shown by the paralysis not improving under treatment, it seems to me desirable that too much time should not be allowed to pass before operative treatment is adopted in one or other of the ways I have described above, as it will lead in a large number of cases, if not in all, to complete restoration of function and thus to saving of otherwise useless limbs.

NOTE ON THE INCIDENCE OF TETANUS AMONG WOUNDED SOLDIERS.

BY

MAJOR-GENERAL SIR DAVID BRUCE, C.B., F.R.S.

The principal object of this paper is to draw attention to the effect of prophylactic injections of antitetanic serum in lowering the incidence of tetanus among wounded men. The chart represents the ratio of cases of tetanus to number of wounded men from the beginning of the war to the end of 1916. This diagram has been made up, not from all the cases of tetanus which have occurred, but only from those which have arisen in home hospitals among the wounded men sent to England from overseas. It does not include cases which have occurred in France or elsewhere.

It will be seen that the ratio in September, 1914, is 16 per 1,000; in October, 32 per 1,000; and that it falls in November to 2 per 1,000, about which figure it remains until the present date. To what cause must this remarkably sudden fall be attributed? No doubt there were various factors at work, but beyond all reasonable doubt the introduction of prophylactic injections of antitetanic serum was the preponderating cause. To understand this it may be well to give a short account of the movements of our army during these first months.

The bulk of the British Expeditionary Force crossed over to France on the night of August 12th and 13th, 1914. On the 18th and 19th the troops began to move northwards into Belgium.

The battle of Mons and the retreat took place between August 23rd and 30th. During the retreat many of the wounded fell into the enemy's hands, and on our own part it was impossible in many cases to carry out any thorough surgical treatment of the wounds owing to the hurried nature of the retreat.

The battle of the Marne was fought between September 7th and 10th.

On September 13th the British Expeditionary Force reached the valley of the Aisne, and remained here until the end of the month. This was the ground where probably many cases of tetanus were infected. The soil is rich and fertile, and has an evil reputation for tetanus. Here in ordinary times it is found necessary to inoculate the farm horses at intervals with tetanus antitoxin as an ordinary measure of precaution.

The troops left the valley of the Aisne about October 3rd, and took up positions at La Bassée and its neighbourhood, where fighting took place from October 11th to the end of the month.

The severe fighting round Ypres, in which a separate division of the army took part, began on October 20th and continued until November 11th, 1914.

It is evident, then, that the 16 per 1,000 cases of tetanus among the wounded which occurred during September, must have been contracted at the battle of the Marne and in the valley of the Aisne. Further, that the abnormally high ratio of cases of tetanus which is found in October must have been to a great extent due to the tetanized soil of the Aisne valley.

There were therefore several factors at work in September and October, 1914, to raise the ratio—the heavy fighting with the consequent heavy toll of wounded, the highly infective character of the soil, the difficulty of collecting the wounded on account of their numbers and the movements of the troops, and finally the difficulty of giving the thorough surgical treatment to their wounds which is so essential in the fight against tetanus.

But there was another, and that the most important, factor which has not been taken into account—the prophylactic injection of tetanus antitoxin. This was not carried out during the first two months of the war. It would appear that in the beginning only a small quantity of antitoxin was taken out with the Expeditionary Force for purposes of treatment, and it was only when the number of cases of tetanus became alarming that steps were taken to ensure that every wounded man should receive a prophylactic dose.

Lord Kitchener telegraphed to the Director of Medical Services, Expeditionary Force, on September 8th as follows:

Earl Kitchener desires information as to whether antitetanus inoculations are being practised for wounded, and whether,