were used in the two cases because two-thirds of the circumference of the colon was injured; in the other case the caecum had two or three small perforations. He had not used ether as a peritoneal dressing.

TWO CASES OF REMOVAL OF A BULLET FROM THE UPPER SURFACE OF THE DIAPHRAGM.

BY

E. D. TELFORD, F.R.C.S., CAPTAIN R.A.M.C.(T.F.),
SURGEON, MANCHESTER ROYAL INFIRMARY.

WHILST the lessons of war surgery have thrown much light on the immediate treatment of penetrating wounds of the chest and have prompted some most valuable additions to surgical literature, the later consequences of these injuries have received little notice.

It is a fortunate fact that not a few of these seriously wounded men recover, but some still carry in their chests the agent, whether shell or bullet, which caused the injury. These foreign bodies show a tendency to fall to the bottom of the thorax, coming to rest there on the upper surface of the diaphragm. In this position they may give rise to no trouble and no surgeon would suggest their removal. There are found, however, from time to time cases in which the foreign body is a cause of persistent and distressing symptoms.

In the first case which I report the man's life was made miserable by a constant "pricking" pain, and in the second case the soldier was quite incapable of any exertion, owing to the fact that an increase of respiratory movement beyond the normal produced at once a crippling amount of pain. The two cases are reported, not merely as examples of removal of foreign bodies from an unusual situation, but rather as instances of the help afforded to the surgeon by a simple method of "positive pressure anaesthesia." Thanks to this method, most ably handled by Major A. Wilson, the recovery of a bullet from the pleural cavity was carried out with perfect safety to the patient and was rendered, from the operator's point of view, little more than a minor surgical procedure. The details of the cases are as follows:

CASE I.

G. W. C., aged 24, was wounded in the right side of the chest by rifle bullet fired at long range. The bullet entered through the third right costal interspace three and a half inches from the middle line. Free haemoptysis occurred at once and continued in considerable quantity for twenty-four hours. He reached England at the end of five weeks, and during this time he had much pain in the lower part of the right chest, with frequent cough and daily haemoptysis. Although he stated, on admission to the 2nd Western General Hospital, that he had lost 2 st. in weight since his injury his general condition was excellent. The chest expanded well, and beyond a few moist sounds there was no physical sign of disease or injury. Cough was frequent, and the sputum always contained a little dark blood. The patient, though apparently well and comfortable in a sitting posture, was not able to lie down with ease, and was rendered very breathless by slight exertion. He complained constantly of a "pricking" pain, which he located deep to the sixth right costal cartilage.

Radiographic examination by Captain Bythell disclosed an intact rifle bullet lying on the upper surface of the diaphragm. The bullet was localized at a depth of two inches beneath a point marked over the sixth right costal cartilage and distant one half inch from the line of the right border of the sternum. As the pain continued after a month in hospital, and the patient's condition remained without change, save that the haemoptysis had ceased, it was decided to attempt the removal of the bullet.

Anaesthesia was induced by chloroform, and the tracheal catheter of the Ehrenfried's positive pressure apparatus having been introduced, the operation was carried out under ether vapour introduced under pressure by bellows. Through a curved skin incision the sixth and seventh right costal cartilages were exposed and denuded of their muscular and tendinous coverings, which were reflected downwards in one piece. A portion, $1\frac{1}{2}$ in. in length, of each costal cartilage was then excised. The opening thus made gave abundant access, and through it the parietal pleura was incised in a horizontal direction. A healthy, well-inflated lung presented itself on opening the pleura. The lung appeared and felt absolutely normal, showing no trace of adhesions.

On packing off the lung with a strip of gauze there was disclosed at once on the upper surface of the diaphragm a rounded eminence, in size and shape like the half of a small walnut shell. An incision into this yielded a few drops of turbid fluid, and the bright, unbroken nose of the bullet at once presented. This was seized and extracted without difficulty; it was lying

on the upper surface of the diaphragm, and had become covered and encysted by pale, friable fibrous tissue.

The gauze was removed and the parietal pleura closed tightly by a continuous stitch of fine chromic gut. The tendino-muscular flap was replaced by chromic gut sutures and the wound closed without drain. The colour and general condition of the patient were perfectly satisfactory throughout the operation.

Beyond a slight tracheitis, probably traumatic, which cleared up completely in three days, the convalescence was uninterrupted. The patient was discharged from hospital in four weeks. The "pricking" pain disappeared from the first and the man was able to move freely without breathlessness.

CASE II.

T. J. E., aged 21, was hit by a shrapnel ball at Suvla Bay. He stated that very free bleeding followed the injury—"blood running out of his mouth without cough." He gradually recovered, and does not appear to have suffered from any complications of the injury beyond occasional slight haemoptysis, which continued up to the time of his admission to the Manchester Royal Infirmary, six months after the injury. At this time the chest showed no abnormal physical signs and his general condition was excellent. The scar of the wound of entry lay over the inferior angle of the left scapula. The patient complained of much pain on exertion and located the pain at a point on the tenth left rib, a little in front of the mid-axillary line; he stated that the pain was very severe on deep breathing.

A radiogram by Captain Barclay showed an intact shrapnel bullet lying deep to the rib and fixed in the angle between the diaphragm and chest wall.

Anaesthesia was carried out, as in Case I, by Major A. Wilson with the help of Ehrenfried's apparatus. The operation was of the simplest nature. A portion of the tenth rib, about $1\frac{1}{2}$ in. long, was resected and the pleura opened by a horizontal incision. The bullet was at once evident and was picked without effort off the upper surface of the diaphragm. The edge of the well-inflated lung was well seen, and as far as it could be inspected through the limited opening the lung appeared to be quite healthy and free. The pleura was closed with fine chromic gut.

The after-history of the case was quite without incident; the patient was discharged in a fortnight in excellent condition, having had neither haemoptysis nor his former pain since the operation.

TIME IN SURGERY.

BY

R. P. ROWLANDS, M.S., F.R.C.S.,
SURGEON TO GUY'S HOSPITAL.

THE priceless value of time in surgery is often overlooked, and fleeting opportunities of successful action are frequently allowed to pass with disastrous consequences. Again and again diagnosis is delayed or operation deferred until it is too late either to save life or to secure the best result. For instance, the early signs of grave abdominal diseases are too often attributed to trivial ailments, and a "wait and see" policy adopted until such irreparable damage is wrought that complete recovery is impossible. In such cases the greatest vigilance is required to determine the proper time to "look and see."

Two typical cases of appendicitis serve to illustrate these points. They were very similar in their early signs, but treated differently and attended by very different results.

CASE I.

- (a) Diagnosis made within twelve hours.
- (b) Operation within thirteen hours.
- (c) Appendix removed, gangrenous, but not perforated.
- (d) General condition good.
- (e) Gridiron incision.
- (f) Abdomen closed.
- (g) Patient up from fourth day; taken to the country on the ninth day. Quite well three weeks from operation.
- (h) Patient has been in perfect health since the operation.

CASE II.

- (a) Diagnosis, abdominal influenza. After nine days, appendical abscess diagnosed.
- (b) Operation after ten days.
- (c) Large abscess opened. Appendix not found owing to the large size of abscess and the grave condition.
- (d) General condition very bad.
- (e) Battle incision, as the surgeon thought he could get more room in this way.
- (f) Drainage on and off for six months; then appendix removed at second operation. Drainage another two months.
- (g) Patient up after ten months. Large gap in abdominal wall with hernia at the end of twelve months. General condition too bad for operation to reconstitute the abdominal wall at present.
- (h) Patient has suffered much pain and misery.

It ought to be more generally known that the mortality of operations for acute appendicitis increases rapidly with the delay. Operations carried out within twelve hours of the onset of an attack are not more risky than those undertaken in the quiescent period. Still, it is common for the surgeon to be asked to see patients for the first time about the third or fourth day, or even later, when abscess, peritonitis, or other serious complications have developed. Similarly, the mortality of rupture of the intestine is only 15 per cent. when an operation is undertaken within four hours, whereas it becomes 70 per cent. when the operation is delayed until after twelve hours.¹ Similarly, the mortality of perforated gastric ulcer when an operation is undertaken within twelve hours is only about 5 per cent., but as high as 60 per cent. when the operation is deferred until after twenty-four hours. The waste of time is almost as serious and even less excusable with chronic abdominal diseases such as cancer of the stomach or large bowel. For less than one-third of cases of cancer of the stomach when first seen by the surgeon is resection possible, and the result of procrastination is nearly as bad in cancer of the rectum.

Avoidance of Waste of Time at Operations.

Time is frequently wasted during operations. It is clear that in order to get good results the surgeon must do his operations well and thoroughly, but it is often forgotten that to get the best results he must also learn to do them quickly and without waste of time. On the other hand, he must never do shoddy work so that he may be quick. While always conscious of the time element, he must not keep one eye on the clock, but both eyes and all his faculties focussed on his task. His aim must be to get so skilful and familiar with his work that he becomes quick almost without knowing it.

It is obvious that the dangers of most operations increase with their duration. If an operation takes an hour when it can be just as well, or better, done in half that time, the disadvantage of the patient is serious. He has to take much more anaesthetic, which obviously adds to the dangers and discomforts still inseparable from anaesthetics. Further, manipulations and exposure are increased and prolonged with marked increase of shock, of post-operative discomfort, and danger of complications. These points do not need to be laboured, for they are evident enough to every one who has eyes to see. When an appendicectomy takes only ten minutes the patient is hardly ill or uncomfortable in any way, whereas if it takes an hour he is miserable and sore for days.

But the value of smart work is much more striking in grave crises such as operations for acute intestinal obstruction or late peritonitis. It is no exaggeration to say that in these cases the mortality multiplies with the time consumed at the operation. The following case illustrates the value of quick work.

Acute Intestinal Obstruction by a Band: Appendix Adherent to Caseous Mesenteric Gland.

A boy, aged 6 years, was seized with very violent pain in the abdomen on the morning of April 10th, 1911. His father gave him two large doses of castor oil, which failed to have the desired effect. He was admitted into Guy's Hospital with a subnormal temperature, blue, cold, and pulseless. The abdomen was considerably distended, but flaccid. It was full in the flanks, and there was a palpable swelling in the right iliac region. On admission at 8 a.m. he was too bad for operation, and was therefore infused into the axillae.

Operation.—The patient was given ether, and the abdomen was opened at 9.30 a.m. The lower part of the right rectus was displaced inwards. The peritoneum seemed to be purplish and clearly contained blood. It was opened, and a large amount of sanious fluid escaped, showing that the condition was one of intestinal obstruction and not appendicitis. Dark purplish, but not gangrenous, coils of small intestine were at once seen. Several coils were withdrawn, and a band was seen consisting of the appendix adherent to a caseous mesenteric gland. This was astride the lower end of the ileum, close to the caecum and completely obstructed it. The adhesion to the tip of the appendix was divided, and the empty caecum drawn away from its former position, which was unusually high. The abdomen was then closed with through-and-through salmon-gut sutures. The operation lasted seven minutes (the patient was only in the theatre ten minutes). He was not much worse at the end of the operation, but the infusion which was carried on throughout the operation was continued. Pituitary extract was injected in 5-minim doses. The patient rapidly recovered.

Our forefathers before the introduction of anaesthetics had no doubt about the value of rapid operating, but later

a mischievous belief became common that, with anaesthesia, time did not matter. Unfortunately, there are still too many who appear to share this opinion. They waste valuable time—which belongs more to their patients than to them—in many different ways. It may be profitable to consider how to avoid some of this serious waste of time.

1. *By Making All Possible Preparations and Calculations Beforehand.*—The skin is always prepared before the patient comes to the operation table. All necessary instruments, ligatures, and sutures threaded on needles are ready for use before the patient is completely under the anaesthetic. The surgeon and his assistant are quite ready to begin as soon as the patient is satisfactorily under the anaesthetic. The surgeon thinks out the various steps of the operation beforehand as far as this is possible. He makes up his mind quickly what he wants to do, and does it without unnecessary delay.

2. *By Improving and Simplifying the Technique of the Operation in Every Possible Way.*—Here a sound knowledge of anatomy and surgery and constant practice are essential. As regards the knowledge of anatomy and surgery, it is assumed that the surgeon will not undertake an operation unless he is familiar with the anatomy of the part and the details of technique of the operation which he contemplates. It is clear from the results of some of the operations for hernia recently undertaken that the operators were not familiar with the anatomy of the inguinal canal, for several cases have come to my notice of recurrence within a few months owing to the inguinal canal never having been opened. It is true enough that a surgeon has to get practice to learn, but a sound knowledge of anatomy and surgery, together with diligent practice on the dead body, will carry the beginner far. Every one who wants to become a good and quick operator should practise on cloth all varieties of knots and sutures. In this way he will become expert and competent in a short time. It is often painful to see a man hesitating about his knots, and uncertain about his method of sewing. Many men waste time by making the skin wounds too small, and thus allowing themselves only an unsatisfactory view of the parts. As a rule, except on the exposed parts, such as the face, there is no object in making a small skin wound. Much time is also wasted in blunt dissection instead of clean cutting. Both of these faults not only waste time, but add to the shock and local damage. Good haemostasis saves much time. For instance, in most operations on the limbs a tourniquet is used, so that no bleeding occurs to obscure the view and delay the work. In other operations the main artery to the part is temporarily clamped or tied. For instance, the carotid is often clamped while operating on the neck. In most operations the important vessels can be secured before they are divided. Complicated instruments are to be avoided, for they waste time. For instance, a needle holder has no advantage in the majority of operations, but adds to the number of movements necessary for a given result. It is, of course, an invaluable and necessary instrument when working in a pit like the bottom of the pelvis. In ordinary work the finger and thumb are much more accurate, adaptable, and infinitely quicker. If one were to judge by the instruments they use, it might be concluded that some surgeons enjoyed making operations as difficult and prolonged as possible. The same may be said of many complicated methods, which are rarely as efficient as simpler ones. Continuous sutures save much time, and they are more effective and haemostatic than interrupted. In grave emergencies an excess of fine detail in sewing is often to be avoided; the sewing should fit the occasion. Lives may be lost through the waste of time involved by an over-refinement of technique.

3. *By Concentrating the Full Attention upon the Operation.*—This is not the time for considering and talking of other matters, nor even for explaining the various steps of the operation to the inquiring stranger. The latter must be content to watch until the operation is completed.

REFERENCE.

¹ *Lancet*, 1908, vol. ii, p. 1143.

THE Italian Minister of Agriculture, Industry, and Commerce has appointed a committee to study the creation and development of industries for the production of chemical preparations likely to be useful for medicinal purposes.