marked susceptibility to thrombosis. One man whose veins were easy to puncture had a local thrombus after each injection. Soft thrombi sometimes form during the injection, and it is then dangerous to proceed. One first sign of local thrombosis in such a case is hindrance to the flow from the needle.

GENERAL OBSERVATIONS.

1. The general tone and appearance of the patient improve so markedly after one or two injections of salvarsan, and the primary sore so soon disappears, that the practitioner may be persuaded by the patient to stop treatment. We have seen several cases in which this has occurred, with the result that a very early relapse has taken place. Care should be taken to continue the treatment until all clinical signs have disappeared and the blood reaction is negative.

2. The speed with which primary sores, secondary rashes, and condylomata disappear is worthy of note; two or three injections will usually remove all outward signs.

3. The effect of salvarsan is not so speedy in cases in which the mucous membranes of the mouth and tonsil are affected, nor in glossitis, and they do not usually heal until the blood has got to a negative phase.

A SIMPLE DIETETIC TREATMENT IN DIABETES MELLITUS.

BY

R. T. WILLIAMSON, M.D.LOND., F.R.C.P.,
CONSULTING PHYSICIAN, MANCHESTER ROYAL INFIRMARY.

WHEN the ordinary dietetic treatment fails to check the glycosuria in cases of diabetes mellitus we have now many new methods of treatment which, temporarily or permanently, are of distinct service. Some of these methods have, however, disadvantages, and occasionally are followed by very unpleasant symptoms.

I desire to draw attention to a simple method of dietetic treatment which I have found of the greatest service, in many cases of diabetes, in checking the glycosuria, at least temporarily. It is free from risk, quite palatable, and can be taken by almost all patients. It is worthy of trial in most cases of diabetes if the ordinary diabetic diet does not promptly check the glycosuria.

The treatment lasts for a week or ten days only; during this period the patient ceases work and rests on the sofa. Every two hours from 8 a.m. to 10 p.m. he receives a small amount of food—eggs, cream, milk, beef-tea, and tea—according to the following diet sheet.

Diet.

- 8 a.m.—Coffee, or tea, with one tablespoonful of cream. One egg (poached, boiled, or buttered).
- 10 a.m.—A glass of warm milk (half a pint).
- 12 noon.—Custard (prepared from one egg and half a pint of milk).
- 2 p.m.—A glass of warm milk (half a pint).
- 4 p.m.—Tea, with one tablespoonful of cream. One egg (poached, boiled, or buttered).
- 6 p.m.—Cream, two tablespoonfuls, in half a pint of warm beef-tea.
- 8 p.m.—A glass of warm milk, or one egg beaten up and added to half a pint of warm beef-tea.
- 10 p.m.—Cream, two tablespoonfuls, in half a pint of warm beef-tea.

The order of these meals may be varied or slightly modified as the patient desires. The patient takes no other food. The bowels are kept regular.

In many cases the glycosuria ceases in a few days, in other cases in a week or ten days, although an ordinary rigid diabetic diet had failed to check the sugar excretion. If the sugar excretion does not cease on this diet of eggs, cream, milk, and beef-tea, carried out as just described, in the course of a week or ten days, I discontinue it and try some other method of treatment. After the glycosuria has been checked I change gradually, at the end of seven or ten days, to a diet of solid food. First, a little diabetic bread* is allowed with the tea and egg at 8 a.m. and 4 p.m., and two of the other meals are diminished or omitted. Then a day or two later bacon and green vegetables are

allowed, followed by custard, in place of the meals at 12 noon and 2 p.m., and the meals at 10 a.m., 6 p.m. and 8 p.m. are omitted, and the tea and egg taken at 5 p.m. instead of 4 p.m. If the glycosuria does not return then the bacon at midday dinner may be replaced by fish, and later by fowl or meat. Later the ordinary diabetic diet may be allowed, and still later, if no sugar returns in the urine, a small amount of white bread may be given.

Of course, this treatment for a week or ten days with the diet of eggs, cream, milk and beef-tea, in the manner just described, does not remove the glycosuria in all cases of diabetes; but it is interesting to note how often it will do so, when a previous treatment with the ordinary rigid diabetic diet has entirely failed to check the sugar excretion.

If the glycosuria returns at a later date, the patient may try again for one week the diet of cream, eggs, milk and beef-tea, and repeat it from time to time, if necessary.

The permanent value of this treatment cannot yet be estimated, but it is a method which is more prompt and powerful than the ordinary rigid diabetic diet.

In the BRITISH MEDICAL JOURNAL of March 13th, 1915, I published a method of treatment with casein or biogene (mixed with cream and water) which I had found of great service. Since that date I have used this method very frequently and often with excellent results, and my further experience has confirmed the opinion I then expressed as to its value. It requires, however, to be carefully watched, and, as I previously pointed out, it occasionally causes unpleasant symptoms, which can, however, soon be checked. Occasionally bad specimens of casein or biogene are obtained which cannot be taken by the patient, and occasionally a case is met with in which a satisfactory preparation of casein or biogene is most unpalatable to the patient and cannot be taken. The casein preparation known as Laitproto No. 6† may then be tried in place of casein, and in the same quantity, and may be well taken. If this should be very unpalatable and cannot be taken, the method of treatment with milk, cream and egg described in this article can almost always be taken quite well.

The dietetic treatment (with cream, eggs, milk, and beef-tea) just described is (1) simple; (2) less expensive than diabetic breads and foods, and specially suitable now on account of the increased price of diabetic foods through the war; (3) it can be taken by nearly all patients, and is quite palatable; (4) it can be taken by patients who cannot take casein or biogene; (5) it does not require any very special care or watching; (6) it is very useful and suitable in the case of diabetic children; (7) it is often very successful, and often removes the glycosuria when an ordinary rigid diabetic diet has failed; in other cases, like all other treatments, it is not successful; (8) after taking this diet for seven or ten days the patient may then often take an ordinary rigid diet, and continue it for a long period without the return of the glycosuria, whilst previously such a diet had entirely failed to check the glycosuria; (9) in some cases it removes the diacetic acid as well as the sugar, but not in many others; (10) probably it is not so often successful as the treatment with casein or biogene (which I have previously published); but if it should fail, the latter treatment may then be tried, or a combination of the two methods may be tried.

The dietetic treatment described in this article should be continued for seven or ten days only. It is to be regarded as a method usually very much more successful than an ordinary rigid diabetic diet in removing the glycosuria temporarily, and its after-effects probably continue much longer. If later the sugar returns in the urine the treatment may be repeated. The earlier the treatment is tried the more likely is it to produce satisfactory results. It is worthy of trial in most cases of diabetes if the ordinary rigid diet does not check the glycosuria in one or two weeks.

† Laitproto No. 6 can be obtained from Casein Co., Limited, Sheepcote Lane, Battersea, London, S.W.

DR. LANPHER VERNON-JONES, of Arlington Street, S.W., left £16,808.

At the first meeting of the American Congress on Internal Medicine, held in New York on December 28th and 29th, 1916, an American College of Physicians was founded. Sixty-five physicians were elected Fellows.

* Biogene bread and ponos biscuits, protene bread, etc., or the coconut and biogene, and other cakes which I have described in the BRITISH MEDICAL JOURNAL, December 23rd, 1916.