operation. Saline given subcutaneously helps very much to combat these. If the temperature rises and there is pain, disinclination for food, and vomiting, the wound should be examined, and fresh spread of sepsis looked for. Should such be present and threaten life, amputation by the rapid circular method should be performed without delay.

If infection still persists in the wound, and is spreading, though not sufficiently to threaten life, treatment will depend upon the organism present. If *B. perfringens* and *S. reptococcus faecalis* (for these two organisms generally coexist) be the cause, then a barrier of hydrogen peroxide or potassium permanganate (strong solution) injected into sound tissue above and below the infected area, and completely encircling the limb, has done good in some cases and sufficed to arrest further spread. Further incision and drainage of the affected part may be needed, and this can be combined with continuous irrigation of the wound with normal or slightly hypertonic saline solution, alternating with eusol.

If the infection be pyogenic, hot antiseptic fomentations, or charcoal and carbolic poultices, will often limit the process.

In a few desperate cases I have tried intravenous injections of eusol, as recommended by Fraser and Bates, but have had no success from its use.

Stimulants may be freely given, and do good in these cases. As soon as possible these men are sent to the base hospital, where more perfect conveniences for subsequent treatment exist.

It must be clearly understood that what has been said in this article refers only to the treatment of fractures between the firing line and the clearing stations. Many cases of serious compound fractures arrive at the clearing stations in a very septic and foul condition. The general condition of such wounded men is grave, and amputation is their only hope of salvation. Attempts to save such limbs have, time after time, ended in disaster, owing to the rapid spread of infection in these fatigued and worn-out men. This will explain the number of cases of amputation reaching the base hospitals from the clearing stations. Further, these amputations have to be performed exceedingly rapidly, as the condition of these men is often desperate. Such operations are purely life-saving operations, and so the same finesse cannot be looked for as can be obtained in later operations when the patient's condition warrants a longer anaesthetic; hence the crude appearance of some amputation stumps when the base hospital is reached.

I am indebted to Mr. R. Brown, of the Kensington "War Hospital Supply Dépôts," for the manufacture of these splints and for valuable suggestions; to Miss Hope and to my wife for the illustrations.

AN APPLIANCE FOR THE AMBULATORY TREATMENT OF FRACTURES OF THE THIGH.

By C. MAX PAGE, M.S., F.R.C.S.,
CAPTAIN R.A.M.C.(S.R.).

THE importance and value of the ambulatory treatment of fractures of the femur at the earliest period practicable is well recognized. The restoration of full function of the limb; the union of the bone, and the repair of any wounds present are all favourably influenced by such treatment. Hardly less important is the improvement in the moral of the patient which occurs once he is able to move about again independently.

The apparatus which I propose to describe is made and applied by the surgeon from inexpensive and readily available material. Its chief advantage over the calliper type of splint is that at a certain period it leaves the knee-joint free.

DESCRIPTION OF THE APPLIANCE.

The appliance in its simplest form consists of two plaster-of-Paris collars; the upper one surrounds the thigh at the level of the great trochanter, and abuts on the tuber ischii; the lower is moulded on to the two condyles of the femur just above the level of the knee-joint. These two points d'appui are united by two

extension screws, which lie just behind the central coronal plane of the femur.

The extension screws, when adjusted, procure long axis extension of the femur, and pass the weight of the body taken on the foot from just above the knee direct to the pelvis.

The principle employed is similar to that of Hackenbruch's "Distraktion-klammer."¹ The fixation at the knee is that used by Professor Delbet² in his spring splint for the thigh.

As indicated above, the appliance leaves the knee-joint free. If it is used in a case where little or no union is present, fixation of the knee-joint and an extension of the lower point d'appui to the ankle should be effected for a short period.

DETAIL OF MATERIAL.

The upper collar or ischial saddle is made from tarlatan. This material (cost about 3d. a yard) is usually sold 65 cm.

wide. All descriptions below will be based on this size. It is formed as follows (Fig. 1): About 5 metres of tarlatan are folded so as to make a parallelogram 65 cm. by 25 cm., and containing twenty thicknesses of the material. This is then cut as shown in Fig. 1.

The raw edges of the boomerang-shaped section are then roughly tacked. It has merely to be dipped into a plaster emulsion before use.

The lower collar is made from one five-yard quick-drying plaster bandage 3 in. wide.

The extension screws are similar to those used for the wooden type of Macintyre splint. The most useful size for adult thighs measures 10 in. when screwed home, and 14 in. when fully extended. The two screws are attached to arches of perforated zinc by means of split rivets (Fig. 2). The zinc collars are cut for each case. The screws should lie symmetrically both above and below, about half an inch behind the coronal plane of the femur shaft.

Tailed Pieces for Attaching the Extension Unit.—These are made from tarlatan. The one for the ischial attachment measures 110 cm. by 4 cm.; the lower one 90 cm. by

FIG. 1.—Ischial saddle, diagram of tarlatan section required; scale 1/10. It should contain twenty thicknesses of material.

FIG. 2.—Extension unit, consisting of two screws attached to zinc arches by means of split rivets.

FIG. 3.—H tail-piece for attaching extension unit to the collars. The one illustrated measures 90 cm. and is for application to the supracondylar collar. It should contain twelve thicknesses of tarlatan.

4 cm. (Fig. 3). Both should contain twelve thicknesses of material. They are cut like an elongated H, and the margins afterwards roughly tacked.

APPLICATION OF THE APPARATUS.

Preparation of the Skin.—The skin around the ischial area and above the condyles should be rubbed with spirit, dried, and powdered daily, for a short period before the application. The thigh should be shaved.

An anaesthetic is only necessary in the case of very recent fractures.

Position.—The patient is placed (Fig. 4) on the operating table, and the trunk raised by resting it on a padded box. The lower edge of this should extend to about the third piece of the sacrum. The man is held in position by a perineal strut pegged into a prolongation from the deep

FIG. 4.—Position of patient, showing body rest, perineal prop, and method of leg extension during application of the apparatus.

surface of the box. The injured leg is steadied and extended by a weight attached to the foot by a cord which passes through a pulley lying on a bracket at the end of the table. A couple of sandbags support the limb just below the knee.

Application of Collars.

The supracondylar collar is formed by winding a 3-inch plaster bandage, moistened in hot water, around the thigh just above the knee. It is applied evenly and without tension, the upper margin lying about one inch above the upper surface of the condyles. The collar is then moulded above the condyles by the steady pressure of the tips of the fingers of either hand. The grooves so made are kept under moderate pressure till the plaster has set, say five minutes.

The ischial saddle is then applied. The section of tarlatan (Fig. 1) is dipped in a warm plaster emulsion of average consistence, and is then wrapped round the thigh, so that the peak or angle lies a couple of inches above and a little to the outer side of the tuber ischii. The whole is then fixed and reinforced by half a dozen turns of a plaster bandage. A groove is then moulded by steady pressure just below the tuberosity.

FIG. 5.—Front view of ambulatory appliance complete; no fixation of the knee.

screws and over the zinc arches. The apparatus so completed is shown in Fig. 5.

Hardening of the Plaster.

This under usual conditions should be complete in about twenty-four hours. For this time a firm cushion must be

kept under the pelvis in order to prevent deformation of the ischial saddle. Extension through the screws should not be applied during this period; if continuous extension be necessary it should be maintained by weight traction. Posterior support of the fracture can be obtained, if necessary, by passing a broad webbing strap from extension screw to extension screw over a firm pad applied to the exposed posterior surface of the femur.

Fixation of the Knee.

In almost all cases it will be found better to fix the knee in an extended position for the first few days that the patient is up. As soon as the muscles controlling the knee-joint (especially the quadriceps extensor) have recovered some function, knee fixation can be dispensed with, if union of the fracture is moderately advanced. For this temporary

FIG. 6.—Front view of ambulatory appliance with Delbet leg extension.

FIG. 7.—Back view of ambulatory appliance with Delbet extension; posterior support to the thigh is not shown.

fixation any well-moulded back splint will serve. In those cases in which little union is present, and therefore in which considerable and continuous screw extension must be maintained, it is better to fix the knee in an apparatus which extends the purchase of the lower point d'appui. I have found Delbet's method of plaster fixation for the leg the most satisfactory for this purpose. It is a little clumsy, but it leaves the ankle-joint free, and gets a firm purchase on the malleoli and back of the heel. The apparatus applied with this extension is shown in Figs. 6 and 7. The extension is made up from the following tarlatan units:

FIG. 8.—Heel-piece.

FIG. 9.—Tibial band.

FIG. 10.—Side-pieces.

Diagrams of tarlatan units for Delbet's leg extension (introduced primarily for use in fractures of the tibia); scale 1/10. Each section is formed of eighteen thicknesses of the material.

1. The heel-piece (Fig. 8) contains eighteen thicknesses of the material, and is cut from a parallelogram 65 cm. by 15 cm., as shown in the figure, leaving a central heel flap 22 cm. wide, and two tails of the same length and about 4 cm. deep.
2. The tibial band (Fig. 9) measures 65 cm. by 6 cm., and contains eighteen thicknesses of material.
3. Two side-pieces 65 cm. by 9 cm. and of the same density.

4. A many-tailed bandage about 50 cm. square. The tails should be about 6 cm. deep and overlap half of the adjacent ones.

Application of Leg Fixation in Extension of the Supracondylar Collar.

The sections of tarlatan (1, 2, and 3) are dipped in plaster.

The heel-piece is then spread out on the table below the foot, with the tails perpendicular to the axis of the leg. The lower margin of the heel flap is flush with the sole of the foot; the tails will lie just above the level of the ankle.

The tibial band is spread out perpendicular to the leg just below the level of the tibial tubercle.

The side-pieces are then applied. The lower extremity lies flush with the sole of the foot, the upper extremity is incorporated with the supracondylar collar and turned back on itself to reinforce the plaster at the knee level.

Completion of Leg Extension.—While the side-pieces are held in position by an assistant, the tails of the heel-piece are wound firmly around, obliquely (Fig. 7), so as to bind the side-pieces. The tibial band is then wrapped round the leg horizontally, performing a like function above. The many-tailed bandage is then firmly applied over all, so as to mould the plaster on to the bony prominences while the plaster is in course of setting.

AFTER-CARE.

As with all plaster appliances in which pressure over bony parts is exerted, the main difficulty is the avoidance of pressure sores. For this appliance no form of padding is required, but sometimes a flannel lining to the lower part of the supracondylar collar may be used. Sores may appear over (1) the ischium, (2) the condyles, or (3) at the lower margins of the supracondylar collar. Their appearance can be avoided in any of these localities if due care be taken. As a first measure the preliminary hardening of the skin over exposed points by means of friction and the application of spirit is important.

The skin under the ischial saddle can be kept in perfect condition by the regular friction, and introduction of powder, effected by the passage of a "scratcher" of flannel. This consists of a strip of flannel passed between the skin and the plaster in front where there is no pressure. The "scratcher" is powdered, and worked backwards and forwards twice daily round the whole circumference of the ischial saddle. A similar appliance can be used for the supracondylar collar. Additional precautions to be taken at this level are (a) not to squeeze inwards too tightly when the collar is being moulded on to the condyles; (b) to cut away the lower edge of the collar at any place where it tends to dig into the skin. The screws should be left as slack as is safe when the patient is in bed.

LIFE OF THE APPARATUS.

From the point of view of rigidity the collars, if they are set properly, will outlast the period of treatment. They may work too loose if they have been applied to a swollen thigh which decreases rapidly in size. Usually, however, the thigh swells a good deal when the patient first gets up; gradually, as this oedema disappears, the musculature improves, and so the collars maintain their hold. The extension screws may work loose in their bedding if this has not been soundly effected. They can easily be reset on the old collars with fresh H tailpieces. As a general rule, it is desirable to change the ischial collar after four to six weeks' use, if the apparatus is still necessary.

TYPE OF FRACTURES SUITED TO THE APPLIANCE.

Simple Fractures.—Any simple fracture in which long axis extension maintains good alignment may be so treated. Plated fractures come under this category.

Gunshot and other Compound Fractures.—I have chiefly used the apparatus for cases of this kind, and have found it well adapted to fractures in the middle third of the femur. If there is a suppurating wound which would come under the ischial collar the apparatus cannot be used. In the case of compound fractures in the lower third of the femur, and in which the wound would be fouled by the supracondylar collar, the same principle can be applied by replacing the supracondylar collar by a plaster taking its purchase from about the ankle-joint. This involves fixation of the knee.

TIME OF APPLICATION.

The apparatus can be used in simple fractures as early as the eighth day. Its use at such a period involves continuous screw extension and fixation of the knee. In gunshot fractures I have found it most satisfactory when applied six to ten weeks after the injury. In such cases there should be fair union between the fragments in good alignment, and it is only necessary to exert strong screw extension while the patient is up and about. The likelihood of the formation of pressure sores is in this way reduced to a minimum.

WALKING AND CRUTCHES.

The rapidity with which men will walk in this appliance without the aid of crutches or stick depends to a remarkable degree on individual mentality. In general it is wiser not to hurry a man to forsake his crutches. So long as he is taking a fair amount of weight on the injured limb, and moving both the ankle and knee joints, its function will rapidly improve.

SUMMARY.

1. The moral and physical value of early ambulation in cases of fractured femur is great. The shortening of the period of convalescence so produced is at the present time very valuable.
2. The apparatus described can be fitted by any surgeon at a small cost.
3. When union of the fracture is fairly firm this apparatus permits of movement at the knee-joint.
4. The appliance is most easily managed when some union is present, and in general should be applied from six to ten weeks after the injury.

I am indebted to Lieutenant-Colonel S. G. Butler, D.S.O., R.A.M.C., for permission to publish these notes, and to Captain J. N. Fergusson, R.A.M.C.(T.C.) for the preparation of the photographs.

REFERENCES.

- ¹ Hackenbruch, *Muench. med. Woch.*, 1912, lix, p. 1487. ² Delbet and Laman, *Presse médicale*, December 2nd, 1915.

THE DRAINAGE OF SEPTIC KNEE-JOINTS.

BY

W. A. CARLINE, LIEUT.-COLONEL, R.A.M.C.(T.),
LINCOLN.

THE very interesting article on the treatment of septic knee-joints by Colonel Andrew Fullerton in the *JOURNAL* of November 25th, 1916, induces me to give some notes of the methods adopted by me for draining the knee-joint.

Having no notes to refer to I can only recall the earlier cases, which impressed me by their frequency, and one recent case I operated upon shortly before going on two months' sick leave.

Septic knees were more common at this hospital in the early days of the war, and in one small ward seven cases were treated by drainage within a few weeks. Six of these recovered, some with good movement and others with a stiff joint. The case that died had a fracture of one thigh and a bullet wound through the other, the latter being very septic; the knee-joint became infected and was drained, but the patient died of septicaemia. At this time there were no conveniences for bacteriological examination nor for electrolysis with chlorine ions.

In a large number of cases of infected knee-joint it is possible to pass a drainage tube through the joint cavity on the inner side without the tube passing between the bones.

The early cases were washed out with 1 in 20 carbolic acid solution. All the tubes were closed by nipping them, except that engaged by the syringe, and the knee-joint fully distended by the disinfectant; the tube nearest the back of the joint was then freed and the disinfectant and pus allowed to escape, the process being repeated until the disinfectant came away free from pus. This treatment will be ineffectual unless the surgeon personally sees that it is carefully and efficiently carried out, and if he does so there should be no danger of "the entrance of microbes other than the original offenders," as suggested by Colonel Fullerton.

In the early Seventies the late Mr. T. Sympton treated cases of pelvic and other deep-seated abscess at the Lincoln County Hospital by a method then called hyperdistension. The abscess was opened by a valvular incision, and the pus