experience illustrated one of the accidental features which affects the prognosis of abdominal wounds.

This is an interesting fact: it has been our experience that uncomplicated wounds of the small intestine which are suitable for treatment by the operation of suture all recover; the operation is no more serious than an appendectomy. Lately we had a case which showed fifteen perforations throughout the length of the small intestine, but all were suitable for suture, and an uneventful recovery was made.

Stomach wounds offer quite a good prognosis when they are uncomplicated, but unfortunately this last desideratum is very rarely present.

Wounds of the colon, even when uncomplicated, demand a grave prognosis. The gravity arises from the liability which there is of infection of the retrocolic and retroperitoneal tissue.

In uncomplicated wounds of the solid viscera a good prognosis may be given.

But, as we have said in a former paper, there is this fact to be recognized, that one must be prepared for repeated most bitter disappointments; yet, when a series of cases is reviewed, the gains seem infinitely greater than the losses. The prognosis, of course, very largely depends on the degree of the injury sustained; but an even more important factor is the length of time which has elapsed since the injury was sustained. Early operation offers the best and surest chance of ultimate success. In reviewing the statistics of the results of such operations as these it is impossible to consider the question *en masse*. Each individual case must be considered, for the chances of success depend upon so many factors that it varies enormously in different instances.

We wish to acknowledge our indebtedness to Colonel Cuthbert Wallace, A.M.S., Colonel Hewatson, A.M.S., Lieutenant-Colonel Wear, R.A.M.C.(T.), Lieutenant-Colonel Frankau, R.A.M.C.(T.), and to the officers of No. 3 Mobile Laboratory.

ABDOMINAL INJURIES IN A CASUALTY CLEARING STATION.

BY

MAJOR A. DON, F.R.C.S., R.A.M.C.(T.).

In civil life, when a surgeon was confronted with an acute abdomen, his advice, if he saw the case in time, would always have been unhesitatingly "exploratory laparotomy." In fact, that term stood for diagnosis, treatment, and prognosis, all in one. Out here we hesitate. Some of the cases get well under expectant treatment, and it is then never known for certain what, if any, injuries they have had; others die, and even if *post-mortem* examinations reveal the causes of death, one cannot be sure what would have been the result of operative interference.

There seems, however, no reason to depart from our previous pre-war attitude. We have learnt a good deal, and advanced considerably, thanks to the work of Surgeon-General Bowlby, Colonel Wallace, and others in this field. An abdominal injury now means to most of us "see and treat," not "wait and see." Even if the patient be *in extremis*, there is a better chance by operation. There is often none after waiting. Even in the presence of advanced peritonitis an operation with drainage has occasionally succeeded, and the suggestion of high enterocolostomy put forward by Mr. Sampson Handley in the BRITISH MEDICAL JOURNAL (April 8th, 1916) seems a distinct advance on the treatment of peritonitis by drainage alone.

In many cases there is merely a continuous haemorrhage going on, and, when that is so, the chance of recovery, even in the most serious case, is good, once the haemorrhage is arrested. Ether anaesthesia and saline always bucks a patient up a little, and haemorrhage from abdominal injuries is by far the commonest cause of death within the first twenty-four hours. Peritonitis is a later danger, arising from rupture of a hollow viscus, or infection of the effused blood, and except in wounds of the stomach and upper part of the small intestine, where the contents are acid and irritating, peritonitis does not produce at first painful and alarming symptoms or so great shock as does simple haemorrhage. The escape of the contents of the

small intestine lower down, and of the colon, is rare in cases seen early in the casualty clearing stations—say during the first twenty-four hours. It seems as if the bowel were so completely paralysed by the blow caused by the rapid transit of the missile that no movement takes place for at least twenty-four hours, and infection only in the track of the wound is found on opening the abdomen. So slight may be the change in position of the bowel that the injured part is almost with certainty located at once, on opening the abdomen, lying just beneath the wound. Perhaps the morphine usually administered in the field ambulance helps to produce this effect, while the motor ambulance journey from the front may help occasionally to cause escape of bowel contents. Wounds of the stomach, the contents of which are often aseptic, may get sealed up by omentum or otherwise during this period of quiescence, and such wounds may heal without operative treatment if the patient is kept quiet and food withheld, but it is a hazardous chance to take.

INDICATIONS FOR LAPAROTOMY.

The indications for laparotomy would seem to be pain and rigidity of the recti, and marked shock or signs of haemorrhage. When these symptoms are marked, the diagnosis of serious intra-abdominal lesion is certain in nine cases out of ten, and in the tenth or doubtful case no harm will come from exploring. But there are ways and ways of exploring, and one of the most deadly is to evert the whole contents of the cavity all at once. It is an excellent method of finding the damage quickly, but it is equally quick in killing the patient by shock.

Pain varies in degree, but haemorrhage pure and simple into the peritoneal cavity has seemed to cause more pain and greater early mural rigidity than any other condition. It is not easy to explain this almost constant feature. There is practically no immediate peritonitis in pure blood effusion. The intestines, away from the point of injury, are normal in appearance in every case. There are no flakes of lymph, and yet the patient may have been alarming the whole ward by his shouts and moans, and have his recti as tense as boards. The amount of blood in the peritoneal cavity is often extraordinary, and bleeding is generally still going on. This last feature is in part due to the pumping effect of respiration in producing a vacuum at every inspiration. For the abdominal cavity becomes a rigid closed cylinder into which the blood is sucked from the wounded vessels by each upward movement of the piston—the diaphragm. Nature's method of minimizing this pumping effect may be to keep the walls of the cavity taut. Wounds of the liver, colon, and spleen give rise to most speedy haemorrhage, but one may find the abdomen full of blood from rupture of one vessel in the omentum or mesentery. The facial expression and pallid appearance is that of haemorrhage more than of shock or of peritonitis—the well known "facies hippocratica." The abdominal wall, tense and retracted at first, becomes later full and plump, suggesting to the eye an ascitic condition. Dullness in the flanks and fluid in the pelvis may be made out, if searched for. The pulse is small, thready, and rapid. Everything suggests urgency. Nothing is to be gained by delay, and these cases are sent to the operating theatre as soon as it can be got ready. One cannot but deplore the loss of time due to the distance the patient has had to travel, the consequent great loss of blood and the shock of a cold and rough journey from the trenches to the casualty clearing station, and speculate on the advantages to these cases of a similarly equipped station a mile or two behind the firing line.

OPERATION.

In the majority of cases I now prefer to enter by the wound, which is made the centre of the incision. The incision itself goes obliquely downward, in the line of the nerve supply, when the wound is outside the rectus, and vertically downwards when in the area of the recti. An elliptical cut excises the wound first completely down to the peritoneum, and sufficiently wide to give a clean edge. The end of the elliptical incision is prolonged, giving an opening 7 in. to 9 in. towards the exit wound, which may even require to be extended in one or both directions later. I do not hesitate to divide both recti straight across if necessary, and to do this quickly I fix the muscles to their sheaths above and below with safety pins before dividing

them; this prevents retraction, and facilitates suturing later. Such wounds heal well.

As already mentioned, the injured organ will usually be found immediately underneath the entrance wound, and this also gives the direction in which one ought to search for injuries. The small bowel is examined from above downwards or from below upwards, according to the side on which the operator is standing. The assistant examines one side and returns the bowel, while the operator examines the side next him as he pulls the intestine out. Not more than 2 ft. of small bowel should be outside at one time. This minimizes shock considerably, and I am convinced that even where the intestines are covered carefully with hot saline cloths there is greater shock. It may be the contact of a foreign substance or merely of the air, for cloths usually become quickly cooled, and the exposed bowel as quickly becomes red and inflamed and very much distended. The loss of heat must be very great, for the shock is considerable, and there is difficulty and delay in returning the dilated bowel and suturing the wall of the abdomen at the end of the operation. It is curious that organs with no apparent sensory nerve supply cause greatest shock in surgical handling. When a rent in the bowel is found, it is immediately dealt with either by simple closure or by excision. Two clamps are put on just tight enough to prevent slipping, one on either side of the part injured, and with their points meeting on the mesentery (Fig. 1). The injured part of the bowel between

FIG. 1.—The dotted line shows the incision; the clamps are brought together in the direction of the arrows for suture.

the clamps is cut away, leaving an inch of bowel and half an inch of mesentery outside the clamp for suturing. An end-to-end anastomosis is done immediately. The operator requires no assistance for this, and the assistant is free to search for and remedy other injuries. I have not had to remove more than 1½ ft. of small intestine at one place, even where there were multiple rents and bruising. Care must be taken when putting in the inner bowel suture otherwise obstruction will occur, as happened in one of our cases (Case ix). I have seen no advantage in excising wide of the injuries, as the bowel is usually quite normal up to the edge of the rent. One's chief aim ought to be speed and short exposure with efficient workmanship.

The traverse of the missile must not be the only area examined for injuries. Several times in my first cases I had to regret not having examined wider, especially when bone had been hit by the missile on entering. Two cases died on consecutive days from perforation of the small bowel by splinters from bone. In the first case (Case xi) there was a pelvic wound involving the urethra. A small perforation of the peritoneum was noted at the operation, and the pelvic portion of the small intestine was bruised, but apparently not perforated. At an autopsy two small rents were found higher up with a spicule of bone sticking in one of them. There was also general peritonitis. The other case had a wound of the left costal margin, where the spleen and stomach were herniated into the pleural cavity through a rent in the diaphragm. Four perforations were found after death in the jejunum, close together, with general peritonitis. In neither of these cases was there much blood in the peritoneal cavity, but both patients had bled much before operation, and were so bad that probably they would not have survived had an abdominal incision and suture been added to the other treatment which had to be done to stop haemorrhage and repair the bladder and diaphragm.

The concussion force of the missile is sometimes so great that organs like the liver, spleen, and kidneys are

ruptured and even completely destroyed without being in what one might designate the traversed area. I have not found a hollow viscus burst by the centrifugal force or concussion wave from the missile, but I have met this condition in civil practice and recognize its possible occurrence.

After thorough search has been made through the entire abdominal cavity and repair effected, normal saline as hot as can comfortably be borne by the operator's arm, is poured in from a jug by an orderly and the whole abdominal contents thoroughly washed. This should be repeated till the saline is almost colourless. Enough fluid is left in to fill the pelvis and lumbar pools, but not to prevent easy closure of the wound by floating the omentum and intestines into the operator's way. The stimulation from this treatment has always been noted by the anaesthetist as very beneficial to the heart and respiratory organs. It restores a patient who has been causing anxiety almost immediately, and I have never noted bad effects from it subsequently.

Drainage by a split tube and gauze wick is used only where there has been a possibility of leakage of bowel contents, not in cases of pure haemorrhage, and latterly often not even in cases of perforation, where there is no evident soiling. Subsequent peritonitis where haemorrhage has been arrested and all perforations closed has been rare.

AFTER-TREATMENT.

Intravenous salines are given before, during, or after operation, as seems advisable. After the patient is returned to bed warmth and quiet is most essential. The bed is raised at either end, according to circumstances. If the patient is suffering from shock the head is kept low, and if there is a liability to peritonitis and a pelvic drain has been left the head is raised. The greater proportion of the cases are treated in the horizontal position right through, and are allowed to lie in the position of greatest personal comfort. After the effects of the anaesthetic have passed, good results and an absence of thirst may be got from Lieutenant Lawrie's drip method of hypodermoclysis, or saline enemata may be administered at intervals. Feeding by the mouth with tea, beef-tea, and milk is begun as soon as patients recover from the anaesthetic. An egg-flip with the juice of half a lemon and a dessertspoonful of sugar is often appreciated, and is given when thirst is excessive and there has been no high perforation of the alimentary canal. Of drugs, few besides morphine before operation and pituitrin when the bandage begins to tighten, have been in use. Eserine seems to possess no advantages over pituitrin, and I have thought its action less certain. A turpentine enema relieves flatulence if given half an hour after the pituitrin.

These patients may be sent to the base after six or seven days if necessary, but in normal times they are retained at the clearing station till all danger is past. They should be well wrapped up and dispatched to the train with the last ambulance car.

WOUNDS OF THE VARIOUS ORGANS.

By the time these cases reach us there is little to guide one as to the organs injured, and it is just as well, since some surgeons advise leaving alone injuries to particular organs. It is nearly always possible to distinguish severe haemorrhage from simple perforation of gut. The symptoms of haemorrhage are much more severe by the time the cases reach us—generally six to twelve hours. Simple rupture is very apt to escape notice when other injuries of a severe nature are present, and the only way to avoid mistakes is to follow every missile likely to have penetrated the abdomen.

Liver.

Haemorrhage is the chief danger, and many cases reach us in a dying state with the abdomen full of blood, and that from comparatively small tears from rifle bullets. The bleeding is generally from the under surface and is difficult to control by suture. Bile is a post-operative complication up here, and as the worst wounds are drained it escapes along the tube track to the outside. The extent of the damage to the liver bears no relation often to the apparent traverse of the missile, especially in tangential wounds on the right costal margin. The concussion force may destroy half the liver, even where the bullet has not

hit the organ at all, and in one of the cases (Case XII), when the bullet just touched the lower border externally, passing along the anterior and outer side of the chest and fracturing the ninth rib, the whole of the liver to beyond the portal fissure was blown to pieces, leaving a few tags of tissue sticking to the almost uninjured upper capsule. The centrifugal force of the concussion waves was also shown in this case by the right kidney being split into four completely separated pieces, and yet the kidney was two inches or more from the nearest point of the bullet track. Liver tissue is quite easily sutured with strong catgut, the needle being entered after the eye. Where there is a gap in the liver tissue a piece of undetached omentum should be stitched into the gap to control the bleeding and make the grip of the sutures more secure.

Spleen.

Wounds of the spleen are not so numerous. I have seen only five. Haemorrhage would seem to be severe at first, as I have found in all but one of my cases the abdominal cavity full of blood, and very little, if any, active haemorrhage. The omentum has in every case formed adhesions so firm that they surprised me at first. The one with little haemorrhage was herniated into the pleural cavity and the vessels strangulated in the opening of the diaphragm. The spleen is difficult to suture and bleeding from the fresh needle holes is serious. A covering of omentum may be tacked on if the spleen is so little damaged as to be useful. It is better to leave it if possible, as blood repair is much more speedy than when the spleen is removed. In the three cases when I removed it, two died, though one had perforation of the small intestine as well, and the other had a completely destroyed left lung with haemothorax.

Stomach.

Wounds of this viscus are frequent, and are supposed not to be so serious, as it has been suggested that operative interference here is unnecessary. They give rise to great pain, and as undigested food and free gas meets one on opening the peritoneum these injuries are not easily overlooked. The posterior rent may escape notice, and is often difficult to repair. In one case I failed to find the posterior rent even after careful search, and almost absolute certainty of its existence from haemorrhage into the lesser sac. It was found *post mortem* at the cardiac opening quite out of reach. I have not had to excise any part of the viscus nor to do a gastro-enterostomy.

Colon.

I have not met with any wound of the middle part of the transverse colon, though several times there has been much blood effusion, both in the mesocolon and into the walls of the bowel. In one case the mesocolon was ballooned with gas, the piece of shell having passed right through it between the colon and stomach. The liver was damaged, and five days after operation food and bile passed out from the tube track. The ascending colon has been torn in several cases, once completely across. This patient recovered and had the colostomy repaired in Scotland. All the other cases were dealt with by simple suture, and with one exception, in which nothing was done owing to injury to the right kidney and shock, recovered. Only one wound in the descending colon has been treated. The chest and spleen were badly lacerated and drained, and a shrapnel bullet passed per anum with the first motion of the bowels. Rectal wounds are fairly frequent, but seldom fatal if the peritoneum is not opened. They are best treated by a colostomy. My fatal case was wounded behind the left trochanter, and the missile passed between the rectum and bladder and tore three rents in the pelvic small intestine (which were repaired), and passed out through the right pelvic wall, tearing the internal iliac vessels and causing severe haemorrhage, which had not completely stopped on the rectal side before the operation.

Bladder.

Injuries have occurred in four other cases, in one with extensive wounds of the small intestine. No treatment beyond catheterization has been necessary in any of them to allow their removal to the base. Suprapubic drainage would be required for more severe lacerations of the viscus.

Small Intestine.

Injuries to the small bowel are very frequent, and in my experience seldom fatal if got in time and uncomplicated by other gross wounds. Lately we have had only one death in uncomplicated cases. In all cases a circular anastomosis is done as described. In the fatal case the anastomosis was impervious, and, though a lateral anastomosis was done several days later, the patient succumbed to paralytic ileus. This would have been a good case to have tried Handley's operation, while doing the second operation, as vomiting continued more or less to the end (Case IX). There is practically no haemorrhage from simple tears of the small intestine and very little evidence of peritonitis from escape of bowel contents.

Accidents.

Twice there has been hernia of small intestine from giving way of a stitch in cases of through-and-through sutures put in for speed. Reposition under an anaesthetic was easy in both cases and both went to the base. I do not recommend the through-and-through suture, except when the patient is very ill and minutes are of importance.

A very collapsed condition from cold may be avoided if hot-water bottles are put round the patients in the field ambulances and the blankets properly tucked in. The stretcher canvas is no protection in itself from cold, and blankets must be put under as well as over the patients. Mistakes are very unusual, but very detrimental to recovery.

Records of Some Typical Cases.

CASE I.

Pte. O. R. Shell wound to right and below the umbilicus. Abdomen very rigid. Condition fair. Pulse 82, temperature 99.6°, respirations 20.

Operation.—Vertical incision through the wound. No free gas in peritoneal cavity, but some flaky fluid. No rent of viscus discovered. A pelvic drain was put in. Temperature never rose above 100° and pulse 120, on the second evening; the temperature was normal on the sixth day, pulse 72. Sent to base on the eighth day.

This was one of my first cases out here. Very little search was made for injury, and the result might be called fortunate in a case that might have done well without surgical intervention. I trusted to the absence of gas and faecal fluid as being an indication that there was no perforation of the bowel.

CASE II.

Pte. G. P. Wound in left thigh three inches below the trochanter major. The track went upwards in front of the femur towards the abdomen. There was no complaint of abdominal pain, but some rigidity of lower half. He was very much collapsed, and a drain was put into the thigh with some saline packing round about. His temperature at night was 99.8°, pulse 80. Next day he complained of slight abdominal pain and rigidity was more marked, and extended to the level of the umbilicus. Temperature 99°, pulse 84. Next day the temperature was 99.6°, and the pulse 90. Operation by paramedian incision; omentum adherent to the abdominal peritoneum, and some matting. Omentum pulled up, and about two pints of faecal and blood-stained fluid welled out from pelvis with some gas. He died the same evening. Two small rents were found in the pelvic part of the small intestine, one being blocked by a large piece of shell 1 in. square by 3 in. thick.

This was nearly a year ago, when operation on all cases was rather deemed inadvisable. This result, which might have been avoided by operation, led to my operating on nearly all cases showing any abdominal symptoms.

CASE III.

Pte. I. B. Accidental bomb wound to left of middle line in lower abdomen; much collapse.

Operation by incision through wound six hours after the accident. A small amount of blood-stained fluid in pelvis; three small tears in pelvic small intestine. Simple suture and drain left in pelvis; no lavage with saline. Second day, temperature 101.2°, pulse 98, respirations 26. He was in fair condition; drain removed. Temperature and pulse became normal on the fourth day, and he was sent to the base two days later. The condition found here was practically the same intra-abdominally as in Case II.

CASE IV.

Pte. B. K. T. Two wounds from trench-mortar shell thirteen hours before admission. Very collapsed, almost pulseless; abdomen very rigid all over and pain severe. Temperature 97.8° and pulse 130. The wounds were in the lumbar region, the larger just outside the erector spinae swelling and midway between the pelvic brim and the twelfth rib; the lesser on the left side of the third lumbar spine.

Operation.—Incision right paramesial, with centre below the umbilicus. Intravenous saline given during operation, about two pints; the peritoneal cavity was filled with blood; a large rent was found on the outer aspect of the ascending colon 1 in. above the ileo-caecal junction; bleeding still free, and some gas escaped as the colon was being brought up into the wound; no faecal escape. There was much bruising and infiltration of gut near the wound, which was closed in the usual way. Missile not found but probably in the colon. Abdominal cavity washed out with hot normal saline. Drain brought out behind through original wound. The temperature on the third day reached 101.6° and pulse 112. Sent to base on tenth day with no abdominal symptoms.

CASE V.

Pte. W. B. Shell wounds of head, left shoulder, and chest at level of the ninth rib in posterior axillary line. Admitted at 7 a.m.; wounded in early morning; too bad for operation when admitted. Pulse imperceptible at wrist, cold, and collapsed. He was warmed up and wounds dressed at 9 a.m. He had two pints of saline intravenously, and warm saline enema with one ounce of brandy. He recovered somewhat, and was taken to the operating theatre at 2 p.m. for examination of the chest wound. This wound was found to penetrate the chest cavity, to pass through a tear in the diaphragm, and to have torn the spleen. The splenic flexure of the colon was badly bruised and infiltrated but no rent was found. There was much blood both in the pleural and peritoneal cavities, and both were drained through the wound; part of the wound was excised and closed. The wound in the back was explored for ten inches and laid open and excised. A large piece of shell was found at its lower end, and pieces of cloth. The head wound was dressed merely, as he was too ill for further operative treatment. The same evening the pulse was 136, temperature 101.2°, and respirations 30. He improved somewhat next day, but on the third day his pulse was 124, temperature 104°, and respirations 32, and there was a good deal of dirty purulent discharge from all the wounds. Intellect clear. He became very emaciated, but temperature and pulse improved till on the tenth day they were 101.4° and 98 respectively; respirations 28. Pus contained *Staphylococcus aureus* and *albus* and streptococci, and a Gram-negative mobile bacillus. He went to the base on the thirtieth day with wounds almost healed.

CASE VI.

Lieut. F. Admitted 1.30 p.m.; wounded by rifle bullet from sniper at 7.30 a.m. Entrance wound in front of and just below tip of twelfth rib, and exit wound in front 2 in. below tip of ensiform in middle line. He was very collapsed and cold, and bleeding was free from the anterior opening. Taken to operating theatre at once. Incision excising anterior wound. A piece of edge of liver between gall bladder, which was intact, and the round ligament, was torn out, extending 2 in. deep into its substance. Severe haemorrhage on removing clot. This was controlled by a plug of gauze, while the rest of the abdomen was explored. Mesocolon much bruised and inflated by gas. Cavity washed out with hot saline. Liver injury sutured after plugging with omentum. Tube put down to depth of wound. No other wounds were found, but the patient was too bad for prolonged search. Intravenous saline given during operation.

Second day: Had fair night, still collapsed, pulse 130, temperature 102.2°. Third day: Temperature 103°, pulse 126. Blood in urine both yesterday and to-day. Much discharge of blood-stained bilious fluid. Fifth day: Tube removed, and being difficult to replace, was left out. Complained of tightness in evening. Wound opened by dressing forceps and about one pint of bilious fluid escaped; smaller tube inserted. Sixth day: Temperature 101.5°, pulse 116, respirations 28. Some milk given by mouth escaped through the wound. The temperature came down to 99° on the ninth day, and he was sent to the base on the tenth day, when he was able to take a fair meal. He wrote three weeks later from England that the wound had closed and he was feeling pretty fit.

CASE VII.

Pte. C. S. Admitted 7.30 a.m. Shell wound half an inch in diameter over outer edge of right rectus, two inches below the umbilicus. Very cold and collapsed. Pulse 120, temperature 96°. Abdomen rigid and very painful; urine drawn off, one pint, clear.

Operation 9 a.m.—Incision round wound. Much blood, one tear in caecum found. No escape of faeces, but still bleeding—sutured; peritoneal cavity washed out with hot saline; drain into pelvis. Had saline hypodermically by Lawrie's method.

Second day: Improved. Third day: Patient cold and blue. Pulse regular but feeble, 118; temperature subnormal. Colour dark and some yellow staining of conjunctiva; vomiting bilious fluid. Diarrhoea. Fourth day: Distinct improvement, bowels

moved twice. Some suppuration from pelvic end of incision. Fifth day: Is much better, no diarrhoea, abdomen soft. Still having saline. Abdominal wound suppurating top and bottom. He continued to improve, and went to the base on the twelfth day.

CASE VIII.

Pte. I. M. T. Shell wounds of abdomen and leg; wounded the day before. Wound over right ilium 1½ in. below crest and on outer side. Other wound through right calf. Tibia shattered, and much gas and haemorrhage. Very collapsed. Abdomen very rigid. Pulse 116, respirations 38, temperature subnormal.

Operation on Admission.—Paracentral incision on right side. Much blood; two tears, each 2 in. in length, at hepatic flexure, one in ascending and one in beginning of transverse colon. Sutured and drained through right lumbar region. Piece of shell not found, but the patient's condition prohibited a thorough search. Second day: Had a fair night; complains of pain in breathing. Pulse 120, temperature 101.4°, respirations 34. Third day: Bronchopneumonia with pleuritic friction sounds. Pulse 116, temperature 102.2°. Sixth day: Temperature 101°, pulse 108, respirations 30. Leg not doing well. Incised for gas and dressed with hypertonic saline packs. Abdomen soft. No discharge from drain wound. Bowels moving daily. He was sent to the base on the seventh day.

CASE IX.

Shell wounds of abdomen and left hip. Abdominal wound half-way between umbilicus and pubis, and just outside rectus on left side. Second wound large tear above and behind trochanter major through glutei. Collapsed; much external haemorrhage. Wounded seventeen hours before admission. Opening made obliquely through oblique muscles and left rectus. Many rents in small intestine in two places; 1 ft. excised at one place, and end-to-end anastomosis done; 3 in. removed at a second spot three feet from the first. No other perforations found, and piece of shell not found. Wound closed without drain. Did well for three days, then vomiting and other signs of obstruction developed. Abdomen re-opened on fourth day. No peritonitis; obstruction at place where most bowel had been removed. Lateral anastomosis done. He improved slightly after the second operation, but gradually got worse again, with signs of ileus, and died on the eighth day. The wound in the thigh was very septic.

CASE X.

Shell wound of abdomen. Admitted 11.30 p.m.; wounded during afternoon.

Operation 3 a.m.—Pulse 120. Much rigidity and pain. Second wound on right side of chest. Incision included chest wound over ninth costal cartilage and passed down the middle of the left rectus. The missile had passed through the liver. Some haemorrhage. Wounds closed by suture. The peritoneal cavity contained lymph flakes and undigested food. A wound found in the anterior wall of the stomach was sutured; and a wound in the omentum and jejunum was closed. The peritoneal cavity was washed out with hot saline and a drain left in the pelvis.

Second day: Very bad, delirious. Pulse 160, temperature 99.5°. Third day: Has had a better night. Getting rectal salines and hypodermoclysis. Fifth day: Improved and conscious. Wound of skin septic; was strapped. Pulse 90, temperature 99°. Very peevish and thirsty and vomits frequently. Tenth day: Has been much better and retains food. Is anxious to get home to England. Twelfth day: Sent to base. Wrote afterwards that he was doing well.

CASE XI.

Captain E. Admitted 2 a.m.; wound in right groin; soaked with blood. Very collapsed. Pulse 126. Almost moribund from loss of blood. Wound, a round hole 1½ inches across, above Poupart's ligament, exposing iliac vessels on either side. Transfusion with saline. Second wound, entrance (?) in perineum, small slit ½ inch long. Injured previous afternoon.

Operation on admission. Great difficulty in arresting haemorrhage from neck of bladder and obturator vessels and cord, which had been divided by bullet. Testicle removed. Two small openings found in peritoneum. Pelvic small intestine bruised; no perforation. Drain through lower wound, upper wound closed. He did not improve after operation, but lived thirty-six hours. **Post-mortem:** Small hole in caecum, gas escaping, and piece of bone blocking hole in small intestine. General peritonitis in lower half of abdomen.

CASE XII.

Corporal S. M. Wound of right side of chest three inches below nipple. Entrance wound one inch below costal margin on right side; exit wound in right lumbar region. The ninth rib was broken. Almost moribund. Pulse weak and uncountable.

FIG. 2.

Diagrams to show position of wounds: E, entrance; X, exit. Fig. 2 shows wounds in front. Fig. 3 shows wounds behind. Numbers correspond to the number of the case in text.

FIG. 3.

Operation.—Wound excised along whole extent. Tear in peritoneum $1\frac{1}{2}$ in. long. Liver examined and sutured along what seemed a torn margin for 8 in. Tube left in. No injury to hepatic flexure. Further examination negated by condition of patient. Did not rally, and died six hours after operation, in spite of intravenous and rectal salines.

Liver found completely destroyed to beyond the gall bladder. Sutures were holding edges of superior and posterior parts of liver capsule. Right kidney split into four pieces. No injury to bowels.

I wish to acknowledge my indebtedness to Lieutenant Connell for taking notes of these cases, and to Captain W. A. Rees and Lieutenant Lawrie for able assistance at operations and for the after-care of patients.

NOTE.—This paper was written and sent for publication in March, 1916. During the first ten days of July, 1916, 42 cases of abdominal wounds were admitted to the casualty clearing station. Owing to the number of wounded many were necessarily not operated on and the statistics are rather curious. Of 12 cases operated on, 8 recovered; of 30 not operated on, 18 recovered—giving a mortality of $33\frac{1}{2}$ per cent. after operation and 40 per cent. without operation. These statistics alone are very misleading, as all the cases selected for operation had been wounded within twenty-four hours of their admission to the station, while the thirty cases not selected for operation had all been wounded for more than twenty-four hours, and most had been lying out for two or three days. No doubt these were the survivors of many more serious abdominal wounds, most of whom had died before the battlefield could be cleared, and no conclusion can be drawn from such statistics as to the value of operation, though one might say that after the first twenty-four hours the benefit to be gained from operation is questionable.

ON THE INDIAN OPERATION OF COUCHING FOR CATARACT.

BEING A SUMMARY OF HUNTERIAN LECTURES DELIVERED AT THE ROYAL COLLEGE OF SURGEONS OF ENGLAND

BY

LIEUT.-COLONEL ROBERT HENRY ELLIOT, I.M.S.(RET.).

THE operation of couching for cataract was practised and described by Celsus at the very beginning of the Christian era, and many contributions to the literature of the subject have since been made. It is probable that it was introduced into Europe from the East, and it was certainly known and practised in the Orient for many hundreds of years before its first whisper in the annals of Western surgery. From the time of the first advent of the British surgeon to India, couching attracted much notice, but it was not till the work of Lister began to bear fruit that the crude and filthy methods of the coucher excited a due amount of horror in the minds of the exponents of modern surgery. The lecturer said that he had studied the methods of these men for over twenty years and had collected carefully-compiled records of 780 cases of couching; in addition he had been able to obtain fifty-four globes on which the operation had been performed. This latter collection he has now presented to the Museum of the Royal College of Surgeons of England.

There are two distinct operations performed—namely, the anterior, in which the lens is attacked through the cornea or limbus from in front, and the posterior, in which the incision is placed behind the ciliary body, well back in the sclera. The latter is by far the more scientific procedure, two instruments being used, and a deliberate attempt being made to break down the suspensory ligament of the lens, before its dislocation is attempted. Owing to the dirty methods of its exponents, and to the crude instruments used, the results of couching are very bad indeed, only 21.64 per cent. of the operations yielding a visual result equivalent to 1/10 or better. It is estimated that the coucher loses 60 per cent. of eyes which might reasonably be saved by better and cleaner methods of operating for cataract. The principal causes of failure are, in order of frequency, iridocyclitis, glaucoma, and imperfect dislocation. In a few cases these men mistake other conditions—such as glaucoma, optic atrophy, and retinitis—for cataract, and operate on them. The lectures dealt with the

pathological side of the question, and took up a number of subjects in turn.

THE VARIOUS DIRECTIONS IN WHICH THE CATARACTOUS LENS MAY BE DISPLACED.

Dislocation forward was only found four times in all the fifty-four globes examined. In one the lens lay impacted in the angle of the chamber; in one it was fixed between the ciliary body and the sclera, having reached that position by way of the broken-down pectinate ligament; in one the capsule of a Morgagnian cataract blocking the angle of the chamber over a wide area was the only evidence of the lens remains; the nucleus and the contained fluid had escaped.

Dislocation backward was the rule. For convenience sake all such dislocations were classified into a number of groups according to the position of the lens and the greater or less amount of fixation it displayed in its new position.

1. The lens was found floating freely in the vitreous, which was little altered in structure, though presenting slight filmy evidences of inflammatory infiltration.

2. The lens was entangled in a more or less consistent exudate, which occupied the anterior portion of the vitreous chamber and which distinctly limited its movements.

3. The lens was fixed to the back of the iris and ciliary body or to some neighbouring part by definitely organized inflammatory tissue, which represented a farther stage of the process sketched in the previous class. The degree of organization present in the shrunken cone of vitreous, which was usually in continuity with the inflammatory tissue fixing the lens, varied very greatly in different cases. In some, definite fibrous tissue was abundantly in evidence.

4. The lens was matted in a dense mass of cicatricial tissue, lying between the inflamed iris and ciliary body in front and the completely detached retina behind; the detachment of the retina had been produced by the shrinkage of the inflammatory exudate poured into the vitreous cavity. In these cases the septic infection had clearly been of an intense character, and the inflammation set up had been correspondingly severe.

5. The lens, though dislocated backward, lay in front of the anterior hyaloid membrane, and therefore outside the vitreous chamber. The septic infection also appears to have been less intense. These two factors distinguish the globes of this group from those just previously considered, with which in other respects they have distinct affinities.

6. The peripheral cortical remains of the lens lie *in situ* imprisoned in the capsule, and are all that is left of the cataract; they correspond to the after-cataracts met with after extraction.

7. In two cases the lens was thrust in its capsule through the retina, and lay between that membrane and the choroid coat.

ACCIDENTAL INJURIES TO OTHER STRUCTURES THAN THE LENS DURING COUCHING.

Though the object of the coucher is to dislocate the lens, he may accidentally injure any of the other structures of the eye. These are now taken in turn.

The Cornea.—Opaque scars on the cornea are frequently met with clinically, but are hard to recognize in formalin-hardened specimens. In one case a fistula of the cornea was present; it had been associated with anterior staphyloma, which had burst along the track of the original injury, leaving a permanent drain behind. In another case there was a capsulo-corneal synechia, and in a third a retino-corneal synechia.

The Sclera.—Injuries of this coat are common, as the incision is made through it in the posterior operation, but they are hard to discover, and there is an element of chance in their detection. They are best recognised by the pigmentation that follows disturbance of the underlying uveal coat. In one case a scleral fistula persisted as a result of the operation, and gave rise to a filtering cicatrix on its surface. In another case the scleral wound had closed firmly by cicatricial tissue, in which the ciliary body was impacted. Lastly, there was an instance of a wound through the limbus, in which the scar could be followed throughout its course in microscopic sections of the part.