PUNCTIFORM HAEMORRHAGES OF THE BRAIN IN GAS POISONING.*

BY

F. W. MOTT, M.D., LL.D., F.R.S., MAJOR R.A.M.C.(T).

(From the Pathological Laboratory of the London County Council, Maudsley Hospital.)

THE subject of punctiform haemorrhages in the brain in gas poisoning has awakened new interest on account of its existence in fatal cases of shell shock with burial, and in cases of death from inhalation of poisonous gases, either as a result of their liberation by explosives in confined spaces, such as mines, trenches, and dug-outs, or as a result of an offensive gas attack by the enemy. I shall endeavour in this communication to explain the cause of these punctiform haemorrhages.

In 1907 I published in the *Archives of Neurology and Psychiatry* (vol. iii) a paper on "Carbon monoxide and nickel carbonyl poisoning." I came to the conclusion that the nickel carbonyl poisoning was really due to the inhalation of CO employed in the manufacture of nickel. Two such cases occurred of which I had the opportunity of examining the central nervous system, and I found multiple punctate haemorrhages throughout the white matter of the brain, as shown in photomicrographs.

In this paper I compared the naked-eye and microscopic appearance of the central nervous system in these cases of nickel carbonyl poisoning, with those observed in a case of suicide by illuminating gas, and I considered them to be identical in nature. I also reviewed in this paper the clinical symptoms and pathology of CO poisoning, in respect to the findings in the central nervous system, and especially the causes which occasioned the haemorrhages. All three cases died in from four to eight days with the complication of pneumonia. Thrombotic occlusion of cerebral arterioles or venules was considered to be the cause of the haemorrhages; in one case from the nickel carbonyl works an organizing thrombus was found in a vessel of the medulla.

In the case of suicide by illuminating gas, admitted under my care at Charing Cross Hospital, signs of cerebral haemorrhage occurred within twenty hours of commencement of the inhalation of the gas, for the limbs became rigid, and a plantar extensor reflex was obtained. At first the temperature on admission was 99°, but when the rigidity of the limbs and the plantar extensor response was discovered, six hours later, the temperature had risen to 105° and the pulse and respiration had become very rapid.

The nervous symptoms pointed to the occurrence of the punctiform haemorrhages found *post mortem* in the internal capsules, and it may be assumed that the rise of temperature might have been due to the toxæmia coincident with the onset of pneumonia; for when the patient died, on the fourth day, pneumonic consolidation was found. Full notes of the clinical symptoms and *post-mortem* findings were reported. Microscopic investigation showed fatty degeneration of the heart and of the epithelium of the convoluted tubules of the kidney.

Punctiform haemorrhages, attributed to hyaline thrombosis of small vessels of the white matter, have been described by Bignami and Nazari¹ in various diseases; for example, aestivo-autumnal malaria, apoplexy, diplococcal meningitis following pneumonia and measles. It is possible, therefore, that pneumococcal toxæmia was productive of, or associated with, the causation of the haemorrhages in these cases of CO poisoning.

But I am inclined to think that the CO poisoning alone would be capable of causing the punctiform haemorrhages, for the following reasons: In both cases from the nickel works there was evidence of old haemorrhages in the form of minute round or oval punctiform patches of softening, indicative of gas poisoning on some occasion previous to the man being obliged to give up work. And it was legitimate to attribute these symptoms they suffered with—namely, giddiness, vomiting, and headache (migrainous attacks)—to the gas poisoning, causing congestive stasis and haemorrhages. It is well to note that these migrainous attacks are frequently met with in men and officers who have been exposed to those conditions in which CO or

other forms of gas poisoning might have occurred without fatal results.

From the facts observed in these three cases of CO poisoning, combined with certain anatomical conditions of the blood vessels supplying the white matter of the brain, to which I shall now direct your attention, an explanation can be offered why these miliary haemorrhages are found in the white matter of the cerebrum and basal ganglia, and not elsewhere in the brain. It must be recognized that a combination of factors may arise in CO poisoning, namely:

1. The heart, owing to the anoxæmia, has to beat faster and to do more work with less oxygen; consequently it may undergo fatty degeneration.

2. There is microscopic evidence of an irritative and degenerative endothelial change in the cerebral capillaries, as shown by mitosis of the nuclei, and a fatty degeneration, made apparent by osmic acid staining. These changes may be due, as Lancereaux suggested, to CO in the serum, but aggravated by the pneumococcal toxin, which is also responsible for a tendency to increased fibrin formation of the blood, and to thrombosis in those vessels in which the anatomical conditions favour the lodgement of emboli, or clotting of the blood from congestive or inflammatory stasis.

The microscopic appearances of these punctiform haemorrhages in known cases of CO poisoning, terminating in death by pneumonia, are similar to those shown in Fig 1, which is a section of the corpus callosum, showing punctiform haemorrhages, from a case of shell shock with probable CO poisoning.

Miliary Haemorrhages in Cases of Shell Concussion and Gas Poisoning.

I may now mention that the microscopic appearances found in these cases of CO poisoning, dying with pneumonia respectively after four days, eight days, and seven days, were in all respects similar to the appearances presented by sections of brains received from France, notified as dying of shell shock with burial and from gas poisoning—with one exception, and that only differed in the fact that a large part of the haemoglobin had been converted into chocolate-coloured pigment granules, which blocked the small vessels in the haemorrhages.

Before proceeding to the description of these cases I will call attention to the anatomical relations of the vessels of the white matter of the cerebrum, where these haemorrhages are found.

Anatomical Relations of the Vessels Favouring Capillary Stasis.

The pia mater covering the cortex sends delicate walled arteries and veins through the cortex to reach the sub-jacent white matter; the arteries consist of short and long vessels which, after giving off fine branches to the inter-lacing capillary network of the grey matter, terminate in a brush of fine arterioles; the short vessels end in this brush just below the cortex; the long penetrate deeper, to end in the corpus callosum and the centrum ovale. Each little artery breaks up into a tree, and forms a separate system of delicate arterioles. Each arteriole ends in a circumscribed area of capillaries, with an emerging vein. These veins do not anastomose. Thrombosis of arterioles or venules would therefore cause capillary stasis and haemorrhage into the brain substance in a circumscribed area, also escape of blood into the perivascular sheaths of arterioles or venules—a condition generally found to occur where there are punctiform haemorrhages. Owing to the thin character of the walls of the arteries, it is difficult to decide whether a vessel in section is an artery or a vein. Punctiform haemorrhages are also found in great abundance in the brain structures supplied by the perforating arteries, especially those in which the opto-striate and lenticulo-striate branches terminate. These vessels give off relatively few branches until they reach their destination in the basal ganglia and internal and external capsules; they then terminate in a brush of delicate walled arterioles. Each vessel supplies, as in the case of the cortical vessels, circumscribed areas of capillaries, and the result of embolism or thrombosis is the causation of similar small limited areas of haemorrhage and softening, which, when numerous, may become confluent. Stasis is favoured also by the narrow lumen of the capillaries of the brain.

* Read before the Pathological Section of the Royal Society of Medicine, February 13th, 1917.

Shell Shock and CO Poisoning.

The brain of a man said to have died from shell shock was handed to me by Professor Keith for examination. The following notes accompanied this brain:

Fatal case of shell shock with burial, from Captain Armstrong, R.A.M.C., No. 7 Mobile Laboratory, B.E.F. Sent on from No. 1 Mobile Laboratory. No. 8 on Captain Armstrong's list.

Brain of man, admitted unconscious, with history of having been buried by shell blowing in parapet. Remained stertorous for two days and died.

Post-mortem Examination.—There is no wound of any kind on his body or head, and no visceral lesion. His ankle on one side was badly "sprained," but there were no fractures. The skull was unfractured, and no fracture of the base could be found. Brain shows multiple punctiform haemorrhages and some slight subpial extravasation. No other particulars. [Photomicrographs illustrating the appearances presented by the brain in this case have been already published to illustrate my Lettsomian lectures to the Medical Society (1916).]

Having regard to the fact that these punctiform haemorrhages and hyaline thrombosis of vessels were identical in their microscopic appearances to those I had observed in CO poisoning, it occurred to me that the man may have been concussed and afterwards gassed while lying unconscious and buried. (Vide Fig. 1.)

It may be argued that these punctiform haemorrhages were due solely to venous stasis and congestion, but I doubt this, for I have neither observed this condition in the number of cases of death from asphyxia occurring in status epilepticus nor after prolonged seizures of paralytic dementia, although I have examined the brains, macroscopically and microscopically, in a great number of instances.

A letter to me from the Trench War Committee confirmed the possibility of CO poisoning occurring when a large shell burst in a confined space, such as a dug-out or a trench, if incomplete defonation of the explosive occurred. Moreover, it must be remembered that CO is odourless, and may be present in trenches or dug-outs without its existence being known. When a mine is exploded considerable quantities of this gas may be formed, and it may travel through the ground considerable distances.

In the memorandum on *Gas Poisoning in Warfare* issued by the Director-General, Medical Services, British Armies in France, it is stated in respect to CO poisoning:

The lungs show no abnormal changes in cases of rapid death. Small punctate haemorrhages may be found in the white matter of the brain, and sometimes ecchymosis in the meninges, if the case has been exposed to a concentration of CO sufficient to cause prolonged unconsciousness.

The fact that CO is not found in the blood when the patient is examined, does not prove that death was not due to CO poisoning, for after some hours of exposure to air it cannot be detected, and there is little opportunity to make the test for some hours or even days.

Captain Dunn read a very interesting paper at the Medical Society on epidemic nephritis, in which he showed hyaline thrombosis of the vessels of the alveoli of the lungs and of the glomerular capillaries of the kidney. In these cases he has observed multiple punctiform haemorrhages of the brain, which he attributed to embolism by hyaline thrombi. These haemorrhages present exactly the same appearances as in CO poisoning or gas poisoning. In a letter he has written to me, he states that he has now observed these haemorrhages in four more cases of nephritis, so that their occurrence in the first case was not

fortuitous. "They are of quite similar appearance to those I have observed in phosgene poisoning." He asks whether haemorrhages of that type are seen in uraemia.

Major Elliot has been kind enough to send me a copy of a memorandum on five autopsies on cases of poisoning by drift gas ($\text{Cl}_2 + \text{COCl}_2$) in which death took place within seventy-two hours, by Captain H. W. Kaye. In this paper he describes evidence of blood destruction in the spleen. Major Elliot also informs me that Captain H. Henry was the first to describe thrombi in the renal vessels and disagreed with Dunn and McNee when they wished to describe what they saw as renal emboli from the lung.

Examination of the Brain in Gas Poisoning.

I have recently had the opportunity of examining the brains of two cases of gas poisoning, in which gas was employed in an offensive by the enemy; and one is of special interest.

The whole of the white matter is peppered over with small dark spots about the size of a pin's head. These are due to haemorrhages, but microscopic examination shows

conditions which I have not found in CO poisoning, nor in other forms of gas poisoning; in fact, I have never seen any condition like this. The red blood corpuscles have been in large measure broken up, and the haemoglobin converted into dark chocolate-coloured pigment granules, which fill the capillaries, arterioles and venules of the white matter of the brain. This is very possibly methaemoglobin, for it is known that exposure to nitrous fumes in concentration will oxidize the haemoglobin, and convert it into methaemoglobin. Phosgene, COCl_2 , has been used by the Germans; it liberates HCl when it comes in contact with a moist surface; it is very irritating and would cause bronchiolitis, and I am informed by my friend Professor Halliburton that it is possible the free hydrochloric acid would convert the haemoglobin into acid haematin.

Similar appearances

were found to those described in CO poisoning—namely, multiple punctiform haemorrhages in the white matter (Fig. 1)—but the blood corpuscles were intermixed with chocolate-coloured pigment granules (Fig. 2). The accompanying drawings illustrate the appearances presented. You may observe a vessel with a hyaline thrombus, stained pink by the fuchsin of the Van Gieson stain, but with a brownish tinge due to the change in the blood pigment.

Possibly, as in capillary fat embolism, we may have embolism by these pigment granules, but, generally speaking, there is definite evidence of thrombus formation, with pigment granules in the thrombus.

Examination with a high-power magnification shows that the smallest vessels are filled with the pigment, which can be seen in the vessels and in the circumscribed areas of haemorrhage as discrete granules packed together. (Fig. 3.)

The specimens show hyaline thrombus formation in the vessel contained in the centre of the haemorrhage or passing to the centre of the haemorrhage; the walls of these vessels are generally so thin as to support the view that they are veins. Some of the small vessels show aneurysmal dilatation. In one section I observed a leash of small arterioles pressed together by the haemorrhage at the side; on one there was an aneurysm

FIG. 1.—Section of corpus callosum from a case of shell shock, showing punctate haemorrhages. In several of the haemorrhages there is in the centre a small white area of brain tissue with a vessel in the middle of it. Under a high power the small vessel is seen to contain a hyaline thrombus stained pink by Van Gieson method. Magnification, 25 diameters.

FIG. 2.—Small vessel blocked with pigment, haemorrhage and pigment granules. Above it is a vessel, probably a vein, filled with lightly brown-stained hyaline thrombus. Magnification 250.

FIG. 3.—Section of optic thalamus showing a vessel bifurcating, packed with pigment granules so closely that the hyaline thrombus can only be seen in one limb. A punctiform haemorrhage with corpuscles and pigment granules is seen above, and on one side there are three small vessels filled with pigment compressed together. Magnification 350.

FIG. 4.—Three small vessels filled with pigment granules and hyaline thrombus, in the midst of three punctiform haemorrhages found in the optic thalamus. One of these vessels shows under a higher magnification an aneurysmal dilatation. Magnification 80.

FIG. 5.—Section of subcortical white matter from a case of shell shock and burial, possibly accompanied by gas poisoning. A punctiform haemorrhage is seen with vessel and portions of smaller branches filled with hyaline thrombus. Magnification 150.

filled with pink-stained thrombus. Amidst the corpuscles are numbers of pigment granules. Three such haemorrhages with occluded vessels proceeding to them can be seen (vide Fig. 4). Nearly all the punctiform haemorrhages show a central vessel, surrounded by an area of necrosed brain tissue, infiltrated usually with leucocytes. The whole of this area is stained pink by Van Gieson stain, and it is more or less difficult to make out the wall of the central vessel. Sometimes a capillary filled with a thrombus can be seen running to the central vessel. It may be filled with chocolate-coloured pigment granules probably embedded in a coagulum as in Fig. 2, or the coagulum may be of a pinkish-brown colour, due to the coagulum being stained by the pigment dissolved in the serum. In this, as in all other cases, there is evidence of an inflammatory stasis and excess of leucocytes in the vessels, and often into the perivascular sheath and tissues around.

Clinical and Anatomical Notes accompanying this Case.

Brain.—Surface veins, large and small, distended with dark blue clotted blood (veins of base of skull were in same condition). Section shows thickly scattered blue-black dots throughout the brain, especially in the white matter; this applies also to the cerebellum, and to much less extent to pons and medulla.

No haemorrhage seen.

Patient was admitted ten hours after being gassed, and died sixty hours after admission from bronchiolitis and failure of right heart.

It is unfortunate that the clinical and *post-mortem* notes of this case are so scanty, for it is one of great pathological interest. The right heart failure and bronchiolitis, from which the patient died seventy hours after inhalation of the gas, would undoubtedly account for the venous congestion and stasis noted *post-mortem*, and for the thrombosis of the small vessels in the white matter of the brain. The blocking of the capillaries, small arteries, and veins by the chocolate-coloured granules of pigment, especially of the capillaries, would, however, suffice to account for the haemorrhages. In some respects the capillary blockage by pigment resembles the condition found in pernicious malaria, in which disease Bignami and Nazari have described punctate haemorrhage of the white matter of the brain; but I am inclined to believe the principal cause of the haemorrhages is inflammatory stasis and hyaline thrombosis of arterioles, capillaries, and venules, the pigment granules being incorporated in the coagulum.

I received the brain of another case in which the bruises on the body, the haematomata in the right lung, and the other conditions described, all suggest that he had been blown up by a shell and buried, and that the injuries of the brain were due to concussion. The fact that there was no CO detected in the blood does not conclusively prove that he was not exposed, while buried, to CO gas.

Clinical Notes of this Case.

Admitted with diagnosis of shell shock. Purple bruises on arm and leg of right side. Stertorous, unconscious, and during the night before death, constant fits. Lived thirty hours in hospital.

Post-mortem: There were two haematomata in the right lung, but no other visceral injury. No haemorrhage of the scalp, and no fractures of the skull. Some slight subdural haemorrhage of the right hemisphere. Fornix destroyed and full of haemorrhages; haemorrhages also seen in corpus callosum. Haemorrhage in both optic thalami; cerebro-spinal fluid tinged with blood. Men admitted with him said he had been buried by a shell. There was no CO in his blood, and the bruising was purple.

Microscopic Examination.—Multiple punctate haemorrhages are seen; hyaline thrombosis of capillaries, arteries, and venules; perivascular sheaths contain blood. (Fig. 5.) Marked evidence of inflammatory stasis. Some of the small veins are filled with blood corpuscles, one-half of which are polymorphonuclear leucocytes, and in the perivascular sheath and tissues around are large numbers of polymorph leucocytes.

Another case, of which full notes were furnished, was due to poisoning from a gas shell barrage. Unfortunately there were no notes of the condition of the reflexes nor the state of tonus of the muscle of the limbs. The clinical notes do not indicate that this patient suffered with pneumonia, nor of any obstruction to the entrance of air to the lungs; there is no statement regarding the cause of the extremely rapid respiration, but the fact that he was given oxygen and diffusible

stimulant for the first twelve hours suggests that there was great respiratory embarrassment due to deoxygenation of blood. Later there is a definite statement: there is no evidence of cyanosis and no respiratory obstruction; the oxygen was stopped. But the respiration still continued very rapid—50 to 60.

After some days his condition greatly improved, and the respiration fell even to 28. Then on the last day in the morning he suddenly developed grave symptoms, and in the evening it is noted that he developed marked nystagmus, internal strabismus, and the right pupil was distinctly sluggish and slightly larger than the left. The conclusions and findings are not inconsistent with CO poisoning, but it may have been some other gas or mixture of poisonous gases.

I was unable to confirm the statement of haemorrhage into the pons and medulla. The vessels were congested, but no haemorrhage was found. The cerebral hemispheres were badly preserved, and I was only able to examine the cerebral cortex of the frontal lobes.

Microscopic Examination.

Portions of the frontal cortex and subjacent white matter showing to the naked eye miliary punctiform haemorrhages were taken, and, as in all the other cases, blocked in paraffin, and sections cut and stained by Van Gieson and haematoxylin-eosin methods, also with polychrome. The punctiform haemorrhages appeared in some of the sections to form a circle of circumscribed, discrete, oval or round areas of extravasated blood, with often a section of a vessel in the centre or proceeding to the haemorrhagic area (Fig. 5). In sections stained by the Van Gieson stain the lumen and the thin-walled vessels so seen appear a pale pink, and this presumably is due to the contained hyaline thrombus.

In some vessels red blood corpuscles are seen with abundant fibrin formation—a similar appearance to that seen in the alveoli in red hepatization; other vessels appear filled with polymorphonuclears and fibrin. Around the central thrombosed vessels of the haemorrhage are seen deeply stained pink areas of necrosed brain tissue infiltrated with polymorphonuclear leucocytes; very often a vessel can be seen filled with blood and extravasated into the sheath, and occasionally the rupture of a thin-walled vessel causing haemorrhage into the perivascular sheath can be seen. This condition of central thrombosis with necrosis of brain tissue around and infiltration of leucocytes is similar to that observed in the CO poisoning from the nickel works, when the patient lived eight days, and is in accordance with what might be expected, seeing that the man lived six days after possible inhalation of the gas.

SUMMARY.

The reason why these punctiform haemorrhages occur in the white matter of the brain in gas poisoning is primarily due to the anatomical condition of the vessels in the white matter of the cerebrum, where the arteries are terminal; each small artery having a separate capillary system, likewise the emerging veins. A tendency to stasis may be brought about in these separate vascular systems by the failure of the heart as a force pump and suction pump, also by those respiratory conditions which lead to right heart dilatation, and interference with the return of blood from the skull. In most cases the two factors are combined. It seems probable, however, that either factor may act independently in causing inflammatory stasis and thrombosis, resulting in multiple punctiform haemorrhages. In the gas case, in which the haemoglobin has been converted into pigment granules, it seems probable that the haemorrhage may be accounted for by embolic occlusion of the capillaries and arteries. It is unfortunate that, with the exception of the case of CO poisoning by illuminating gas, I have not had the opportunity of examining the organs of the body.

It is quite probable that, as in that case, fatty degeneration of the heart, the kidneys, liver, and vessels of the brain would be found to exist.

In conclusion, I desire to acknowledge my indebtedness to my assistant, Mr. Charles Geary; also to Miss Munro and Miss Watson for the assistance they have afforded me, and to the Board of Control for a grant to enable me to give them an honorarium for their services.