Below the knee the Thomas splint can nearly always be used, except in those cases in which the fracture is near the ankle. Even here it is often possible to use it by the aid of the sole extension as devised by Sinclair.

In the case of fractures of the upper extremity the Thomas splint has not proved so satisfactory, but only for the reason that the straight posture of the arm is unsuited to transport except under special circumstances, as in transit by barge. The form of Thomas splint for the bent arm has not proved a success. For transport the form of internal angular splint, with a hinged back piece for the upper arm as devised by Captain Colin Clarke, is probably the best.

The development of the operative side of the casualty clearing station and the provision of x rays has been of inestimable benefit to the patient. There can be no doubt that the chance of the patient recovering with a good limb and of escaping a long period of suppuration depends on the attention that can be paid to his wound in the first instance. No amount of after-care can ever make up for the want of it at the first moment. A thorough and deliberate operation is all-important. There must be a free opening; the cavity must be explored by the eye, and not only by the finger, otherwise dead tissue and possibly foreign bodies will be passed over.

When first received, the wound is dirty, but the number of pus-producing bacteria is comparatively few. In a few days it is probable, no matter what treatment is adopted, that they will have greatly increased in number. If the first operation has been incomplete, a second may be necessary at the very time that the wound is in the worst possible state, and the procedure necessary to supplement the primary operation may be disastrous in exposing fascial planes to infection from a wound teeming with bacteria.

The early, deliberate and efficient cleansing of the wound is the basis of success, no matter what chemicals are used after it is completed.

REFERENCE.

BRITISH MEDICAL JOURNAL, August 7th, 1915.

PENETRATING WOUNDS OF THE CHEST AT THE CASUALTY CLEARING STATIONS.

BY

COLONEL SIR WILMOT HERRINGHAM, C.B.,
CONSULTING PHYSICIAN.

THE number of chest wounds admitted to the clearing stations is about 2 per cent. of the whole number of wounds admitted. The most favourable cases are those in which a bullet has gone right through the chest; the least favourable, those made by a piece of shell which is retained within the chest.

In cases where the missile has involved both the chest and the abdomen the prognosis is very unfavourable. Occasionally the whole stomach, or part of the intestine, is drawn up into the pleura through a wound of the diaphragm. In other cases the abdominal viscera are so injured as to prevent recovery. In any case the addition of a wound of these organs to the temporary loss of the use of one lung, which is the usual result of a wound of the chest, produces a condition from which very few recover. Injury to the spinal cord is a still more fatal complication.

The following notes are chiefly drawn from a consecutive series of 211 cases, of which careful observations were made and recorded at the time.

The patients are often much collapsed at time of admission, so that in many cases the heart cannot be felt, and in some cannot even be heard for the first twenty-four hours. With warmth, rest, and morphine, they improve greatly by the second day.

Few cases bleed dangerously from the external wound. When this occurs, it can usually be stopped by plugging the wound. There is often much distress if the external wound admits the free entry and exit of air in respiration. It is almost at once relieved if the wound be made airtight with strapping over the dressing. For these large openings into the pleura, in which sometimes three or four ribs are smashed, and infection from the open air would be almost inevitable, a form of procedure has recently been adopted which promises well. After careful paring of the wound, removal of loose bone, and blunting of sharp

edges, the skin and, where possible, the muscles are drawn together and stitched over the aperture, leaving either only a small hole for a drainage tube or none. If a tube is left in, the cavity is then filled with an antiseptic. One surgeon is using emerald green, 1 part to 1,000 of solution of gum tragacanth, for this purpose.

In all but fourteen of the series there were signs of haemothorax. It is rare for the effusion to increase in extent under observation. The chief safeguard against continued haemorrhage is collapse of the lung. In one case which died from repeated external haemorrhage the lung was found adherent to the pleura throughout. Collapse had not in consequence taken place. In another (not in this series) that died with an increase of the haemothorax, by repeated internal haemorrhage, a piece of metal was found lodged in a large pulmonary vessel which it had partially but not completely severed. The retraction of the vessel was thus prevented.

When the effusion is moderate in size, reaching not above the middle of the scapula, nor further forward than the mid-axillary line, it does not cause serious distress. These cases form the great majority. In them by the third day the pulse falls to 84, the respirations to 28, and the temperature will be falling also. Since the observations of Bradford have shown that fresh haemorrhage hardly ever occurs after the lapse of seventy-two hours from the wound, such cases were, as a rule, evacuated at the end of that time. Information from the base showed that this policy was not attended by any bad results.

When the effusion is greater than this, specially if it is complicated by pneumothorax, the patient usually shows distress. The heart is displaced, the pulse is above 100 and the respiration is over 32. Such cases are not fit to travel. They should be aspirated, and about a pint of blood and as much air as possible should be withdrawn. Some were rendered comfortable by this procedure and were able to travel without damage on the fourth or fifth day. It must be noted that the hospital was close to the train, and that though the journey might last even to thirty hours the conditions were comfortable.

It was not thought desirable at so early a stage to remove the fluid completely with the aid of oxygen replacement; that procedure was therefore left for the base hospitals.

In the latter part of the year 1916 a new method of treatment was adopted in a limited number of cases, especially in patients in whom a missile was found by x rays to be retained in the chest. On the second day after the injury, ribs were resected or a costal flap turned back, the pleura opened, and the missile removed. The pleura was then thoroughly washed out, and the whole wound carefully closed. The number of cases so treated is as yet insufficient to enable definite conclusions to be drawn, but experience is so far favourable.

The complications, other than mere size of the haemothorax, which prevented early evacuation, were either septic infection of the effusion on the wounded side, or some disease of the lung on the opposite or unwounded side. In many cases a missile rakes the chest and enters both pleurae. Neither lung can then be called unwounded. The term is confined to cases where one pleura alone has been injured. These complications will now be considered.

In some cases the patient may be comfortable while at rest and have no fever, but on examination there may be the signs of consolidation of the unwounded lung, and movement may produce shortness of breath and some cyanosis. In such cases the condition is that of massive collapse of part, usually at the base, of the unwounded lung. The side is often contracted, the heart is drawn over, and the x rays, if available, show the diaphragm raised and motionless. This condition, familiar after abdominal operations in civil life, was found by Bradford to be a frequent complication of chest wounds. It clears up in about a week.

In other cases there may be an increasing cyanosis and distress, even at rest, for which the condition of the wounded side does not account. On the unwounded side there may be the signs of bronchopneumonia. Some of these are true cases of that disease. But more of them are due to engorgement of the unwounded lung which affects the back and the lower lobe chiefly. The whole of the affected part is solid with blood, and on section presents a glistening surface of dark crimson colour. Its

causation is not clear, but the situation argues failure of the pulmonary circulation.

Unquestionably, however, if the cases be followed through, the most common complication of chest wounds is infection of the haemothorax. When cases are sent down on the fourth day not many infected cases are seen at the clearing station. It is, however, important that the medical officer should be quick to mark its symptoms. The face is pale, though there may be a local flush, the expression anxious, the tongue dry, the appetite bad, and there may be vomiting. The pulse and respiration usually quicken and the temperature rises. In many cases gas formation within the thorax is shown by the alteration of the physical signs and the displacement of the heart. But this does not always occur, nor are any of the other symptoms constant.

Briefly speaking, whenever the medical officer is dissatisfied with the progress of the patient and cannot otherwise explain it, he should always suspect septic infection. If suspicion is aroused, aspiration should be performed at once. Sometimes the fluid will be found to stink, or it may be seen to contain pus, or the froth may remain permanently crimson from haemolysis. Any of these signs is sufficient to indicate that a free opening should immediately be made. If doubt remains, it is well, after drawing off a pint of the effusion, to leave the patient for a day, and, unless he has by then obviously improved, to operate. Bacteriological evidence, when forthcoming, is valuable as a confirmation, but it is on the one hand uncertain, for the infection may be confined to a certain part only of the pleura, and on the other, may cause considerable delay. Clinical evidence should always be trusted, and action taken upon it without hesitation.

Five cases of septic infection occurred before the fifth day after the wound in the present series of cases. In all a free opening was made at the clearing station, and all these patients, though some of them were very ill for a time, recovered sufficiently to be evacuated.

In the whole series there were twenty-two deaths, of which two were due to wounds of the heart, ten to conditions of the pleura and lungs, and ten to wounds of the spine or abdomen. The remainder were evacuated.

THE ADMINISTRATION OF ANAESTHETICS AT THE FRONT.

BY

CAPTAIN GEOFFREY MARSHALL, R.A.M.C.(S.R.).

FROM the point of view of the anaesthetist, wounded men may be divided into three main classes:

1. The lightly wounded.
2. Those suffering from serious wounds with more or less shock and haemorrhage, the two factors being commonly associated.
3. Those suffering from a severe degree of sepsis, especially anaerobic infection.

The choice of anaesthetic depends on which of these classes the patient belongs to, as well as on the region of the body injured.

1. THE LIGHTLY WOUNDED.

These patients are good subjects for anaesthesia, so that the chief desiderata are safety, speed, and convenience. The ideal anaesthetic is one with which induction is rapid, and recovery complete a few minutes after operation, so that the patient is in fit condition for early evacuation by ambulance train.

Gas and oxygen anaesthesia meets these requirements best. With its help a greater number of these cases can be dealt with satisfactorily in a limited time than with any other anaesthetic. When this method is not available, ether should be used. The ether is best administered as a warm vapour by "Shipway's apparatus," as both induction and recovery are more rapid than with the open method.

Local anaesthesia can only be employed in a small number of cases, on account of the multiplicity of wounds and their lacerated and soiled condition.

II. THE SERIOUSLY WOUNDED.

In the more serious cases the one consideration is safety. In other words, we require an anaesthetic which will not be harmful to a patient who is still suffering from the shock of injury, and one which will minimize the shock of operation.

Spinal Anaesthesia.

It has been urged that spinal anaesthesia would meet these requirements, especially in wounds of the legs and thighs, and would therefore be of great value in military surgery. In practice, however, it is found that the intrathecal administration of stovaine has dangers of its own when applied to men whose wounds are recent. In a large

In the charts the continuous line represents blood pressure in millimetres of mercury. The dotted line represents pulse rate per minute.

CHART I.—Spinal anaesthesia. Group A. Stovaine 0.075 gram. Wounds of leg and buttock twenty-one hours. Haemoglobin 85 per cent. Operation conservative. Death twenty-four hours later from gas gangrene.

proportion of these cases the administration is followed by a great fall of blood pressure and symptoms of cerebral anaemia, that is, pallor, vomiting, loss of consciousness, and occasionally convulsions. The syncope is sometimes fatal. It is in the man who has lost blood, and whose wounds are less than forty hours old, that spinal anaesthesia is

CHART II.—Spinal anaesthesia. Group B. Stovaine 0.1 gram with glucose. Wounds of thigh sixteen hours. Haemoglobin 102 per cent. Operation conservative. Recovered.

dangerous. This is shown by an analysis of fifty consecutive cases operated on at a clearing station under stovaine spinal anaesthesia.