velocity, but I have seen quite a number of cases where small pieces of shell or shrapnel have pierced the scalp in an oblique direction, and the fracture to the skull has occurred at some distance from the wound of entrance in the scalp.

One of the last operations I did illustrated this point. The piece of metal lay on the dura, without having torn it, but in getting there it had pierced both outer and inner tables of the skull in the thickest part of the occipital region in an oblique direction, and was found fully an inch away from the entrance of the scalp wound.

A flap avoids all risk of missing a skull fracture, gives an ample field for operation, and the septic wound of entrance can be adequately dealt with.

In closing the wound I make no attempt to tie any arteries of the flap, relying upon accurate apposition and firm stitching.

If the medical and surgical literature of this war is to possess any value, it must, in my view, be based solely upon individual experience. There can be no such thing as a hard-and-fast rule binding our practice. If it were so, surgery would cease to be an art. It would become a trade.

The utterance of pious opinions, unsupported by careful observations, can have no place in these records of facts, the examination and sifting of which may be destined to prove of supreme importance to future generations of soldiers who fall wounded in the fight. Our views as to technique may change, time may modify our practice, but, whatever else may alter, the principle of applying to each case full deliberation and well-considered judgement should "stand like stone."

NOTES ON THE TREATMENT OF HERNIA CEREBRI.

BY

S. SMITH, M.B., B.S., CAPTAIN (TEMP.) R.A.M.C.

DURING eighteen months' experience at a base hospital in France, during which time between six and seven hundred cases of gunshot wound of the head have passed through my hands, one of the most difficult problems which have arisen has been the prevention, or where this has been impossible, the treatment, of hernia cerebri.

The life of the patient is in grave jeopardy as long as this serious complication exists, especially if it shows a tendency to progress, constituting as it does both a warning and a menace.

It warns us of increased and increasing intracranial pressure, and, unless a state of equilibrium be speedily established, the patient will die from compression, causing paralysis of his vital centres. The progressive protrusion of brain matter through a small cranial wound may also produce further or new functional disturbance, not only because the herniated brain is disintegrated, but because the fibres in its neighbourhood are displaced, stretched, and often ruptured. It is not, for instance, rare to find hemiplegia supervene where a hernia develops either in the frontal or post-parietal region.

A hernia, too, especially if large, acts as a direct menace, by reason of the tendency for it to become gravely infected, and for this infection to spread to the underlying brain substance, leading to a generalized cerebritis, or to the spread of infection into the lateral ventricle. The danger of this latter event is all the greater because often there is a diverticulum of the ventricle in the hernia.

Serious infection of the hernia is most likely to occur in those cases where it is strangulated, as so often happens when the bony opening through which it protrudes is too small, the familiar "button" hernia, with its attenuated and constricted pedicle, being thus produced; such strangulation of the protrusion is followed by softening and sloughing, and thus the facility for deep infection is increased.

It has seemed to me that the brain, perhaps because normally it is so effectually covered and protected by its bony envelope, evinces, more than any other tissue of the body, a tendency when exposed to become ulcerated and sloughy, and this should be given great weight when the question of operative measures comes to be considered.

1. Limitation and Protection of Exposed Brain.

It has been our constant endeavour, therefore, to limit as far as possible the amount of brain exposed, making wherever feasible the plastic floor, and draining by rubber tubes passed in at the angles of the flap, a procedure introduced and advocated by Lieutenant-Colonel Sargent, R.A.M.C.¹ The technique of this operation is simple, and in suitable cases it gives excellent results. The larger the dural defect, the larger is the extent of the exposed brain and the greater the risk of infection. Consequently, I consider it of importance, apart from the danger of opening up the highly susceptible subarachnoid space, that the dural opening already produced by the wound should not be enlarged.

Still more important is it not to incise the dura if it be found intact, as this procedure is accompanied by grave risk of infecting the damaged, but not septic, brain underneath, thus leading to the very condition of affairs favourable to the formation of a hernia that we are doing our utmost to prevent.

As a means of diminishing the cerebral exposure we have found suture of the edges of the scalp wound after excision of the edges of the original wound only possible in a relatively small proportion of our cases, the wound generally having been far too extensive and septic for the performance of such a desirable procedure. Moreover, where such a method of closure has seemed practicable, the tension necessarily produced would have made efficient drainage very difficult, and there is an obvious objection to a tight suture line lying pressed up against the exposed brain, which we are trying both to protect and to drain.

2. Antisepsis and Drainage.

It goes almost without saying that another eminently desirable factor in the prevention or limitation of hernia cerebri is the effective treatment of the ever prevalent sepsis, both by the use of efficient drainage with frequent dressings where necessary, and also by the use of antiseptic lotions.

After many experiments with various lotions, I have used latterly a mixture consisting of equal parts of carbolic lotion (1 in 20), hydrogen peroxide (10 vols.), and water, as suggested to me by Mr. Parry Morgan, M.B. (late Captain (temp.) R.A.M.C.). The hydrogen peroxide is specially valuable by reason of its faculty of breaking up and facilitating the removal of sloughs, clots, and disintegrated brain matter that may have blocked up the lumen of the lateral rubber drains, and may so be interfering with the efficient drainage of the underlying brain. This lotion has also been found most useful in the treatment of those cases in which an exposed hernia cerebri already exists. It is very necessary that old dry dressings should be "wetted" off such a hernia and not torn away, for not only does a hernia from which a dressing has been roughly torn evince a tendency to troublesome haemorrhage, which is often difficult to stop, but frequently once the thin outer wall of the hernia is removed there is a marked tendency to the onset of rapid sepsis with the subsequent breaking down of the septic mass.

Before settling down to the routine use of the peroxide-carbolic mixture, I used at one time saline solutions of various strengths, even putting saline tablets rolled up in gauze in direct contact with the brain in certain cases; saline, however, was not very satisfactory, and had a marked tendency to render the brain in contact with it "boggy," with increase in the size of the hernia. Various spirit solutions—among them mercury biniodide in spirit, formalin, and even absolute alcohol—were also tried with a view to drying up the excrescence, but although in many cases the hernia itself dried up satisfactorily enough, the general hardening and fixing of its surface produced by the spirit did not allow the septic foci from bone fragments, etc., at any depth to come to the surface; and although the hernia looked clean and shrunken, the patient frequently died from meningitis or cerebritis extending from such bottled-up foci.

3. Postural Treatment.

It is important to make use of gravity in the treatment of hernia cerebri and also in its prevention, and with this end in view we always, when possible, sit our patients up

as soon as they are sufficiently recovered from the effects of the operation; this procedure is, unfortunately, impracticable in many cases, as, for instance, when the patient is unconscious or very restless, or where some such complication as a fractured femur exists. In those cases, however, in which it can be done, this manoeuvre has a remarkable effect in limiting the hernia, and will even cause it to subside altogether in many cases. It is specially efficacious in herniae of the frontal region or near the vertex, where gravity has full play and the tendency for the brain is to fall away from the wound. It is also important that, by the free administration of morphine, heroin, or similar drug, the patient should be kept quiet and so prevented from bruising his hernia by knocking it against the head-rail—a by no means rare occurrence; with this end in view, too, it is well to keep the hernia, if protruding to any extent, surrounded by a sufficiently large “buffer” of wool to prevent injury.

4. Lumbar Puncture.

In spite of every care, however, hernia cerebri will always be met with in a certain number of cases, both following operation and also in those in which no operation has been performed. It is here that the systematic use of lumbar puncture is of the utmost value.

For many months past we have been treating hernia cerebri by repeated lumbar puncture, using the sitting posture as a valuable auxiliary. Although many cases so treated have died, our results on the whole have been more satisfactory than before these two simple and logical lines of treatment came into general use.

Increase in the intracranial pressure, which is the fundamental factor in the production of hernia cerebri, may be due to:

(a) An actual swelling of the brain substance itself, commonly due in the early stages to traumatic oedema cerebri. This may be succeeded at a later stage by that common and fatal sequel—spreading cerebritis. In these cases, where increase in intracranial pressure is due rather to increase in the actual volume of the brain itself, owing to the oedema and consequent swelling of the individual cells composing it, rather than merely to increased accumulation of cerebro-spinal fluid in the ventricles and subarachnoid spaces, lumbar puncture, by itself appears to do little good. It is a desperate disease requiring a desperate remedy, and often a contralateral subtemporal decompression, combined with repeated lumbar puncture, are the only measures holding out any prospect of success; although there is some risk in conveying infection to the subarachnoid space, some cases thus treated have lived and gone home in a satisfactory condition who must otherwise inevitably have died from cerebral compression.

(b) Haemorrhage, either extra- or intra-cerebral, is rarely sufficient in cases that reach the base to give rise to symptoms; but secondary haemorrhage into the damaged brain, in or beneath the hernia, is not uncommon, especially in cases sent down to us too soon after operation. In these cases a patient, who appears to be progressing satisfactorily, will quite suddenly evince symptoms of increased intracranial pressure, and the accident is evidenced externally by a rapid and enormous increase in the size of the protrusion, from which dark blood and disintegrated brain escape in large quantity. This is specially liable to occur when infection by *B. perfringens* is present. Death is usually too sudden in these cases of secondary haemorrhage into the brain substance to allow of any operative measures.

(c) Some degree of meningitis will cause increase of fluid in the ventricles and subarachnoid space. It is, of course, this third factor in the production of increased intracranial pressure which can often be so effectually treated by repeated lumbar puncture.

Character of Cerebro-spinal Fluid.

The character of the fluid drawn off by lumbar puncture in these cases has been found to fall into one of the following groups:

1. Quite clear though under increased tension, containing no cells and showing no evidence of meningitis. This is probably the type of fluid associated with the early traumatic oedema.
2. Definitely turbid, containing many morphonuclear

and mononuclear cells, often tinged with blood, and showing a stray staphylococcus, streptococcus, or degenerated *B. perfringens* in one or two cells amongst those examined. These I have labelled cases of reactive meningitis, and they are probably associated with a localized or generalized meningitis of low virulence.

3. Purulent, crowded with leucocytes, with many organisms in the field, staphylococci, streptococci, *B. coli* or *B. perfringens* growing on culture. *B. perfringens* is rarely met with as a primary infection of the meninges but is common as a terminal event, and is often super-added to a staphylococcal or streptococcal meningitis towards the end. The explanation appears to be that the *B. perfringens* (*B. welchii*) is of low “serophytic” power, and it is only when the patient’s power of resistance has well nigh vanished that this organism is able to make any real headway.

4. Definitely blood stained, often presenting the appearance of almost pure blood. This is usually evidence of haemorrhage, commonly near the base, and indicates an unfavourable prognosis.

Both in Class 1 and in Class 2 we have had many successes with repeated lumbar puncture; and although, so far, we have not succeeded in saving a case in Class 3, we kept one patient with a virulent streptococcal meningitis alive for nearly a month.

Out of one series of twenty cases with varying degrees of hernia cerebri from which cerebro-spinal fluid has been drawn—

In 5 the fluid was clear (in two cases tinged with blood).

In 9 the fluid was turbid, containing many polymorphonuclear cells and a few stray pyogenic organisms.

In 4 the fluid was frankly purulent (staphylococcus 2 cases, streptococcus 2 cases).

In 2 the fluid was purulent though no organisms grew on culture.

Method of Employing Lumbar Puncture.

In every case, except where the patient is unconscious or dull, general anaesthesia, preferably chloroform, has been used. Our reasons for employing general anaesthesia for such an apparently trivial operation is that usually the patients, during the stage in which we see them, are very restless and extremely sensitive, often arching their backs violently even when touched with the hand, and, in our experience, local anaesthesia with novocain has not been sufficient to overcome this repugnance. Further, such violent straining naturally leads to the protrusion of more brain and the consequent destruction of more functional tissue. In some few cases in the past we found it possible to insert the needle under local anaesthesia at the first visit, but the patient violently resisted any subsequent tapping. It is fortunate in this respect that head cases do not resent the frequent lapses into oblivion necessitated by this procedure as do normal individuals.

Having inserted the needle, and before running off any of the fluid, we estimate the pressure of the cerebro-spinal fluid by means of the simple manometric apparatus described originally by Page and Cassidy.² Percy Sargent has also described its use in connexion with head injuries.¹

The instrument consists simply of about 3 ft. of narrow rubber tubing, connected at one end to a narrow glass barrel funnel some 6 in. in length, and at the other to a small metal fitting with a tap, allowing it to be connected up with the special lumbar puncture needle provided with the apparatus, and which is also fitted with a tap. The tubing and about half way up the glass funnel are now filled with warm normal sterilized saline, and the apparatus fitted on to the needle.

When connected up it will be found—if the needle connect with the intrathecal canal and is not blocked by a plug, as of fibrin—that the saline in the glass funnel will undergo small regular oscillations of an amplitude of from 2 to

5 mm. The glass funnel is then held vertically over the point where the needle enters the patient's back, and moved up and down in a vertical plane until the upper level of the saline column is stationary, save for the small oscillations already referred to. A measurement is then taken with a centimetre scale from the point where the needle enters the back to the upper level of the fluid. Normally, we have found that the column of saline thus supported varies from 15 to 25 cm. The amount of fluid to be withdrawn will roughly depend on this reading, and varies from 4 to 8 fluid drachms, according to the initial pressure recorded. Even where the initial pressure is very high (in one of our cases it amounted to 108.5 cm.) it is not necessary to remove more than from 4 to 8 fluid drachms, as the pressure recorded after the removal of this quantity will rarely register much above the normal. It is important, too, that the fluid be withdrawn slowly, almost drop by drop, as otherwise the hernia, with the too sudden relief of pressure, may sink back through the bony opening into the brain, leaving a deep cavity where before there was a large hernia. This is dangerous, for adhesions may be broken down and a septic meningitis lighted up. Before we realized the danger of this phenomenon, one of our patients, who until then had progressed favourably, suddenly developed fatal meningitis soon after his hernia had involuted in the manner above described—as the result of lumbar puncture where too great a quantity of fluid was removed at too rapid a rate. After removal of the fluid into a sterile test tube for subsequent examination the pressure is again taken and recorded; if still too high, a small amount more fluid may be removed.

The intervals allowed to elapse, between which repetitions of the puncture are necessary, depend on many factors, some of which will now be discussed.

If after the first puncture the hernia shows no sign of decreasing, and compression symptoms do not decrease, it has generally been our practice to repeat the tapping on alternate days, until the pressure as shown by the manometer readings is brought down to within reasonable limits; in a few cases of high pressure with rapidly increasing hernia and progressive signs of compression we have repeated the operation daily, though this, in my experience, is rarely necessary, the performance of the tapping on alternate days giving equally good results. If the fluid is clear, thus showing that there is, in all probability, no inflammatory process at work, it is permissible to wait longer before repeating than if it is turbid, crowded with leucocytes, and, maybe, giving growths of organisms on culture. One cannot assume, however, that clear cerebro-spinal fluid drawn off by lumbar puncture necessarily means absence of meningitis, for more than one of our patients have died in whom lumbar puncture gave clear fluid until the last, though subsequent autopsy revealed extensive suppurative meningitis.

In this connexion it is interesting to note that during the early days of the war I had a case (not a wound of the head) die from what autopsy revealed to be well-marked meningococcal meningitis, with much pus both at the base of the brain and also along the cord, which, when cultured, gave a pure growth of meningococci. Lumbar puncture was performed on the day preceding death, and gave clear fluid, containing a few stray cells, but no organisms visible in films or growing on culture.

The following two cases are illustrative of several treated in our ward by repeated lumbar puncture:

CASE I.

Private A., admitted November 22nd, 1915; suffering from gunshot wound of vertex.

Wounds.—Entry: Dirty wound, 3 in. in length, stretching obliquely across vertex, 6 in. behind nasion. Exit: Larger, more explosive wound, with much septic and disintegrated brain oozing from it, 2 in. to left and 1 in. behind wound of entry.

Condition on Admission.—Dull, difficult to examine, exhibiting weakness of right side of face. His tongue protruded to right. He had paralysis, with slight general rigidity of right arm and hand. Both legs were paralysed, and in that condition of marked rigid extension and adductor spasm typical of wounds near the vertex. Trephined November 23rd; usual flap made, many bony fragments driven into brain were removed, the wounds in his scalp were joined and excised and the flap replaced. The flap was drained by two small rubber tubes, one passed in at each lateral angle of the flap. The scalp was in too septic a condition to allow of the plastic floor being made.

Manometer Readings.

Date.	Clinical Condition at the Time of Lumbar Puncture.	Initial Pressure shown by Manometer.	Quantity and Character of Fluid Removed.	Pressure after Removal of Fluid.
1915. Nov. 29	Slight hernial protrusion through wound	72 cm.	3 vi of slightly turbid fluid removed. No organism. Fair number of polymorphous and mononuclear cells. Sterile on culture	30 cm.
Nov. 30	Slightly better. Protrusion the same	21 cm.	3 vi of slightly blood-stained and turbid fluid, presenting same characters as before	11.5 cm.
Dec. 2	More hernial protrusion	24 cm.	3 v of slightly turbid fluid still containing many cells. Sterile on culture	12.5 cm.
Dec. 3	More protrusion	53 cm.	(No note made.) Fluid removed slightly turbid. Many cells	26 cm.
Dec. 5	Hernia has receded			
Dec. 6	Hernia again protruding	50 cm.	3 vi of slightly blood-stained and turbid fluid removed. No cells apart from blood	17.5 cm.
Dec. 7	Hernia slightly larger	50.5 cm.	3 vi of quite clear fluid removed. Contained no cells	20 cm.
Dec. 13	Hernia, which had receded after last tapping, again increasing	36.5 cm.	3 vi of clear fluid removed	13.5 cm.

No further tapping was required. Patient sent home December 27th, 1915, with hernia flush with skull opening. General condition greatly improved, although his paralysis persisted.

CASE II.

Patient B. was admitted on July 13th, 1915, with a history of having been wounded the day previously.

Wound.—Septic gutter wound 3 in. in length of right parietal region. Much septic brain substance extruded. Patient was trephined on July 14th. There was much fracturing, with comminution of the skull under the wound. A deep track led from the wound deeply into the brain substance. This contained many indriven bone fragments, which were removed. The cavity was then washed out and the track drained by one of Sargent's perforated metal tubes 2 in. in length.

Manometer Readings.

Date.	Clinical Condition and Condition of Wound.	Initial Pressure.	Quantity and Character of Fluid.	Pressure after Removal of Fluid.
1915. July 17	Wound clean. Some protrusion around tube. Patient restless. Temperature 103.8° F.			
July 27	Hernia increasing, tube pushed out by pressure of brain. Pulse remains rapid. Temperature down			
Aug. 2	Very drowsy. Hernia larger and tense	47 cm.	Fluid clear, 3 vij were removed. Contained many polymorphous cells. No organism present	23.4 cm.
Aug. 4	Rigor, followed by cramps in legs after last tapping. Hernia slightly larger. Some tendency for it to break down	41 cm.	3 v of clear fluid removed. No cells or organism present	17 cm.
Aug. 6	Condition slightly better	42 cm.	3 vij of clear fluid removed	12 cm.
Aug. 8	Slightly more protrusion. General condition improved	19 cm.	3 iv of clear fluid removed	7.5 cm.
Aug. 14	Hernia breaking down, and slightly larger	27.5 cm.	3 vi of clear straw-coloured fluid removed	11.5 cm.
Aug. 18	Hernia much smaller. Temperature and pulse normal. General condition much improved. Sent to England			

Both of these two cases were of a type with progressive hernia cerebri, which would, in our experience, have died

if their increased intracranial pressure had remained unrelieved.

The following case is interesting as showing how a patient may drift on from week to week with a streptococcal meningitis.

CASE III.

Private M. was admitted on August 8th, 1915, with a history of having been wounded seven days previously by shell fragments.

Wound.—A dirty gutter wound 2 in. in length immediately behind pinna of left ear.

X-ray examination showed a bone defect of considerable extent under wound, with indriven pieces of bone and many shell fragments lying in the brain substance at varying distances from the wound of entry. The patient was trephined on August 9th, 1915. Fragments of bone were removed from the brain substance underlying the wound. No attempt was made to reach the shell fragments; wound too septic to allow of its being floored in.

Manometer Readings.

Date.	Clinical Condition and Condition of Wound.	Initial Pressure.	Quantity and Character of Fluid Removed.	Pressure after Removal of Fluid.
1915. Aug. 14	Complaining of intense headache. Some hernial protrusion through wound	46.5 cm.	5 vi of turbid fluid removed, containing many leucocytes. Streptococci in pure culture	22 cm.
Aug. 15	Still has intense headache. Temperature high	66.5 cm.	3 x of turbid fluid mixed with blood removed	Not taken.
Aug. 17	High, irregular temperature. Hernia increasing. Headache severe	58 cm.	3 (?) removed. Fluid straw-coloured, slightly turbid. It contained many cells. A few colonies of streptococci on culture	
Aug. 19	—	51 cm.	3 ij (all that could be got to run) of turbid fluid, containing many fibrinous strands, removed. Fluid contained many leucocytes	
Aug. 21	Patient no worse	44.5 cm.	3 v of almost clear straw-coloured fluid removed. Very few colonies of streptococci on culture	
Aug. 26	Semi-conscious. Increasing hernia cerebri	108.5 cm.	3 v removed. Fluid opaque and straw-coloured. It contained a few flakes of dense white fibrinous material	16.5 cm.
Sept. 1	Semi-conscious. More hernia cerebri	65 cm.	3 viij of turbid fluid containing many white shreds	
Sept. 5	Patient unconscious, temperature high, pulse weak. Patient died			

Our comment on this case would be that it is only rarely that a case of streptococcal meningitis lives as long as three weeks. His condition appeared to improve steadily up to August 21st, 1915, when the fluid, which had been turbid, was almost clear, and if only he had been tapped on one or two occasions between this date and August 26th, when the pressure was so high (108.5 cm.), we might conceivably have saved him.

It has seemed to us, after a lengthy experience of these more chronic cases of septic meningitis, that the virulence of the disease, both as manifested by the patient's clinical condition, and also by the number of cells, organisms, etc., in the fluid drawn off by lumbar puncture, shows a tendency to wax and wane, as if, from time to time, owing to the breaking down of adhesions, fresh showers of organisms were conveyed from the local site of infection—that is, the wound—to the meninges at large; just as in the more chronic cases of miliary tuberculosis successive showers are conveyed at varying intervals from the local lesion, it may be a caseating patch in the lung or a broken-down gland, to the blood stream and the various organs of the body.

It is with this factor in view that we should endeavour to secure and maintain efficient external drainage of the wound, even when there is a generalized meningeal infection.

That lumbar puncture, even, is not always without its attendant risks may be illustrated by the following fatal case following its use:

CASE IV.

Private S. was admitted on December 31st, 1915. The wound, a fairly clean punctured wound, was situated in the frontal region.

X-ray examination showed a track leading from the wound of entry down to the occipital region, where lay several shell fragments embedded in the under aspect of the right occipital pole.

An operation was performed on January 3rd, 1916. The wound of entry was cleaned up and pieces of bone removed from the brain substance. No effort was made to remove the shell fragments. The patient was well until January 7th, 1916, when he complained of severe headache. After that he had a steadily rising temperature and some hernial protrusion through the wound of entry. Lumbar puncture was performed under general chloroform anaesthesia. The initial pressure of the fluid was 53 cm. After about 5 fluid drachms of clear fluid had been removed, it began to become increasingly blood-stained, the last drachm containing almost pure blood. Almost immediately the needle had been withdrawn the patient suddenly became intensely cyanosed and his respiratory movements spasmodic, long irregular intervals occurring between each aspiratory effort. In spite of this, however, the pupils remained contracted and the pulse-rate normal and of good volume. Artificial respiration and the inhalation of oxygen produced no effect, and finally, about an hour after the commencement of the attack, he died, the pupils having dilated widely, and the pulse having given out a moment before death.

The autopsy, performed by Captain Henry, R.A.M.C., showed a septic track leading from the wound of entry to the under surface of the right occipital pole, where there lay several large fragments of shell casing. The under surface of the right occipital lobe was bathed in a large quantity of recently-shed blood. Secondary haemorrhage had occurred from the ragged torn end of a small cortical vessel lying on the under surface of the track.

We may reasonably assume that in this case it was the sudden removal of a certain quantity of supporting cerebrospinal fluid that had just been sufficient to start the haemorrhage.

Except for this unfortunate accident, and the other case already described, in which fatal meningitis followed the rapid involution of a hernia after lumbar puncture, we have had no death attributable, either directly or indirectly, to the performance of lumbar puncture.

I am much indebted to Lieutenant-Colonel Sargent, R.A.M.C., and Lieutenant-Colonel Gordon Holmes, R.A.M.C., for their help in the preparation of this paper, which is largely a result of their work. My thanks are also due to Lieutenant-Colonel Ellery, R.A.M.C., Officer Commanding the Hospital, for permission to make use of the notes of cases treated there.

REFERENCES.

¹ Sargent and Holmes, Treatment of Penetrating Wounds of the Skull, *British Journal of Surgery*, vol. iii, No. 11, 1915. ² Proceedings of the Royal Society of Medicine, vol. iv, pt. 1, Clinical Section, p. 56.

TREATMENT OF FRACTURED MANDIBLE
ACCOMPANYING GUNSHOT WOUNDS.

BY

H. P. PICKERILL, M.D., M.D.S. BIRM.,

MAJOR, N.Z.D.C.,

ASSISTANT DIRECTOR DENTAL SERVICES (N.Z.); PROFESSOR OF DENTISTRY AND DEAN OF DENTAL FACULTY, UNIVERSITY OF OTAGO, N.Z.

THE object of this article is to call attention to some methods of treating fractures of the mandible resulting from gunshot injuries, the more so since very few if any of those described in textbooks are suitable.

In a fractured mandible caused by gunshot there is practically always an external wound or wounds, though in not a few cases the bullet seems to have entered through the mouth. The fracture is always compound, and there is more or less loss of bone through comminution or actual carrying away of a portion of the mandible.

The result of such an injury is primarily to exaggerate considerably the typical displacements usually seen in civil practice. Any splint or other remedial treatment should be designed to correct such displacements, and it is necessary to recognize what these are likely to be.

Fractures of the anterior portion of the jaw involving the incisor region lead to (1) narrowing of the arch of the jaw; and (2) twisting inwards of each horizontal ramus, brought about chiefly by the action of the mylohyoid muscle.