THE ORIGIN OF ORTHOPAEDIC SURGERY.

BY

A. ROCYN JONES, M.B., B.S., M.R.C.S.,

SURGICAL REGISTRAR, THE ROYAL NATIONAL ORTHOPAEDIC HOSPITAL,
LONDON.

THERE was a work, in two volumes, entitled, "*Orthopaedia; or, The Art of Correcting and Preventing Deformities in Children*"; translated from the French of M. Andry, Professor of Medicine in the Royal College, and Senior Dean of the Faculty of Physick at Paris." It was printed by A. Millar at Buchanan's Head in London in 1743. M. Andry explains his choice of the term as follows:

As it is not within the reach of every Reader to understand the Title of Orthopaedia, which is affixed to this Book, I shall begin with explaining it: after which I shall give an Account of the Book itself. But in the first place I must take notice of two Pieces which have been published upon a subject akin to this, though they differ a good deal from it; the first in the year 1584, by Scèvele de Saint-Marthe, and the second in the year 1656 by Claude Quillet. As to the Title, I have formed it of two Greek Words—namely, *ὀρθός*, which signifies straight, free from Deformity, and *παῖδιον*, a child. Out of these two Words I have compounded that of Orthopaedia, to express in one Term the Design I propose, which is to teach the different Methods of preventing and correcting the Deformities of Children.

The expression seemed to me the more allowable, that the two celebrated Authors above cited, have made use of Terms of the same kind: the first in giving the Title of Paedotrophia to a Treatise upon the Manner of suckling Infants; and the second that of Callipædia, to a Poem upon the Method of getting beautiful Children: both which Titles are likewise taken from the Greek: the first from *παῖς*, an Infant, and *τροφή*, Nourishment; and the second from *καλός*, beautiful, and *παῖδιον*, a Child.

In a supplement at the end of the second volume mention is made of a series of critical articles, published weekly, by the Abbé Desfontaines, under the title of "Observations upon Modern Writings," and Andry says that, "In this particular one of October 14th, 1741, he (Desfontaines) confines himself to my Orthopaedia." The term "orthopaedia" was, therefore, probably used for the first time in 1741, a view corroborated by William Adams, who states that Andry published his book in Paris in that year.

Hippocrates used bandages for the cure of varus, and the only advance on this method, for centuries, was the employment of mechanical appliances. Orthopaedic surgery can only be said to have really begun with the introduction of tenotomy. The first record of tendon cutting was that performed by Lorenz, on the advice of Thilenius, in 1784. He cut the tendo Achillis through an open wound for talipes equino-varus. The first account of subcutaneous tenotomy is by Petit in 1799, who cut the tendo Achillis for a retraction of the heel after Chopart's operation. In 1816 Delpech, of Montpellier, conceived the idea of treating talipes equinus by cutting the tendo Achillis subcutaneously after first transfixing the leg just in front of the tendon. Owing to a mishap in his first case he did not pursue the operation, although in his work, *De l'Orthomorphie* (Paris, 1829) he lays down very sound rules for the operation and subsequent treatment. Stromeyer, who greatly benefited by Delpech's rules, divided the tendo Achillis by subcutaneous puncture in 1831, and by a series of successful cases established the division of tendons on a secure basis, and ensured its reception as a standard operation in surgery.

Stromeyer published his results in his book on operative orthopaedic surgery (Hanover, 1838). A young English physician in London, W. J. Little, who had talipes equino-varus following infantile paralysis in early childhood, had consulted Sir Astley Cooper and most of the approved surgical authorities in London, but from none did he receive the slightest prospect of cure. In June, 1836, he went to Hanover, and his tendo Achillis was divided by Stromeyer, who later afforded Little the opportunity of practising the operation himself. Going afterwards to Berlin, he taught the operation to Dieffenbach, and treated upwards of thirty patients in that city; Little was therefore the first to introduce subcutaneous surgery into the Prussian capital. Through Johannes Müller's kindness he was also allowed to dissect numerous deformed fetuses contained in the Berlin Museum. In Stromeyer's own words, Little became "the apostle of subcutaneous surgery," for until the conversion of Dieffenbach the German universities regarded the operation with dis-

favour, although afterwards Stromeyer himself was elected professor of surgery at Erlangen.

On his return home Little at once introduced the operation, performing his first tenotomy in London on February 20th, 1837, and a few weeks later attended a discussion at the Royal Medical and Chirurgical Society on this new surgical procedure. The following year he was instrumental in founding the Royal Orthopaedic Hospital, the first public meeting of which was held in 1840 with the Earl of Eldon in the chair. This was the beginning of orthopaedic surgery in this country. In 1839 Little, who was now assistant physician and lecturer on comparative anatomy at the London Hospital, published the first English book on this branch of surgery, *A Treatise on the Nature of Club Foot and Analogous Distortions*, and dedicated it to Sir Astley Cooper, who had displayed a warm interest in what Little had been able to demonstrate. Little continued to write much, and in 1853 published another important work, *On the Nature and Treatment of the Deformities of the Human Frame*. His description of spastic diplegia in infants reveals his power as a clinician, this affection being often referred to as Little's disease.

Of the many workers who have appeared in this field since Little, two stand out prominently—William Adams and Hugh Owen Thomas. William Adams followed closely in Little's steps, and did as much as pre-antiseptic days would permit to advance the subject. He joined the Royal Orthopaedic Hospital as assistant surgeon in 1851, and did much good pathological work. His book, *Club Foot: its Causes, Pathology, and Treatment*, based on his Jacksonian Prize Essay of 1864, was published in 1866. He devised operations for ankylosis at the hip-joint, and for Dupuytren's contraction, and introduced several ingenious corrective appliances. Among other works published by him was one on *The Reparative Process in Human Tendons*, based on post-mortem examinations and experiments on rabbits. John Hunter and Paget had previously studied tendon repair in animals, the former, it will be remembered, being induced to do so through rupturing his tendo Achillis when dancing.

Hugh Owen Thomas of Liverpool was practically a contemporary of Adams. He was a man of extraordinary originality of view. In 1869 he published a well-illustrated monograph on *A New Method in Applying the Wire Ligature in Compound Fractures of the Lower Jaw*, in which he anticipated much of the present day method of dealing with this lesion. His now famous hip and knee splints were figured for the first time in *Diseases of the Hip, Knee, and Ankle* (1875). In this book Thomas adopted the great principle of mechanical and physiological rest, in the treatment of surgical diseases, laid down by John Hilton in his famous lectures at the Royal College of Surgeons, 1860-1862. Hilton very successfully applied his principle to diseases of the hip-joint by means of a splint, which consisted of a leather case from the pelvis to below the knee, with a posterior steel extension and foot-piece to fix the rest of the limb. The Thomas splint was an improvement on this, in that it was simpler and could be variously modified for correction purposes in cases with flexion.

It took many years, however, for Hilton and Thomas to change surgical opinion upon the treatment of joint tuberculosis. Thomas was the first to point out the cardinal principle of relaxing paralysed muscles, and thus preventing over-stretching; he carried it out in practice with his metal splint for dorsiflexion at the wrist, and with his caliper for maintaining extension at the knee.

THE USE AND ABUSE OF BONE GRAFTS.

BY

T. P. McMURRAY, M.B., B.CH., CAPTAIN R.A.M.C.(T.),

SURGEON, NORTHERN HOSPITAL AND INFIRMARY FOR CHILDREN,
LIVERPOOL; MILITARY ORTHOPAEDIC BASE HOSPITAL,
ALDER HAY, LIVERPOOL.

THE operation of bone grafting for non-union of fractures is by no means a new one, and it is one which has undergone many alterations since its inception.

The first bone-grafting operation was done in 1809, when Mercer obtained successful healing of bone plates in the skull after a trephine operation.

A great number of substances, such as solidified milk,