war experience in which we employed almost exclusively and with satisfaction plaster-of-Paris dressings, that, except in the occasional case, equally good, perhaps better, results may be obtained by the use of the Thomas and Jones splints.

The suggestions made in this note are in no way modifications of these splints, but only adaptations which may possibly be found to increase their usefulness.

In fractures of the head or neck of the humerus and in injuries to the bones or muscles of the shoulder girdle we have been impressed with the value of fixation with extension in the abducted position. In our experience there are a surprisingly large number of these injuries. Not only may better alignment and more speedy subsidence of the septic process be gained by the abduction, but the

comfort of the patient seems greater in this position. There can be no doubt that future function is favoured also. As Sir Robert Jones has pointed out, even a stiff shoulder with the arm in at least 60 degrees of abduction provides a surprisingly useful limb. As soon as the septic process begins to subside and the temperature is under control, ambulatory treatment increases the resistance of the patient, and greatly simplifies his care.

It is with these ends in view that these adaptations have been planned. Sufficient trial has been given in the wards and in transport to England to seem to prove their practicability. Many other adaptations may easily be made, indeed have been made. These figures with their legends are merely suggestive and demonstrative of certain forms we have found useful.

A SPLINT FOR THE TREATMENT OF GUNSHOT WOUNDS INVOLVING THE SHOULDER-JOINT.

By J. CAMPBELL, CAPTAIN R.A.M.C.(S.R.).

THE treatment in a base hospital in France of a large number of gunshot wounds involving the shoulder-joint has impressed on me in the first place the great necessity for securing adequate rest and fixation, combined with extension, and in the second place the absence of any useful method for attaining them. Many cases, that came under my care last year after being treated for some time, showed a considerable degree of toxæmia, and not infrequently had the elbow close-in to the side, a bad position for ankylosis owing to the subsequent limitation of movement in the limb. The method of treatment I formerly adopted was merely to bring the arm to the fully abducted position, and secure it there by means of weight extension over a pulley. The abducted limb was slung in place from a Balkan frame placed crossways of the bed, at the level of the shoulders. Care was taken that the patient was never

the uppermost malleable band, is also turned forwards in the same way. To the extremities of these rigid, forwardly curved bands are attached the side pieces of an ordinary Thomas arm splint, in a position of almost complete abduction with very slight forward flexion. The splint is applied in the following way: The patient is placed on his back on the padded frame with the affected shoulder region lying over the gap in the frame, and securely fixed by bending the malleable iron bands to the contour of his body (Fig. 3). He is prevented from slipping up or down on the frame by shoulder and perineal bands fastened as shown. The affected arm is supported by rigid bands of perforated zinc, and fixed extension is applied, not to the forearm, but to the upper arm. In this way the elbow can easily be passively moved and thereby prevented from becoming stiff—a most important consideration in treatment. By passing a bandage, impregnated with French chalk, from side to side between the body and the frame several times a day, any tendency to a sore back is easily overcome.

By the use of this splint I have found that the patient can be easily nursed and the dressings changed without pain or discomfort, and without the general reaction described above. Certainly such cases as I have treated on this frame settled down more rapidly and required less interference than had formerly been the case. Furthermore, patients can be transported in it with ease.

FIG. 1.—Frame, front view.

FIG. 2.—Frame, back view.

FIG. 3.—Frame applied.

moved. The dressings were changed only once in five to ten days after slightly raising the patient's shoulders and pushing the bed from underneath towards the sound side, till the patient was lying on the edge of the bed with the scapular region exposed behind. The dressings could then be done with but little movement; in spite of this, however, there nearly always followed a not inconsiderable rise in temperature and pulse rate, lasting several days.

Accordingly the idea suggested itself that a splint made after the principle of a Jones's abduction frame for the lower limb would prove useful for such cases, and after having used it in several instances I am quite convinced of its efficacy.

The splint consists of an iron frame, as shown (Figs. 1 and 2), covered by a thick, firm pad. The frame and pad are cut away in such a manner as to allow of free access to the greater portion of the scapular region. Three pairs of malleable iron bands, running transversely, embrace the body (a) just above the trochanters, (b) at the level of the lower ribs, and (c) at or just above the level of the nipples. The upper and lower bands are joined together in their respective pairs by straps and buckles. On the side for which the splint is intended the uppermost malleable iron band is replaced by a rounded rod of stouter metal carefully padded—this acts as the point of counter extension. The band of the frame on this same side also is curved round above the shoulder and then turned forwards, and a stout bar, running transversely at the level of

After the period of acute sepsis subsided the cases were put up in plaster with the limb still in the abducted position, but with the elbow flexed, and windows were cut out to allow of access to the wounds. The cases were then sent to England as walking cases.

My thanks are due to Surgeon-General Sir George Makins, K.C.M.G., for the help he kindly gave me in having the splints made at the Base Dépôt Medical Stores, Boulogne.

THE Paris house of Masson announces that it has made arrangements to publish in two annual volumes papers founded on the work done at the Ambulance de l'Océan, La Panne, Belgium. The publication will be directed by Dr. Depage, with the assistance of Drs. A. P. Dustin and G. Debaisieux.

SCHWARZE (*Deut. med. Woch.*, August 2nd, 1917), in some comments on methods of treatment of bacillary dysentery in the field, is rather pessimistic as to the so-called specific treatments, such as antiserum, bismuth, etc. He has had the best results with daily injections of adrenalin up to six doses, combined with two or three injections of atropine. The colic and tenesmus were greatly relieved, and the stypitic action of adrenalin was of value in severe and debilitated cases.