Pyorrhoea alveolaris is often present.
Stomatitis and gingivitis.
Quinsy is a not uncommon complication.
Gastro-enteritis cases have occurred in which, *post mortem*, abundant bacilli and spirochaetes have been found in the intestines (Vincent).
Nephritis and albuminuria. The urine of these patients often contains a trace of albumin, and in some cases acute nephritis may occur. To quote one example:

A doctor working at a base hospital was three weeks in hospital with Vincent's angina, and on recovery was given three weeks' sick leave. Whilst in England it recurred on the same tonsil, the recurrence only lasting a few days. On return to duty he had a third attack on the opposite side, and was admitted to hospital, his urine containing a trace of albumin. Three days after admission he developed acute nephritis. Twelve days later the albumin and casts had entirely disappeared.

Other complications are ulceration of the pharynx, on some occasions causing perforation of the carotid artery (Vincent), bronchitis, laryngitis, with ulceration of the vocal cords, pleurisy, empyema, otitis media, cachexia, endocarditis.

A case of gangrene of the vulva and perineum caused by *B. fusiformis* and Vincent's spirochaete in large numbers has been recorded by Spillmann.

An association of Vincent's microbes complicating a syphilitic chancre of a phagedaenic tendency has been noted by Laimois, and the same association has been observed in soft chancre. A case of thrombosis of the left saphenous vein with gangrenous ulceration of the leg, complicating Vincent's angina, has been recorded by Guinsberg.

I have seen one case in which the uvula was the only part attacked, leading to complete destruction of the posterior portion and tip, a very small part only remaining. Osteomyelitis has been observed by Gilberti, and pernicious anaemia by Mayer.

Symptoms.

In an uncomplicated case the symptoms are sore throat, headache, and pain behind the eyes; there may be no symptoms after the first day or two except slight malaise. In the more severe cases there may be loss of appetite, wasting, and great prostration.

Treatment.

Drugs which have proved of benefit are calomel in repeated small doses, but Vincent states that mercurial treatment aggravates the condition when the patient is not a syphilitic.

Arsenic and a mixture of sodium salicylate and potassium chlorate have been used, and tonics locally. Mercury perchloride and glycerin (1 in 500), silver nitrate, methylene blue, neo-salvarsan, solution or powder applied daily for three days, and frequent gargles of hot hydrogen peroxide have been recommended.

The treatment which Vincent himself finds most effective is a thorough painting with tincture of iodine 6 per cent., after well scrubbing with a tampon to remove the membrane.

A dose of antidiphtherial serum has in some cases proved beneficial. The importance of treating the unaffected side should be borne in mind to prevent occurrence on that side. Salvarsan intravenously (one dose) has been said to modify the disease and prevent recurrence.

The condition must be diagnosed from diphtheria, follicular and other forms of tonsillitis, syphilitic ulceration, primary and tertiary, and scarlet fever. In every case a culture should be made to establish the absence of Loeffler's bacillus.

The Wassermann reaction in Vincent's angina is negative. Certain skin conditions, such as erythematata and bullous eruptions, have been reported, but are probably due to coincident streptococcal infection.

Deductions.

Vincent's angina is a more serious complaint than is often supposed; its general effect on the system is often severe, probably from abundant absorption of toxins. Recurrence may take place even three weeks after the first attack, in which case complications are more liable to occur.

REPORT ON SIX CASES OF TONSILLECTOMY IN DIPHTHERIA CARRIERS.

BY

C. C. BALLANTYNE, M.B.TOR., CAPTAIN C.A.M.C.,

AND

B. S. CORNELL, M.B.TOR., LIEUTENANT C.A.M.C.

DURING June and July, 1917, about fifty cases of diphtheria and diphtheria carriers were treated at Moore Barracks Canadian Hospital, Shorncliffe. The so-called carriers particularly presented a problem in that persistent treatment failed to eliminate the germ. Accordingly, tonsillectomy was resorted to in some of these carriers, and a report of six cases follows. While no originality is claimed for the treatment, the results may perhaps prove interesting.

The tonsils were completely enucleated by the dissection and snare method, and swabs were at once taken from the oral surfaces of the removed tonsils. Then, after thoroughly cleansing the tonsils externally with alcohol and ether, cross sections of the gland were made at three different levels with sterile knives. Swabs were at once taken from the crypts at each of these different levels, the first being one-eighth of an inch beneath the oral surface, the second at the centre of the tonsil, and the third one-eighth of an inch from the capsule. Hypertrophied adenoids, when present, were also removed, sectioned and swabbed in a similar manner. The swabs were immediately planted on Loeffler's blood serum media, incubated for twelve hours, and the resulting growths examined for the *Bacillus diphtheriae*.

CASE I.

Pte. W., admitted suffering from peritonsillar abscess, which readily subsided after incision. On May 24th, as a result of routine swabbing, he was found to be a diphtheria carrier. Until June 12th swabs were invariably positive, and on that date tonsillectomy was performed. Swabs from the tonsils showed:

Right: Surface, positive; top section, middle, and bottom, negative.

Left: Same as right.

Adenoids: Surface and section, positive.

Subsequent throat swabs: Three successive negatives on June 15th, 18th, and 20th.

CASE II.

Pte. M., admitted as a carrier; for six weeks numerous swabs were repeatedly positive. Tonsillectomy on June 12th.

Right: Surface and top section, positive; middle and bottom section, negative.

Left: Surface, top section, and middle, positive; bottom section, negative.

Subsequent throat swabs: Successive negatives on June 15th, 18th, 20th.

CASE III.

Pte. S., admitted with mild tonsillitis, which readily subsided in forty-eight hours without antitoxin. Routine swab June 3rd positive for diphtheria, and repeated subsequent swabs positive. Tonsillectomy June 12th. Swabs from tonsils showed:

Right: Surface, top section, and bottom section, positive; middle section, negative.

Left: Positive throughout.

Three successive negative swabs were not obtained until June 26th, 27th, and 28th.

CASE IV.

Pte. T., admitted June 17th as a carrier. Swabs persistently positive. No constitutional reaction. Tonsillectomy on June 29th. Swabs from tonsils showed:—Right: Positive throughout. Left: Positive throughout. Three successive negatives obtained July 6th, 7th, 9th.

CASE V.

Pte. O'H., admitted May 25th as a diphtheria carrier. Swabs invariably positive. Tonsillectomy June 21st. Swabs from tonsils showed both tonsils strongly positive throughout. Three successive negatives obtained June 23rd, 24th, and 25th.

CASE VI.

Pte. Z., admitted June 5th with mild diphtheria, which readily responded to 12,000 units of antitoxin. Swabs positive persistently till June 29th. Tonsillectomy June 29th. Swabs from tonsils showed both tonsils strongly positive throughout. Successive negatives obtained July 3rd, 4th, and 5th.

In these six cases the operation obviously accomplished the object for which it was undertaken. Although in some the organisms persisted for a few days after operation they disappeared with the healing of the wound. In all these cases saline irrigations and gargles had been continuously used until the time of operation, and their failure is explained by the pathological findings, since in four of the six cases the bacilli were found in the very

depths of the crypts, and surface applications could not be expected to reach them.

Living cultures of *Staphylococcus pyogenes albus* had also been employed as sprays over the tonsils in the hope of outgrowing the diphtheria bacillus, and it was usually found that swabs taken next day were negative for Klebs-Loeffler organisms, but were loaded with staphylococci. The bacilli, however, reappeared in large numbers within a day or two. The probable explanation is that the large number of staphylococci collected on the first swabs had outgrown the bacilli on the media. If, however, the staphylococci outgrew the bacilli on the tonsils, then fresh organisms were soon resupplied from the tonsillar crypts.

CONCLUSIONS.

1. That in apparently normal individuals the *Bacillus diphtheriae* may be harboured in the depths of the tonsillar crypts.
2. That complete enucleation of the tonsils seems to be a successful means for eliminating diphtheria organisms from carriers, although it appears too drastic a measure for routine adoption.

PNEUMOTHORAX WITH HERNIA INTO LEFT PLEURAL CAVITY.

BY

T. H. SANDERSON-WELLS, M.D.LOND., M.R.C.S.ENG.,
L.R.C.P.LOND.,
WEYMOUTH.

THE following case seems sufficiently rare to merit publication:

The patient, aged 26, was admitted to the Princess Christian Hospital, Weymouth, in the late evening of August 21st, complaining of urgent and persistent vomiting. There was a previous history of indigestion for some years; he had never been jaundiced.

The patient was wounded in the chest ten months previously. It was apparently a perforating gunshot wound, followed by the spitting of blood, and fracturing the eighth, ninth, tenth, and eleventh ribs in the left mid-axillary line. The wound was quite healed and sound. The patient had apparently had previous attacks, lasting as much as six hours, but they had become milder during the last three months.

The present attack followed violent exercise on August 17th. Complaint was made of pain in the region of the pylorus; it became very acute forty-eight hours later and had continued ever since. Vomiting occurred thirty-five minutes after the onset of the pain and persisted. The vomit at first was green, but soon became black, with an unpleasant, but not fetid, odour. It recurred during the time he was watched every four minutes in amounts of a few ounces. There had been constipation since the onset of the pain, but he passed flatus on August 20th. He attributed his condition to a diet of fish, but no solid food was taken for forty-eight hours; water swallowed returned almost immediately as black fluid.

On admission he was very collapsed and nearly pulseless, and Dr. Quackenbos, who handled him throughout with the greatest care and skill, administered 10 minims of adrenalin hypodermically. An enema led to the passage of a small quantity of flatus, but no faeces.

The abdomen looked extraordinarily empty, and had an appearance of scaphoid retraction. There was no tenderness, but a little lump to be felt over the pylorus, which was tender and which eventually proved to be an enlarged lobe of the liver. The liver dullness appeared, however, to be normal. Reflexes were present.

There were obvious physical signs of a pneumothorax. The heart's apex beat was visible beneath the right nipple; dullness was present on the left side up to the fourth rib; breath sounds absent; tinkling fluid splash sounds. The patient, whose condition was very bad, was given saline, $\frac{1}{10}$ grain of atropine, and 10 minims of adrenalin.

Operation.

A median incision was made above the umbilicus, and the abdomen explored. The bowel felt, and appeared to be, completely collapsed, and much of the abdominal contents absent. The stomach could not be found, the great omentum and transverse colon had also disappeared. The liver was pushed downwards, and presented in the wound. Further exploration with the hand traced the stomach, omentum, and colon up towards the left half of the diaphragm, and through a hole at the back of the diaphragm into the left pleural cavity. Reduction could not be effected, and the diaphragm was incised from behind forwards, the hole enlarged, and the fingers passed through into the pleural cavity; firm adhesions were discovered between the pleura and its contents, and reduction was obviously impossible. The abdominal wound was quickly closed and the left thorax opened by an osteoplastic resection of the ninth and tenth ribs on the left side in the mid axillary line. Adherent

omentum covered the pleura over the wound and had to be tied off; then through the opening collapsed colon and an enormously dilated stomach presented. A Paul's tube was inserted into the stomach, and $1\frac{1}{2}$ to 2 pints of black fluid drained off. The tube was packed round with gauze, and the skin held in position with a few stitches. The patient was removed to bed, his condition apparently better than when he was brought down. His breathing was much easier, and he seemed very comfortable. He became restless, however, and, despite the exhibition of all usual remedies, he died in the small hours of the morning.

A partial investigation was made the following day by Dr. Quackenbos, who reports as follows:

The heart was situated behind the sternum, the apex being in the parasternal line on the right. An aperture was disclosed in the body of the left diaphragm, which had been enlarged by incision, but which appeared to be of some standing. The liver was displaced downwards, and to the left. The left lung was collapsed, and showed evidence of bronchopneumonia and adhesions at the apex. The stomach was very much enlarged, and, with the transverse colon, the great omentum and four feet of the jejunum, was in the left thorax. The omentum was firmly adherent to the aperture in the diaphragm and elsewhere to the pleura. There were adhesions of the gut to the pleura in places.

The explanation of the condition is difficult. The old-standing appearance of the hole in the diaphragm might even suggest a congenital weakness, but it seems more likely that the perforating bullet wound which broke the ribs injured the diaphragm, and that the negative pressure in the thorax drew up abdominal contents to take the place of collapsed lung.

Whether, had the patient's condition been better, it might conceivably have been possible to separate the adhesions and reduce the hernia it is difficult to say, but in any case the future function of the organs would not have been satisfactory.

A CASE OF ECTOPIA VISCERUM.

BY

P. R. COOPER, M.D., B.Sc., F.R.C.S.,
BOWDON, CHESHIRE.

THIS case is of sufficient rarity to merit a brief description, especially as the question of surgical intervention arose.

A male child, 16 hours old, was brought to the Altrincham Infirmary by Dr. Young of Broadheath. An umbilical sac or cyst, larger than the abdomen itself, was attached at the umbilicus by a broad pedicle. Its covering was translucent and composed of apparently the same gelatinoid material as the umbilical cord, with which it was incorporated. The sac was fairly tense and contained some darkish fluid, but the wall was not sufficiently transparent to allow of the contents being identified. The child was apparently full time, but not quite of normal size. It was markedly cyanosed and sluggish in its movements, but it evinced no signs of bodily discomfort. The head and limbs were well shaped and apparently normal. The thorax was, however, very narrow and poorly developed. There was very little abdominal wall apart from the umbilical protrusion. The testes were undescended, but the penis and scrotum were well formed.

It was doubted very much whether the condition could be remedied by surgical means, but as the child had practically no chance of survival without surgical intervention, and as the possibility of such intervention could not be ascertained without exploring the contents of the sac, it was decided to operate immediately. Dr. Young gave the anaesthetic, and the abdomen was prepared as for laparotomy. The outer wall of the sac (that is, amnion) was pinched up with forceps, snipped, and a quantity of blood-stained watery fluid released; this reduced the size of the sac quite one-half, and a second covering was exposed by extending the opening in the outer layer. Both layers were extremely thin, membranaceous, and non-vascular, except just at the lower part, where the umbilical vessels proceeded to the stump of the umbilical cord.

On incising the inner covering (? peritoneum) the large dark solid mass of the liver at once presented; this was situated entirely outside the abdomen proper. The intestines were also exposed, and were for the most part entirely outside the abdomen proper. The opening of the abdominal wall proper—that is, where the gelatinoid material of the sac abruptly joined the normal skin—formed a ring of about 2 in. diameter, and through this an attempt was made to reduce the extruded viscera. With great difficulty the intestine was put back, but it was found quite impossible to replace the liver. There was not only no room for it inside the abdomen, but there was not enough abdominal wall to have covered it even had one been able to reduce it. All that could be done was to reapply Nature's inadequate covering of the viscera, stitch it in position, and dress the parts aseptically. The child lived for twenty-seven hours after the operation. It had two actions of the bowels, but so far as we know did not pass water. A little fluid was given by the mouth, but vomiting occurred shortly before death, which presumably resulted from cardiac failure.