some instances. These pulmonary lesions probably possess some significance in relation to the frequent occurrence of dyspnoea.

For the supply of clinical records and pathological material we must thank collectively a very large number of officers throughout the armies in France.

We are especially grateful to Surgeon-General W. W. Pike, who placed all the records of cases of nephritis in his army at our disposal, and to Colonel Sir Wilmot Herringham, who has all along taken the greatest interest in the work and given us much valuable assistance and advice.

ON THE HAEMAGGLUTININ REACTION AS A TEST OF THE TOXICITY OF VARIOUS ANTISEPTIC REAGENTS,

AND ON THE ASSOCIATION (IF ANY) BETWEEN THE HAEM-
AGGLUTININ CONTENT OF THE BLOOD SERUM AND
PUS, AND CAPACITY TO RESIST INFECTION.

(With Special Plate.)

BY

C. J. BOND, C.M.G., F.R.C.S., COLONEL A.M.S.,

HONORARY CONSULTING SURGEON, NORTHERN COMMAND; HONORARY
CONSULTING SURGEON TO THE LEICESTER ROYAL INFIRMARY;
MEMBER OF THE MEDICAL RESEARCH COMMITTEE.

It is known that different individuals vary in the capacity of their blood serum to agglutinate the red blood cells of other individuals and of animals of different species. It is supposed that somewhere about 80 to 90 per cent., possibly more, of all individuals possess blood serums containing one or other of the different haemagglutinins. The following inquiry was undertaken to test (by utilizing the agglutinative reaction) the toxicity of various antiseptic reagents to the blood serum and the red corpuscles, and to ascertain whether patients whose blood serum contained any of these haemagglutinins resist infection better than patients with non-agglutinative serums.

During a previous investigation into the influence of antiseptics on the activities of leucocytes and the healing of wounds¹ it became necessary to examine the blood serum in order to test the toxicity of different serums to my own leucocytes during incubation. This examination, while revealing facts of interest concerning the action of different serums on the white blood cells, also confirmed the previously known fact of the existence of iso-agglutinins in the serums of a considerable number of patients taken from a hospital population. Fortunately several different patients were found possessing blood serums of four distinct haemagglutinative types (Fig. I), namely:

1. A serum agglutinative to human (C. J. B.) red cells but non-agglutinative to sheep's red corpuscles.
2. A serum agglutinative to human (C. J. B.) red cells and also to sheep's red cells.
3. A serum non-agglutinative to human (C. J. B.) and to sheep's red cells.
4. A serum agglutinative to human (C. J. B.), sheep's and guinea-pig's red cells.

Without entering into the wide problem of the relative

The Medical Research Committee have provided us with some of the apparatus necessary for histological work, and we are indebted to them also for permission to use the drawings made by Sergeant A. K. Maxwell, R.A.M.C.

REFERENCES.

- ¹ Sir John Rose Bradford, *Quarterly Journal of Medicine*, vol. ix, No. 34, January, 1916, and *Journal of the Royal Army Medical Corps*, October, 1916.
- ² Langdon Brown, *Lancet*, February 19th, 1916.
- ³ Abercrombie, *Journal of the Royal Army Medical Corps*, August, 1916.
- ⁴ Thornley, *BRITISH MEDICAL JOURNAL*, December 16th, 1916.
- ⁵ Cf. Aschoff, *Pathologische Anatomie*, Jena, 1913, and McCallum, *Textbook of Pathology*, Philadelphia, 1916.

incidence of these different kinds of agglutinative serums in man and in the animal species it is evident that such a sharply defined divergence in agglutinative capacity affords a means (when employed under controlled conditions) of investigating (a) the relative effect of different antiseptic reagents on agglutination, and (b) the value, if any, of the iso-agglutinative factor in the production of acquired immunity against infection.

The Iso-agglutinative Reaction as a Test of Antiseptic Toxicity.

While the bactericidal capacity of an antiseptic reagent is of the greatest significance, it is becoming increasingly evident that the relative toxicity of different reagents to the blood and tissue cells is also a matter of equal importance.

From the theoretical point of view it is desirable to test this toxicity in regard to (a) the white corpuscles, (b) the red corpuscles, (c) the tissue cells, (d) the plasma and the blood serum. I propose, however, in this communication to refer only to the effect of different antiseptics on the capacity of any given blood serum or pus to clump human (my own) washed red corpuscles and those of the sheep and guinea-pig, and to refer to the power of my own leucocytes and the living phagocytes in different samples of pus to ingest red corpuscles, so agglutinated or opsonized.

Experiment 1.—Six separate drops of a serum agglutinative to C. J. B. red cells were placed on a glass plate in horizontal series (see upper row, Fig. II). Six corresponding drops from a non-agglutinative serum form the lower row. To each separate drop of serum in both rows one drop of an antiseptic reagent was added in the following order:

- (1) 1 drop 0.9 per cent. saline.
- (2) 1 drop 5 per cent. hypertonic saline.
- (3) 1 drop 4 per cent. chloramine-T solution.
- (4) 1 drop 2.5 per cent. carbolic acid solution.
- (5) 1 drop 0.1 per cent. mercury biniodide solution.
- (6) 1 drop 0.1 per cent. acriflavine solution.

The reagent was first mixed with the serum, and then to each drop was also added a measured quantity of human (C. J. B.) red corpuscles previously washed in normal saline.

The corpuscles were then thoroughly mixed with the antisepticized serum in each drop.

This experiment shows that when certain antiseptics are added to blood serum the agglutinative effect on the red cells varies according as the serum is or is not an agglutinative serum. Thus in the upper row (an agglutinative serum) the addition of normal saline shows the normal agglutination and serves as a control. The hypertonic saline, chloramine, and carbolic acid give no

DESCRIPTION OF SPECIAL PLATE.

COLONEL C. J. BOND.

FIG. I.—Showing types of haemagglutinating serums. *Blood Serum W*: Non-agglutinative to human (C. J. B.) and sheep's cells. *Blood Serum X*: Agglutinative to human (C. J. B.) and sheep's red cells. *Blood Serum Y*: Agglutinative to human (C. J. B.), non-agglutinative to sheep's red cells.

FIG. II.—Showing the action of antiseptics on an agglutinating serum (upper row), and on a non-agglutinating serum (lower row). 1, Normal saline; 2, 5 per cent. saline; 3, chloramine-T 4 per cent.; 4, carbolic acid 2.5 per cent.; 5, mercury biniodide 0.1 per cent.; 6, acriflavine 0.1 per cent.

FIG. III.—Showing the action of antiseptics on human red corpuscles (C. J. B.) without blood serum. 1, Normal saline; 2, 5 per cent. saline; 3, chloramine-T 4 per cent.; 4, carbolic acid 2.5 per cent.; 5, mercury biniodide 0.1 per cent.; 6, acriflavine 0.1 per cent.; 7, iodine 0.1 per cent.

FIG. IV.—Showing the reagglutination of red cells by agglutinating and non-agglutinating serum (upper row), after previous treatment by an agglutinating serum and the removal of the agglutinins by washing in saline. In the lower row the first agglutination was produced by acriflavine.

FIG. V.—Showing the effect of removal (by saturation) of the specific haemagglutinin, for human and sheep's red cells respectively, from a serum (5) agglutinative to both human (C. J. B.) and sheep's red cells. Upper row: A, after saturation with sheep's cells. Lower row: B, after saturation with human cells.

FIG. VI.—Showing the effect of different antiseptics on a heated agglutinating serum (upper row), and on a heated non-agglutinating serum (lower row). 1, Normal saline; 2, 5 per cent. saline; 3, chloramine-T 4 per cent.; 4, carbolic acid 2.5 per cent.; 5, mercury biniodide 0.1 per cent.; 6, acriflavine 0.1 per cent.; 7, iodine 0.1 per cent.

FIG. VII.—Showing the action of different antiseptics on an agglutinating liquor puris (upper row), and on a non-agglutinating liquor puris (lower row). 1, Normal saline; 2, 5 per cent. saline; 3, mercury biniodide 0.1 per cent.; 4, chloramine-T 2 per cent.; 5, iodine 0.1 per cent.; 6, acriflavine 0.1 per cent.

agglutination; mercury biniodide, acriflavine, and iodine (in a less degree) all give a well-marked agglutination.

In the case of the non-agglutinating serum in the lower row treated with the same reagents none of the drops show any agglutinative change.

This shows that it is essential to ascertain the agglutinative capacity of any serum before using it as a test of the action of antiseptic reagents. In the case of the hypertonic saline a transient initial agglutination occurs with an agglutinating serum, but this soon disappears. Chloramine, carbolic acid (in the strengths used) both cause destruction of the red cells and laking of the haemoglobin. This also occurs with all the other antiseptics after a lapse of time with the exception of the hypertonic saline.

Before drawing any inference it is necessary to check this observation with a similar series showing the effect of the same antiseptics on washed red cells without the addition of any blood serum. Fig. III shows the result. To each drop of different antiseptic solution a measured quantity of washed and sedimented (C. J. B.) red cells was added in the following order:

- (1) Normal saline.
- (2) 5 per cent. saline.
- (3) Chloramine 4 per cent.
- (4) Carbolic acid $\frac{1}{2}$ per cent.
- (5) Mercury biniodide 0.1 per cent.
- (6) Acriflavine 0.1 per cent.
- (7) Iodine 0.1 per cent.

None of the reagents in the strengths used, except acriflavine, cause agglutination. Iodine 0.2 per cent. also does so; but it must not be inferred from this that acriflavine and iodine are more toxic to red cells than the other antiseptics. As a matter of fact they are less toxic, and the absence of agglutination in the case of chloramine and carbolic acid is due to the destructive action of these reagents on the red cells.

The Capacity for Reagglutination shown by Red Cells which have been Previously Agglutinated by Serum or by Antiseptic Reagents.

There is also another way in which we can test the action of antiseptics on red corpuscles. A measured quantity of washed human (C. J. B.) red cells was added to known volumes of the different antiseptic solutions to be tested. Corpuscles and solution were thoroughly mixed and allowed to stand together for half an hour. The mixture was then centrifugalized and the sedimented corpuscles washed free of all antiseptic admixture in several changes of normal saline. The result of submitting red cells previously so treated to the action of an agglutinating and a non-agglutinating serum respectively seems to vary somewhat according to the antiseptic used, and whether it does, like flavine, or does not, like chloramine, itself produce an initial agglutination, and whether it does or does not destroy the red cells. In general terms we find that red cells which have been treated with antiseptic reagents (which do not destroy the corpuscles or lake the solution) do, after washing in saline, reagglutinate when mixed with an agglutinating serum. This is true of normal saline, hypertonic saline, acriflavine, and hydrogen peroxide. In some cases the effect of the antiseptic seems to be to sensitize the corpuscles and render them more susceptible to reagglutination. Thus red cells agglutinated by acriflavine in some specimens reagglutinate when mixed with a non-agglutinating serum. It is of great interest to find (see Fig. IV) that this is also true of red cells previously agglutinated by contact with an agglutinating serum without any antiseptic. The experiment also shows that (as far as haemagglutinins are concerned) it is possible to wash away the agglutinating substances from the surfaces of the corpuscles, leaving them again susceptible, and in some cases more susceptible, to the influence of a second dose of agglutinating serum. Bearing in mind the very close analogy between the haemagglutinins and the bacterial agglutinins it is probable therefore that this is true also of the latter bodies. The serum used to agglutinate and sensitize the red cells was also tested after centrifugalization and was found to have lost all its agglutinative content.²

The Effect of Antiseptics on Heated Serums.

We have seen that the effect of an antiseptic on the agglutinative capacity of a blood serum depends on the

inherent capacity of that serum to agglutinate red cells. But it also depends on whether the serum is in the normal condition or whether it has been heated. We know that the haemagglutinins are thermostable at a temperature of 56° C., so that the difference in result cannot be simply due to destruction of haemagglutinins. Fig. VI shows the effect of different antiseptics on a heated agglutinating serum (top row) and on a heated non-agglutinating serum (lower row).

The effect of heating for half an hour at a temperature of 56° C. on an agglutinating serum is to increase agglutination. It now occurs with carbolic acid and hypertonic saline when red cells are added, and even with chloramine an initial slight clumping takes place. In the case of the non-agglutinating serum (lower row) agglutination is absent in all the drops.

In order further to analyse the effect of heating an agglutinating serum, the influence of acriflavine and iodine, when added to the serum in different degrees of dilution, was examined in detail. Dilutions in normal saline of a non-agglutinating serum in strengths of 1 in 10, 1 in 15, 1 in 20, 1 in 40, 1 in 80, 1 in 100, 1 in 200, 1 in 300, 1 in 400 were made. The point was then found at which the addition of an equal volume of this diluted non-agglutinating serum neutralized the agglutinative effect of a drop of acriflavine 0.1 per cent. and iodine 0.1 per cent. respectively when mixed with the serum. This point was found to lie just below 1 in 300 in the case of the iodine solution, and 1 in 400 in the acriflavine solution. Evidently it only requires an extremely small quantity of a non-agglutinating serum to prevent the agglutinating action of these antiseptics.

It is interesting to consider for a moment how this inhibition may be brought about. Unlike serum, which acts specifically on the red cells, the antiseptic acts non-specifically both on the serum and on the red cells, combining with protein elements in both. If there are available proteins in the serum, other than agglutinins, capable of combining with the unsatisfied element in the antiseptic, then the corpuscles may escape injury. If, on the other hand, there are no available proteins in the serum, or if they have been destroyed by heat, then the full effect of the antiseptic will fall upon the red cells. Antiseptic reagents vary in the avidity with which they combine with serum proteins, and thus in their toxicity to the red cells.

An attempt was also made to ascertain the amount of non-agglutinating serum that must be added to an agglutinating serum in order to neutralize or inhibit its agglutinative power. An agglutinating serum was diluted four times with a non-agglutinating serum. At this stage of dilution, although it agglutinated red cells, the reaction was incomplete. Clumping occurred on gentle movement, but on standing the agglutination disappeared, to reappear again on movement, just as a freshly formed precipitate may redissolve in excess of the reagent used.

The Specific Character of the Different Haemagglutinins.

It is known that an agglutinating serum may contain different haemagglutinins, each separately capable of clumping the red cells of another individual or those of a certain species of animal. The following experiment shows that it is possible to remove one kind of haemagglutinin from a serum without removing the others. Two separate measured volumes were taken of a serum agglutinative to human and sheep's red cells; one volume (a) was shaken up with washed sheep's red cells, and the other volume (b) with washed human (C. J. B.) red cells. Both were centrifuged, and the clear serum tested for agglutinative capacity. Volume (a) was found to have entirely lost the power of agglutinating sheep's red cells while still clumping human corpuscles, while volume (b) had retained its capacity to agglutinate sheep's cells but had lost the power to agglutinate human cells. As in the case of the bacterial agglutinins, therefore, the haemagglutinins are specific for different varieties of red cells, and the same serum may contain a number of haemagglutinins. Fig. V shows the result of this experiment.

The Action of Haemagglutinating Serums on "Whole" Foreign Blood—"Drop and Pool" Method.

So far we have only dealt with the influence of different antiseptics in favouring or inhibiting the tendency of red

blood corpuscles to clump together under suitable conditions. Before we draw any inferences as to the toxic action of these reagents on the serum, the red corpuscles or the leucocytes, it is necessary to ascertain what action different serums of different agglutinative capacity exercise on the whole blood of another individual (C. J. B.) without the addition of any antiseptic (Fig. VIII).

FIG. VIII.

One or more drops of the serum to be tested are placed within a vaseline or plasticine cell. Into the centre of the little pool of serum so enclosed a small drop of blood from the finger is allowed to fall; the cell is closed by a cover-slip. After incubation the cover-slip and cell wall of this emigrating chamber are removed, the blood clot and serum are gently washed away with normal saline, and the film of emigrated leucocytes fixed and stained with 1 per cent. iodine solution. The extent to which the toxic or agglutinative action of the serum affects the activities of the leucocytes from the periphery towards the centre of the drop of blood affords a rough quantitative test of the toxic capacity of the serum by which the drop of blood is bathed, and the extent to which the foreign serum has neutralized the normal serum of the blood tested.

If the foreign serum used be an iso-haemagglutinating serum then the (C. J. B.) red blood cells clump, but a further change also takes place. The white blood cells which have emigrated from the clot on to the slide will not only be covered with attached clumps of red cells, but some of the leucocytes will contain ingested red corpuscles. The agglutination of the (C. J. B.) red cells occurs in relation to their own (C. J. B.) leucocytes, and the adhesiveness is mutual, not only between individual red cells, but also between red cells and the surface of the leucocytes.³ This clumping and ingestion of red cells by native leucocytes in the presence of a foreign iso-haemagglutinating serum throws light on an important biological problem, namely, the relation of the agglutinins to the opsonins, and will be referred to later. Meanwhile, another fact of importance must be mentioned. A given serum may exercise a powerful agglutinative effect on the red cells and yet be relatively innocuous to the white blood cells. These, when incubated with such a serum, may emigrate and adhere to the slide in large numbers, and exhibit a considerable capacity for elaborating iodophil substances.⁴ On the other hand, a non-agglutinating serum may prevent emigration and the formation of iodophil substance. It may poison the leucocytes and yet fail to agglutinate the red cells. In fact, the leucotoxic effect and the haemagglutinative effect of a serum may vary independently.

Haemagglutinins in Pus.

Samples of pus have been collected from a considerable number of wounds dressed with normal saline in order to avoid the complicating effect of antiseptics. The pus has been centrifuged, and a drop of the liquor puris placed on a glass slide and a given volume of washed red cells, human, sheep's, or guinea-pig's added. The result has been to show that in most cases the liquor puris agrees in agglutinative content with the blood serum of the patient. If this is haemagglutinative, then the liquor puris is haemagglutinative also. Not only is this true of the effect of the liquor puris on human red cells, but it is true also of the red cells of different species of animals.

The Effect of Heat on Liquor Puris.

The effect of heating liquor puris for half an hour at 56° C. is, to a certain extent, the same as in the case of the corresponding blood serum. An agglutinative liquor puris retains its agglutinative capacity, while a non-agglutinating liquor puris remains non-agglutinating.

The Action of Antiseptics on Liquor Puris.

Not only does the purulent discharge from a wound faithfully reproduce the agglutinative characteristics of the blood serum of the patient, but it continues to do this, within limits, independently of the kind of infection present in the wound, or the kind of antiseptic reagent with which

the wound is dressed. It is evident that this retention of agglutinative capacity in a fluid like pus affords a means of ascertaining the degree of decomposition going on in the discharge and the extent to which its vitality has been affected by the growth of micro-organisms in the wound or by antiseptic dressings. I have been surprised at the tenacity with which an agglutinating pus from a heavily infected wound will retain its power of agglutinating human red cells. Even pus recovered from a piece of gauze dressing soaked in normal saline which had lain on a granulating wound for twelve hours yielded a liquor puris agglutinative to human red cells; but this will not be the case, of course, unless the patient's serum is also haemagglutinative. These observations are of interest as confirming the fact, not only that the purulent discharge from most wounds contains many living pus cells, but also that, from the humoral point of view, pus may be sufficiently living to contain agglutinins.

Fig. VII shows the action of different antiseptics on an agglutinative liquor puris (upper row) and on a non-agglutinative liquor puris (lower row). The reagents added to the liquor puris are the same as those used in the case of the blood serum, and there is a striking similarity in the reaction in both cases. In the upper row agglutination occurs with normal saline, mercury, and acriflavine, and slightly with iodine, but no agglutination is seen in any of the drops similarly treated in the lower row in which the liquor puris is itself non-agglutinative.

Summary of the Action of Antiseptics on the Cellular and Humoral Elements of Blood and Pus.

1. On the red blood corpuscles. It is, I think, clear that we may divide antiseptic reagents into two groups: (a) Those which agglutinate red blood cells without destruction of the corpuscles; (b) those which haemolyze the red cells without producing agglutination. Acriflavine and 5 per cent. saline are examples of the first group, and to a less degree mercury biniodide and iodine, though these produce a certain amount of laking. Chloramine and carbolic acid belong to the second group. The test of the absence of haemolysis is the capacity of previously agglutinated red cells to reagglutinate, after thorough washing, when again mixed with an agglutinating serum. Acriflavine and hydrogen peroxide seem to favour this result.

2. On the leucocytes and pus cells. It is useful to distinguish between those reagents which inhibit emigration of leucocytes from the blood clot when incubated with the whole blood in a closed cell and those which allow of free emigration and the elaboration of iodophil substances by the phagocytes. Judged by this test, acriflavine in strengths below 1 in 2,000 in normal saline seems to be only slightly injurious to the leucocytes when incubated with whole blood. An interesting fact has been observed when serum or liquor puris has been incubated with acriflavine in 1 in 2,000 concentration in normal saline. Any leucocytes or pus cells or organisms present in the serum seem to become, when incubated with acriflavine in serum, peculiarly sensitive to iodine when this reagent is subsequently added in 1 per cent. solution. The combination of serum, or liquor puris, acriflavine, and iodine together brings about a blue-black staining of the leucocyte granules. The organisms also show up as black bodies when ingested by the leucocytes, and the blue-black colour of the granules suggests the colour reaction formed when iodine is added to certain carbohydrate substances. With regard to pus cells, capacity to ingest pigment granules and to elaborate iodophil substances in the presence of antiseptic reagents affords a working test of vitality, and this has been used as the test of leuco-toxicity in the antiseptics used. The test has been applied in two ways: (1) *In vivo*, by the addition of sterilized starch grains or pigment granules to the antiseptic used for dressing the wound. The number of pus cells which contain starch grains or pigment affords evidence of phagocytic activity in the presence of the antiseptic in the wound. (2) *In vitro*, pus cells from the wound have been incubated with the antiseptic and the starch grains in a closed cell. Judged by this test acriflavine compares favourably with mercury, carbolic acid, chloramine, and hypertonic saline.

3. The action of antiseptics on the blood serum. The inhibition of the capacity of an agglutinating serum to agglutinate washed red cells provides a test of disintegrative action on the serum by the antiseptic employed. Here

also reagents can be divided into two groups—(a) those which destroy the haemagglutinins in the blood serum, and (b) those which do not. Chloramine, carbolic acid, and hypertonic saline belong to the former, and acriflavine, hydrogen peroxide (in certain concentrations), and iodine to a less degree, to the latter.

The Relation of the Haemagglutinins to the Bacterial Agglutinins.

There does not appear to be any essential connexion between these two constituents of the blood serum. From a small number of observations kindly made for me at the 5th Northern Base Hospital by Dr. Agnes Porter we find that a serum of high haemagglutinative capacity may have very little power of agglutinating certain varieties of bacteria and cocci, and vice versa.

The Relation (if any) between Haemagglutinin Content and Leuco-toxicity of a Blood Serum.

Reference has already been made to the fact that different serums (from healthy persons) vary greatly in their action on the white blood corpuscles of another individual when these are incubated with such a foreign serum, and the question arose whether any association could be traced between high haemagglutinative content and marked leuco-toxicity, or vice versa. As far as my observations go, I have not been able to trace any such association. It is probable, therefore, that leuco-toxicity is associated with the presence of other bodies in the serum and not the haemagglutinins.

The Relation of the Haemagglutinins to the Opsonins.

I have already spoken of the mutual reaction which takes place between leucocytes (or pus cells) and red corpuscles when the latter agglutinate in the presence of the former. Many samples of pus from infected wounds have been centrifuged. If the liquor puris proved to be agglutinative to my own or to sheep's red cells, some of the pus cells after thorough washing in saline have been incubated in a closed cell with human (C. J. B.) or sheep's washed red cells. As a general rule, ingestion of red cells does not take place under these conditions by thoroughly washed pus cells or by washed leucocytes; but if a drop of the agglutinating liquor puris be added before incubation the pus cells will take up the red cells. If ingestion is about to take place the red cells clump and adhere to the leucocytes as well as to each other. I have, however, found that thoroughly washed pus cells from an agglutinative pus will sometimes ingest sheep's red cells in the absence of any liquor puris. This suggests that sheep's cells may require less opsonizing than human cells, or that the pus cells elaborate fresh haemagglutinins.

A further point also requires notice. When red cells are added to healthy pus from a patient whose blood serum contains haemagglutinins, not only do the red cells clump together and adhere to the pus cells, but the pus cells also agglutinate in masses. This can be seen by mixing a drop of such pus on a slide with normal saline. Washed red cells are then mixed with the thinned pus. After a short incubation period a gentle rocking movement of the slide will bring about not only a clumping of the red cells but also a massing of the leucocytes in groups.

The effect of agglutinative liquor puris has also been tested on whole blood by the "drop and pool" method. Liquor puris obtained by centrifuging pus from a wound in a patient with a haemagglutinating blood serum is placed in a closed cell. Into the centre of the pool of liquor puris a drop of blood (C. J. B.) from the finger is allowed to fall and the cell closed and incubated. On removing the cell walls and gently washing away the central blood clot the opsonizing effect of the liquor puris on the red cells can be observed. If highly toxic, no emigration of leucocytes takes place, but if the liquor puris is healthy (that is, from a well drained, healthy wound), numbers of cells will emigrate on to the slide, some will show the iodophil reaction, and some, generally in the peripheral zone, will be covered with clumped red cells, while a certain number will have ingested some of the red cells which adhere to them (Fig. IX).

Thus, by the addition of a haemagglutinin to a liquor puris or a serum, we can induce the leucocytes to phagocytose native red cells—that is, red cells from the same blood—when incubated together. Iso-phagocytosis of red

cells by pus cells may also be observed in pus from wounds in patients whose blood serum is haemagglutinative.

Thus we arrive at this position. Pus cells from "whole" pus (if it contains haemagglutinins of the right kind) will ingest washed human red cells more freely than washed

FIG. IX.—Showing the phagocytosis of (C. J. B.) red cells by C. J. B. leucocytes in a drop of C. J. B. blood incubated in agglutinating liquor puris; "drop and pool" method. (Drawing from microscope slide.) 1, Agglutinated red cells; 2, red cells in phagocyte.

pus cells from the same pus will digest them, although the washing in saline actually increases the activity of the cells by removing the bacterial toxins and tryptic content of the liquor puris. The difference in the two cases depends on the fact that haemagglutinins are present in the "whole" pus and absent in the washed cells. Moreover, no ingestion of red cells occurs if the "whole" pus is non-agglutinative.

Putting all these facts together, we must, I think, conclude that, as far as the phagocytosis of red cells by leucocytes is concerned, haemagglutinins also act as opsonins and prepare the red cells for ingestion.⁵

The question remains as to the origin of the haemagglutinins in the blood serum. The fact that washed agglutinative pus cells may clump and ingest sheep's red cells in the absence of liquor puris suggests that in some cases the phagocytes can elaborate their own agglutinins and opsonins, but further work on these lines is required in this difficult field.

The Relation (if any) between the Presence of Haemagglutinins in the Blood Serum and Pus and Capacity to Resist Infection.

It is a matter of great interest to ascertain whether patients whose blood serum contains any of the various haemagglutinins for human or animal red cells show a greater or a less power of resisting infection than patients whose blood serum and pus are not agglutinative.

The subject is a difficult one. I have examined the pus, after confirmatory examination of the blood serum, from a considerable number of patients from this point of view, together with, in some cases, a bacteriological examination of the wound discharges. In the absence of recorded facts about the blood serum before the injury, I cannot satisfy myself that any constant association exists between haemagglutinin content of serum or pus and resistance to infection in any given case. At the same time there can be little doubt that a detailed examination of the blood serum and pus from a large series of cases, checked by bacteriological data concerning the nature of the infection and clinical findings bearing on recovery, would not only solve this question, but would throw valuable light on fundamental problems connected with the production of the immune state.

There is, however, some evidence that the blood serum of a patient previously known to be non-agglutinative to human (C. J. B.) red cells may, under the influence of a heavy systemic infection, become agglutinative to the same red cells. This fact, if confirmed in other cases,⁶ is important, because it is possible that the increase in

⁵ This has now been confirmed in quite a number of patients.

haemagglutinin content may serve as an indication that other collateral changes—that is, increase in antibodies and opsonins—are also occurring in the same blood serum.

Further, pus cells from a wound in a patient whose blood serum contains haemagglutinins frequently contain ingested red corpuscles. The pus cells in the presence of the iso-haemagglutinins (or the opsonins if these are separate bodies) tend to become auto-haemagglutinative and ingest their own red cells. Pus cells from wounds in patients whose blood serum contains no haemagglutinin do not phagocytose their own red cells.

It is probable that the red cell destruction and anaemia which accompany systemic infection in certain cases are associated with a blood serum and liquor puris of a haemagglutinative character.

In connexion with this inquiry, I wish to acknowledge kind help from my colleagues in military hospitals and from Dr. Mackarell, pathologist to the Leicester Royal Infirmary. The figures in the special plate are reproduced from photographs.

REFERENCES.

¹ BRITISH MEDICAL JOURNAL, June 3rd, 1916, December 23rd, 1916, and February 3rd, 1917. ² See also Dean, *Proc. Roy. Soc.*, 1905, B, 76, p. 506; and *Proc. Roy. Soc.*, 1907, B, 79, p. 399. ³ See Keith, *Proc. Roy. Soc.*, 1906, 77, p. 537. ⁴ BRITISH MEDICAL JOURNAL, February 3rd, 1917. ⁵ See Keith, *Proc. Roy. Soc.*, 1906, 77, p. 546; and Barratt, *Proc. Roy. Soc.*, 1905, 76, p. 624.

ON WAR NEPHRITIS AND SOME CIRCUMSTANCES INCIDENTAL THERETO.

BY

SIR THOMAS OLIVER, M.D., F.R.C.P.,

NORTHUMBERLAND WAR HOSPITAL, GOSFORTH, NEWCASTLE-UPON-TYNE.

It was not until fully eight months after the commencement of the war that nephritis loomed large in the sickness returns of soldiers. In the case of the German army the outbreak occurred a few months later. The fact that the winter campaign had been got over and the spring and early summer months had come was advanced by some physicians as a reason why exposure to cold could not have been the cause of the disease. The malady has fallen upon our troops in France and Flanders with unequal incidence and with unequal severity. Occasionally the number of men from one convoy admitted for nephritis under the care of my colleague Dr. Dagger and myself has equalled 40 per cent. of the total, while on other occasions there would be only 10 per cent. or less. It is sometimes said that the number of cases of nephritis among our soldiers at the front has not been larger than that occurring among the civil population. I can hardly subscribe to this opinion, and I base my dissent upon experience of admissions into the Royal Victoria Infirmary, Newcastle, extending over three decades. In the summer of 1915 few of the cases of nephritis admitted into the Northumberland War Hospital presented acute symptoms. This might be explained partly by the fact that the severer and more acute cases had been retained in hospitals on the Continent. Beyond albuminuria and complaints of headache and backache, there were few physical signs and symptoms. Oedema, with us, has been an infrequent physical sign compared with the experience of Colonel J. Michell Clarke,¹ who found it in fifty-seven out of seventy-four patients. It was during my first visit to France in September, 1915, that I saw in the stationary hospitals there cases of nephritis with acute symptoms, including dropsy and uraemia.

War nephritis, to give it the name by which it is generally known, differs from ordinary nephritis in that there is a greater tendency for blood, and blood in fairly large quantity, to appear in the urine, for the specific gravity in many cases to be well maintained, for the urine to be not so scanty, the blood pressure to be lower, also for the malady to be attended by fewer cardio-vascular changes, and to be followed by fewer complications than are found in ordinary nephritis.

We have had no deaths from war nephritis in the Northumberland War Hospital. Hurst² states that death in this malady is extremely rare, not more than 3 or 4 per 1,000 patients, and that when death takes place it is usually through uraemia. A low death-rate has also been noted in the Italian army. On the other hand, and as illustrating the absence of uniform experience, three of Colonel Michell

Clarke's 68 patients died. A fair percentage of our patients at the Northumberland War Hospital have recovered, but in men over 30 years of age albuminuria has frequently persisted, the urine has become pale and watery, and its specific gravity has decreased. Many of these men have become the subjects of chronic kidney disease. In a considerable number of the younger men recovery has only been apparent, for albuminuria after having been absent for days or weeks has returned, without any exaggeration of the symptoms, other than fresh complaint of backache and the presence of blood in the urine. The length of time required for the disappearance of the albuminuria varies—it may be three weeks or it may be as many months, but the longer it persists the more likely are the kidneys to become the seat of chronic and permanent structural changes. While error in diet is a cause of recrudescence of symptoms, muscular exercise—for example, tennis playing and too long walks—may also conduce to their return.

I have alluded to the fact that in war nephritis there are fewer symptoms and cardio-vascular signs than in the usual forms of nephritis with which we are familiar; hence the comparative absence of well-marked hypertrophy of the left ventricle of the heart and of loud accentuation of the second sound over the aorta. The short duration of the illness may partly, but it cannot altogether, explain the comparative infrequency of cardiac hypertrophy, and of thickened arteries.

I have not been able to ascertain how far the previous occupation of the soldier plays any part in the development and course of the malady. On the Italian front it has been observed that men whose previous occupations exposed them to hardships and who were intemperate have suffered from nephritis in larger numbers than men engaged in more protected occupations and who are of temperate habits.

Some of the soldiers were slightly anaemic, but they seldom exhibited the well known features of patients suffering from Bright's disease. I have not found any well marked reduction in the number of red blood corpuscles, nor any increase or decrease of leucocytes, nor has basophilia been a feature of the blood examinations. Of the presence of eosinophile corpuscles in the urine and their excess in the blood as described by Bernhardt I have no experience. A positive Wassermann reaction was not obtained in a series of 13 of my cases. McKenzie Wallis obtained a positive reaction in 18 out of 56 patients. Since other physicians have secured the reaction, and seen relief follow antisyphilitic treatment, it is possible that syphilis is more than an incident in some forms of war nephritis. The blood pressure has varied from 70 to 130 mm. Hg; in only a small percentage did it rise to or over 180. Some of the soldiers were admitted into the hospital not for kidney trouble at all, but for trench fever, P.U.O., or on account of having been gassed.

SYMPTOMS.

The disease usually commences with a feeling of malaise, pains, especially in the limbs and back, headache, and occasionally sickness with oedema of face and feet. Among German soldiers the disease is said to be ushered in with cough, worst at night, and with shortness of breath, attended by oedema of face, and by thirst and scanty secretion of urine. Our own men, too, suffer at the onset from bronchitis and dyspnoea. By the time the patients have reached the North of England dropsy, if ever present, has usually disappeared. Anasarca we never see. In none of our patients has vision been affected, nor have there been uraemic convulsions. One patient developed acute pulmonary oedema, and, after a serious illness, made a good recovery from the lung trouble, but albuminuria still persisted. It would be interesting to know what the experience of the R.A.M.D. is in regard to the future of the men who have been treated for nephritis, who recovered and were, in the early months of the war, again sent back to the front. Since in a fairly large percentage of patients who have been suffering from nephritis relapses occur, no case should be regarded as cured until, after several months have passed, it is found that, on full diet and after muscular exercise, albumin does not reappear in the urine.

ETIOLOGY.

The causation of war nephritis is obscure. Although the urine and the blood of my patients have on several occasions been examined bacteriologically, in none of them