peripheral perfusion. It is important to appreciate that the blood pressure may be normal or raised in the presence of severe hypovolaemia in these children—because of increased plasma renin concentrations produced by renal ischaemia and increased plasma vasopressin concentrations.⁶

There is debate over the pathogenesis of nephrotic oedema.⁷ It is thought that the “underfill” mechanism is more common in childhood nephrotic syndrome. Hypoalbuminaemia leads to reduced plasma oncotic pressure, hypovolaemia, and secondary renal salt and water retention. Infusions of 20% albumin are therefore appropriate in hypovolaemic patients.

We recommend 20% albumin at a dose of 1 g/kg ideal body weight over four hours. We consider that one to two hours, recommended by the British Association for Paediatric Nephrology,¹ is too rapid and increases the risk of fluid overload. During the infusion it is essential to monitor the pulse, blood pressure, and respiratory rate and to look for developing jugular venous engorgement and dyspnoea. These observations should continue for two hours after the infusion, as fluid shift may continue. Blood pressure may actually fall to a more normal level.⁷ Frusemide should not be given while clinical assessment continues to suggest hypovolaemia.

Occasionally nephrotic syndrome may lead to profound hypovolaemia and shock. Rapid expansion of the intravascular volume with 4-5% human albumin solution 10-20 ml/kg over 30-60 minutes is appropriate. Subsequent infusions of 20% albumin may then also be needed.

If the child is assessed as normovolaemic and has severe symptomatic oedema we recommend 20% albumin at a dose of 1 g/kg ideal body weight over four hours, with frusemide 1 mg/kg given two hours into the infusions.

If signs of fluid overload do develop, frusemide 1 mg/kg should be given intravenously. If this produces no improvement 5 mg/kg should be given. Further deterioration may necessitate venesection and ventilation and referral to a centre with facilities for paediatric intensive care and haemofiltration.

As intravascular volume may change daily until remission occurs it must be assessed before each infusion of 20% albumin. Advice over the telephone from a paediatric nephrologist may help in difficult cases.

1 Report of a Workshop by the British Association for Paediatric Nephrology and Research Unit, Royal College of Physicians. Consensus statement on management and audit potential for steroid responsive nephrotic syndrome. *Arch Dis Child* 1994;70:151-7.

2 Luetscher JA, Hall AD, Kremer VL. Treatment of nephrosis with concentrated human serum albumin. I. Effects on the proteins of body fluids. *J Clin Invest* 1949;28:700-12.

3 Weiss RA, Schoeneman M, Greifer I. Treatment of severe nephrotic edema with albumin and furosemide. *N Y State J Med* 1984;84:384-6.

4 Report of the International Study of Kidney Disease in Children. Minimal change nephrotic syndrome in children: deaths during the first 5 to 15 years' observation. *Pediatrics* 1984;73:497-501.

5 Haws RM, Baum M. Efficacy of albumin and diuretic therapy in children with nephrotic syndrome. *Pediatrics* 1993;91:1142-6.

6 Houtman PN, Shah V, Barratt TM, Dillon MJ. Reduction of hypertension in hypovolaemia (letter). *Lancet* 1990;336:1454.

7 Humphreys MH. Mechanisms and management of nephrotic oedema. *Kidney Int* 1994;45:266-81.

(Accepted 5 June 1995)

Controversies in Management

Should backache be treated with spinal fusion?

Spinal fusion is the only treatment for discogenic pain

John P O'Brien

Backache indicates pain originating in the vertebral column (most commonly in the lowest two mobile segments of the lumbar spine). It is felt mainly in the lower back and buttocks, often being referred to the lower limbs. Referred pain in the limb can be confused with radicular pain (sciatica) from nerve root compression.

Fusion is a non-specific term. It comes from the Latin *fundere*—to pour or to melt. In the context of the spine, fusion entails welding or stiffening two or more vertebral bodies. Arthrodesis might therefore be a more appropriate word (removing of the articular surfaces and securing bony union¹).

History

Hippocrates in 380 BC observed spontaneous fusion of the facet joints in a case of spinal tuberculosis. It was nature's attempt to halt the progress of the deformity. Hibbs read the work of Hippocrates, and in 1911 surgically fused the posterior spinal elements in young patients with spinal tuberculosis.² This successfully prevented subsequent deformity. Later Hodgson and Stock used anterior spinal fusion for tuberculosis and then the correction and fusion of scoliosis³; this became standard surgical treatment for these problems.

Fusion today

Today fusion is an important part of the conventional surgical management of spinal tumour, defor-

mity, and infection as well as trauma resulting in an unstable compression fracture visible in an x ray film. The role of fusion is questioned, however, when trauma produces a symptomatic tear of the annulus not visible in an x ray film.

There are few published descriptions of the rationale for fusion of the lumbar spine, but consider the following two:

Firstly, movement of a motion segment is painful. Therefore elimination of that motion between vertebrae should eliminate the pain.⁴ Depending on the preference and training of the surgeon, the fusion may be performed through the facet joints or between the vertebral bodies (the latter is mechanically preferable), between the laminae (obsolete), or between the transverse processes (traditional).

Secondly, pain is shown by discography to be due to a torn annulus—so-called discogenic pain. Removal of the annulus is a logical clinical decision, optimally through an anterior approach. But the spine is clearly destabilised by this disc clearance. A bone graft to replace the disc by filling the defect and maintaining the height of the disc space is therefore appropriate and logical.

Why use fusion?

The controversy surrounds primary fusion for backache. The indication for fusion is generally agreed for spondylolisthesis or postsurgical instability after extensive laminectomy.

What is the problem, then, with fusion for back pain? One problem is that fusion is used to treat a comparatively unusual lesion which was not well described until recently.⁵ The pain has often been regarded as inexplicable (if not psychiatric), and consequently neurosurgeons have concentrated on the nerve roots and problems arising from canal stenosis while orthopaedic surgeons have been distracted by the success rate and ease of joint replacement surgery.

It is now clear, however, that a severe injury or repeated small injuries to a motion segment will commonly produce a tear in the annulus of the disc.⁵ The outer annulus has a rich sensory innervation but the disc is relatively avascular and so has poor healing potential.⁶ Partial healing with breakdown of inferior scar tissue is therefore common, and patients often experience recurrent attacks of back pain.

The correct surgical principle is to identify the anatomical source of the pain by discography (magnetic resonance imaging can miss annular tears). The source of pain, commonly an annular tear, should then be removed surgically through an anterior approach and the disc replaced with bone plugs to stabilise the spine.

Success of surgery

Wiltse noted many patients in whom fusion was unquestionably solid but severe pain persisted.⁶ It is not uncommon for a patient with a technically successful posterior fusion to have persisting disabling discogenic pain.⁸ These symptoms are heightened by neovascularisation and inflammatory cell infiltration of the anterior annulus (A J Freemont, personal communication).

Palpation of the anterior annulus of these discs through the abdomen will produce severe pain radiating through to the back, which is often the only positive clinical sign.⁹ Recent observations by Kuslich *et al* confirm that the outer annulus is likely to be responsible for symptoms in most cases of low back pain in which the nerve root is not trapped.¹⁰ The main fault of many types of fusion is that they do not specifically address the pain source.⁶

Fusion for backache is not an alternative to expert rehabilitation with muscle retraining and emphasis on the abdominal muscle mechanism. It is only when these measures have failed that surgery should be considered. If inadequate surgical training, poor surgical technique, poor patient selection, wrong level surgery, and incorrect diagnosis prevail fusion for backache must stop. But with the right preparation, for the right patient, it is the right treatment because it is the only treatment of the painful lesion.

- 1 Blakiston's *Gould Medical Dictionary*. New York: McGraw Hill, 1972:137.
- 2 Hibbs RH. An operation for progressive spinal deformities. *N Y J Med* 1911;93:1013.
- 3 Hodgson AR, Stock FE. Anterior spinal fusion—a preliminary communication on the radical treatment of Pott's disease and Pott's paraplegia. *Br J Surg* 1956;44:266-75.
- 4 Farfan HF, Kirkaldy-Willis WH. The present status of spinal fusion and the treatment of lumbar intervertebral joint disorders. *Clin Orthop* 1981: 198-214.
- 5 Park WM, McCall I, O'Brien JP, Webb JK. Fissuring of the posterior annulus fibrosus in the lumbar spine. *Br J Radiol* 1979;52:382-7.
- 6 Yoshizawa H, O'Brien JP, Smith WT, Trumper M. The neuropathology of intervertebral discs removed for low back pain. *J Pathol* 1980;132: 95-104.
- 7 Wiltse LL. Surgery for intervertebral disc disease of the lumbar spine. *Clin Orthop* 1977;129:22-45.
- 8 Weatherley CR, Prickett CF, O'Brien JP. Discogenic pain persisting despite solid posterior fusion. *J Bone Joint Surg (Br)* 1986;68:142-3.
- 9 O'Brien JP. Anterior spinal tenderness in low-back pain syndromes. *Spine* 1979;4:85-8.
- 10 Kuslich SD, Ulstrom CL, Michael CJ. The tissue origin of low back pain and sciatica. *Orthop Clin North Am* 1991;22:181-7.

The case for spinal fusion is unproved

James Wilson-MacDonald

Back pain affects about 80% of the population at some time in their life, and it is generally agreed that most people are best treated conservatively. Most episodes of acute back pain will settle either spontaneously or with treatment, but in a few people the pain does not resolve and they may be considered for spinal fusion.

Spinal fusion was originally developed for the treatment of tuberculosis, poliomyelitis, and scoliosis. Because it seemed successful for treating back pain arising from these conditions, it was assumed that any mechanical back pain could be successfully treated with spinal fusion. More than 20 000 lumbar spinal fusions are performed annually in the United States, and there is a direct relation between the number of spinal operations performed in any one area and the number of orthopaedic and neurosurgeons.¹ About eight times as many spinal surgical procedures are carried out in the United States per capita as in Britain.

Conservative treatment

Prospective studies have shown reasonably convincingly that time off work in patients with first episodes of acute back pain can be reduced by 30-50% by early physiotherapy.² Chiropractic treatment has also been shown to have a similar success rate.

Treatment for chronic back pain is not so convincing. Koes found that only four of the randomised controlled trials of exercise therapy for back pain were of an adequate standard to draw any definite

conclusions, and it was impossible to decide whether exercise therapy was better than any other type of treatment, or whether one type of exercise was better than another.³ However, some studies have suggested that conservative treatment can be at least as successful as any surgical treatment. Kohles *et al* showed that up to 85% of patients with chronic back pain can return to work with intensive retraining methods.⁴ They used a method called "functional restoration" in which a team of professionals assesses and treats patients with chronic low back pain. Trunk strength and spinal range of motion were used as objective measurements of progress and success.

Back pain is the commonest cause of loss of work among people under 45, with huge economic cost, but there is little incentive in the health service to apply resources to intensive retraining programmes for these patients.

Spinal fusion

Spinal fusion for back pain has never been compared with non-surgical treatment in a randomised controlled trial.⁵ The only non-randomised prospective trials of fusion have compared surgery with and without fusion for herniated discs. Of all the papers on spinal fusion for back pain, only 47 had adequate follow up, and although 68% of patients reported in these papers had a satisfactory outcome from fusion, the success rate varied between 16% and 95%.⁵ There

Spinal Unit, Nuffield
Orthopaedic Centre,
Oxford OX3 7LD
James Wilson-MacDonald,
consultant orthopaedic surgeon