What is the problem, then, with fusion for back pain? One problem is that fusion is used to treat a comparatively unusual lesion which was not well described until recently.⁵ The pain has often been regarded as inexplicable (if not psychiatric), and consequently neurosurgeons have concentrated on the nerve roots and problems arising from canal stenosis while orthopaedic surgeons have been distracted by the success rate and ease of joint replacement surgery.

It is now clear, however, that a severe injury or repeated small injuries to a motion segment will commonly produce a tear in the annulus of the disc.⁵ The outer annulus has a rich sensory innervation but the disc is relatively avascular and so has poor healing potential.⁶ Partial healing with breakdown of inferior scar tissue is therefore common, and patients often experience recurrent attacks of back pain.

The correct surgical principle is to identify the anatomical source of the pain by discography (magnetic resonance imaging can miss annular tears). The source of pain, commonly an annular tear, should then be removed surgically through an anterior approach and the disc replaced with bone plugs to stabilise the spine.

Success of surgery

Wiltse noted many patients in whom fusion was unquestionably solid but severe pain persisted.⁶ It is not uncommon for a patient with a technically successful posterior fusion to have persisting disabling discogenic pain.⁸ These symptoms are heightened by neovascularisation and inflammatory cell infiltration of the anterior annulus (A J Freemont, personal communication).

Palpation of the anterior annulus of these discs through the abdomen will produce severe pain radiating through to the back, which is often the only positive clinical sign.⁹ Recent observations by Kuslich *et al* confirm that the outer annulus is likely to be responsible for symptoms in most cases of low back pain in which the nerve root is not trapped.¹⁰ The main fault of many types of fusion is that they do not specifically address the pain source.⁶

Fusion for backache is not an alternative to expert rehabilitation with muscle retraining and emphasis on the abdominal muscle mechanism. It is only when these measures have failed that surgery should be considered. If inadequate surgical training, poor surgical technique, poor patient selection, wrong level surgery, and incorrect diagnosis prevail fusion for backache must stop. But with the right preparation, for the right patient, it is the right treatment because it is the only treatment of the painful lesion.

- 1 Blakiston's *Gould Medical Dictionary*. New York: McGraw Hill, 1972:137.
- 2 Hibbs RH. An operation for progressive spinal deformities. *N Y J Med* 1911;93:1013.
- 3 Hodgson AR, Stock FE. Anterior spinal fusion—a preliminary communication on the radical treatment of Pott's disease and Pott's paraplegia. *Br J Surg* 1956;44:266-75.
- 4 Farfan HF, Kirkaldy-Willis WH. The present status of spinal fusion and the treatment of lumbar intervertebral joint disorders. *Clin Orthop* 1981; 198:214.
- 5 Park WM, McCall I, O'Brien JP, Webb JK. Fissuring of the posterior annulus fibrosus in the lumbar spine. *Br J Radiol* 1979;52:382-7.
- 6 Yoshizawa H, O'Brien JP, Smith WT, Trumper M. The neuropathology of intervertebral discs removed for low back pain. *J Pathol* 1980;132: 95-104.
- 7 Wiltse LL. Surgery for intervertebral disc disease of the lumbar spine. *Clin Orthop* 1977;129:22-45.
- 8 Weatherley CR, Prickett CF, O'Brien JP. Discogenic pain persisting despite solid posterior fusion. *J Bone Joint Surg (Br)* 1986;68:142-3.
- 9 O'Brien JP. Anterior spinal tenderness in low-back pain syndromes. *Spine* 1979;4:85-8.
- 10 Kuslich SD, Ulstrom CL, Michael CJ. The tissue origin of low back pain and sciatica. *Orthop Clin North Am* 1991;22:181-7.

The case for spinal fusion is unproved

James Wilson-MacDonald

Back pain affects about 80% of the population at some time in their life, and it is generally agreed that most people are best treated conservatively. Most episodes of acute back pain will settle either spontaneously or with treatment, but in a few people the pain does not resolve and they may be considered for spinal fusion.

Spinal fusion was originally developed for the treatment of tuberculosis, poliomyelitis, and scoliosis. Because it seemed successful for treating back pain arising from these conditions, it was assumed that any mechanical back pain could be successfully treated with spinal fusion. More than 20 000 lumbar spinal fusions are performed annually in the United States, and there is a direct relation between the number of spinal operations performed in any one area and the number of orthopaedic and neurosurgeons.¹ About eight times as many spinal surgical procedures are carried out in the United States per capita as in Britain.

Conservative treatment

Prospective studies have shown reasonably convincingly that time off work in patients with first episodes of acute back pain can be reduced by 30-50% by early physiotherapy.² Chiropractic treatment has also been shown to have a similar success rate.

Treatment for chronic back pain is not so convincing. Koes found that only four of the randomised controlled trials of exercise therapy for back pain were of an adequate standard to draw any definite

conclusions, and it was impossible to decide whether exercise therapy was better than any other type of treatment, or whether one type of exercise was better than another.³ However, some studies have suggested that conservative treatment can be at least as successful as any surgical treatment. Kohles *et al* showed that up to 85% of patients with chronic back pain can return to work with intensive retraining methods.⁴ They used a method called "functional restoration" in which a team of professionals assesses and treats patients with chronic low back pain. Trunk strength and spinal range of motion were used as objective measurements of progress and success.

Back pain is the commonest cause of loss of work among people under 45, with huge economic cost, but there is little incentive in the health service to apply resources to intensive retraining programmes for these patients.

Spinal fusion

Spinal fusion for back pain has never been compared with non-surgical treatment in a randomised controlled trial.⁵ The only non-randomised prospective trials of fusion have compared surgery with and without fusion for herniated discs. Of all the papers on spinal fusion for back pain, only 47 had adequate follow up, and although 68% of patients reported in these papers had a satisfactory outcome from fusion, the success rate varied between 16% and 95%.⁵ There

Spinal Unit, Nuffield
Orthopaedic Centre,
Oxford OX3 7LD
James Wilson-MacDonald,
consultant orthopaedic surgeon

was so much variability in the methods of treatment, selection, and assessment of patients that it was impossible to compare the results or to draw any conclusions about the appropriateness of fusion for back pain.

Lumbar fusion is probably contraindicated in many circumstances. For example, the association of smoking with pseudarthrosis and failure of spinal fusion is well documented. A recent study of people seeking compensation for a work induced injury showed that the back pain and lifestyle of most deteriorated after spinal fusion.⁶ Fusion rates can be improved with new techniques such as segmental spinal stabilisation using pedicle screws,⁷ and there is a correlation between fusion rate and resolution of pain.⁵ These implants make the surgery more technically demanding and more satisfying to the surgeon, but new techniques often bring with them new complications. In a survey of over 600 surgical cases, 21.2% of patients had serious complications such as nerve root damage.⁸ Hospital costs alone for failed spinal surgery in the United States are estimated to be about \$2bn (£1.3bn) a year.

Some evidence suggests that many conditions which cause back pain can be successfully treated with spinal fusion. For example, in young patients with spondylolithesis spinal decompression with simultaneous spinal fusion was reported to give more satisfactory results than decompression alone.⁹ Similarly spinal fusion for back pain and spondylolithesis can give good results, especially in children.¹⁰ However, even these procedures have not been subjected to randomised trials. Surgery to replace the lumbar discs or operations which change the balance of the spine with posterior ligament devices may prove more effective than spinal fusion in the long term treatment of patients with back pain.

Conclusion

The case for spinal fusion for back pain has not been proved. Excellent results have been obtained with non-surgical treatments, and adequate resources have not been applied to fully assess these methods. Spinal fusion probably has a place in the treatment of back pain in certain circumstances, but many patients selected for spinal fusion at the present time might be better treated in other ways.

A multicentre randomised prospective clinical trial is currently being planned, and this will answer some

of the questions regarding the efficacy of spinal fusion and other surgical procedures in the treatment of patients with back pain. Total joint replacement has almost replaced fusion of the hip or the knee for arthritis, and in future spinal fusion may be replaced by other more effective treatments in much the same way.

- 1 Cherkin D, Deyo R, Loeser J, Bush T. *An international comparison of back surgery rates*. Marseilles: International Society for the Study of the Lumbar Spine, 1993.
- 2 Mitchell RJ, Carmen GM. Results of a multicenter trial using an intensive active exercise program for the treatment of acute soft tissue and back injuries. *Spine* 1990;15:514-21.
- 3 Koes BW, Bouter LM, Beckerman H, van der Heijden GJMG, Knipschild PG. Physiotherapy exercises and back pain: a blinded review. *BMJ* 1991;302:1572-6.
- 4 Kohles S, Barnes D, Gatchel J, Mayer TG. Improved physical performance outcomes after functional restoration treatment in patients with chronic low-back pain. *Spine* 1990;15:1321-4.
- 5 Turner JA, Ersek M, Herron L, Haselkorn J, Kent D, Ciol MA, et al. Patient outcomes after lumbar spinal fusions. *JAMA* 1992;19:907-12.
- 6 Franklin GM, Haug J, Heyer NJ, McKeefrey SP, Picciano JF. Outcome of lumbar fusion in Washington state workers compensation. *Spine* 1994;19:1897-904.
- 7 Zdeblick TA. A prospective randomized study of spinal fusion. *Spine* 1993;18:983-91.
- 8 Esses S, Sachs BL, Dreyzin V. Complications associated with the technique of pedicle screw fixation. A selected survey of ABS members. *Spine* 1993;18:2231-8.
- 9 Herkowitz HN, Kurz LT. Degenerative lumbar spondylolithesis with spinal stenosis: a prospective study comparing decompression and decompression and intertransverse process arthrodesis. *J Bone Joint Surg Am* 1991;73:802-8.
- 10 Johnson JR, Kirwan E O'G. The long term results of fusion in situ for severe spondylolithesis. *J Bone Joint Surg Br* 1983;65:43-6.

Commentary

P C Rubin

Backache is common but not well understood; spinal fusion is expensive, potentially hazardous, and of no proved benefit—these were the central messages that I took from these articles. On the one hand we have an enthusiastic advocate of spinal fusion putting forward reasons why spinal fusion should work—such as that movement of a motion segment is painful. On the other side is the simple fact that the technique has not been subjected to controlled clinical trial and may be associated with important complications in as many as a fifth of cases. Hard choices are having to be made about priorities for expenditure in the NHS. On the evidence available I would find it hard to support the purchase of spinal fusion for backache.

Department of Medicine,
Division of Therapeutics,
University Hospital,
Queen's Medical Centre,
Nottingham NG7 2UH
P C Rubin, professor

ANY QUESTIONS

How should a wasp sting causing swelling of the tongue be treated?

As with all cases of anaphylaxis, the treatment will be dictated by the severity of the effects, the site, and the speed of the progression of symptoms. Oxygen should be given and a nasopharyngeal airway inserted if available. Adrenaline should be given: this usually prevents further swelling.¹ The intramuscular route is preferable in an unmonitored patient without dyspnoea, but in a monitored patient or one who has collapsed the intravenous route is preferable.²

There are two other potential routes of administration. Inhaled adrenaline has been advocated,³ but it has the disadvantage of requiring equipment, and in patients with airway oedema the distribution or the ability to use the inhaler may be impaired. It is less effective if other symptoms of anaphylaxis occur.⁴ The adrenaline could also be injected directly into the tongue, but the possible

consequences of causing bleeding in the swollen tongue probably preclude this.

The patient should be taken to hospital for observation. An H₁ antihistamine should be given when the patient is stable as late recurrence of angio-oedema has been documented. If the patient has collapsed intubation or emergency cricothyroidectomy should be performed. —MALCOLM FISHER, head, intensive therapy unit, Royal North Shore Hospital, St Leonards, New South Wales, Australia

- 1 Fisher MM. Clinical observations on the pathophysiology and treatment of anaphylactic cardiovascular collapse. *Anaesth Intensive Care* 1986;14:17-21.
- 2 Fisher MM. Treating anaphylaxis with sympathomimetic drugs. *BMJ* 1992;305:1107-8.
- 3 Muller U, Mosbech H, Blaauw P, Dreborg S, Malling HJ, Przybilla B, Urbanek R, et al. Emergency treatment of allergic reactions to hymenoptera stings. *Clin Exp Allergy* 1991;21:281-8.
- 4 Ewan PW. Route of administration of adrenaline for the treatment of anaphylactic reactions to bee or wasp stings. *Clin Exp Allergy* 1991;21:753-5.