overtaken and training requirements will become important.

If pretest counselling were more complete and informed consent required then the public acceptance and uptake of genetic tests might not be as great as figures from the current neonatal screening programmes suggest.

ADRIAN EDWARDS
Clinical fellow

Department of General Practice,
University of Wales College of Medicine,
Llanedeyrn Health Centre,
Llanedeyrn,
Cardiff CF3 7PM

- 1 Wilson JMG, Junger YG. *Principles and practice of screening for disease*. Geneva: WHO, 1968. (Public health papers No 34.)
- 2 Streetly A, Grant C, Pollitt RJ, Addison GM. Survey of scope of neonatal screening in the United Kingdom. *BMJ* 1995;311:726. (16 September.)

Pitfalls in the interpretation of tumour markers

EDITOR,—We do not agree with H S Pandha and colleagues that poor sensitivity and specificity of assays cause a major problem in the measurement of α fetoprotein concentrations.¹ Measurements of α fetoprotein are used in several clinical situations, including to diagnose and monitor neoplastic disorders (such as germ cell tumours and hepatomas) and in antenatal screening for Down's syndrome and neural tube defects. Modern immunoassays for α fetoprotein have adequate sensitivity, and it is important for clinical laboratories to use an assay that is specific for α fetoprotein but that adequately detects all its molecular forms in view of the variety of clinical settings in which α fetoprotein is measured. In specific cases, identification of the particular form of α fetoprotein secreted may be valuable, but this is not generally required.

Although concentrations of both α fetoprotein and human chorionic gonadotrophin were initially raised in the case reported, the authors do not comment on the absence of any increase in the concentration of human chorionic gonadotrophin concomitant with the second increase in α fetoprotein concentration. This might have suggested that this increase in α fetoprotein concentration was not due to recurrence of the primary tumour. However, recurrent tumour deposits or metastases may not secrete the same tumour markers (or may not secrete them in the same proportions) as the primary tumour, and this may cause particular difficulty with heterogeneous tumours such as teratomas.

D A OLEESKY
Senior registrar in medical biochemistry
R FIFIELD
Consultant (grade C) biochemist

Suprarenal Protein Reference Unit,
Medical Biochemistry Department,
Cardiff Royal Infirmary,
Cardiff CF2 1SZ

- 1 Pandha HS, Wasan HS, Harrington K, Waxman J. Failure of normalisation of α fetoprotein concentration after successful treatment of teratoma. *BMJ* 1995;311:434-5. (12 August.)

Risk factors for acanthamoeba keratitis

EDITOR,—We wish to respond to the letters¹ about our case-control study of risk factors for acanthamoeba keratitis in people who use contact lenses.² We undertook the study to determine the reasons for the increase in the number of cases of acanthamoeba keratitis seen at Moorfields Eye Hospital. Aware of the multiplicity of risk factors for the disease, we performed multivariable analysis and commented on factors identified as significant. David V Seal and John Hay do not

identify the "sweeping generalisations" or "misleading comments" they say that we made, and we are not aware of having made any.

We did not claim that frequent replacement of the lens case alone would prevent growth and replication of acanthamoeba when chlorine based disinfectants are used, but we suggested that monthly replacement of the case, which we have promoted since the data were collected, "may reduce the prevalence of contamination of storage cases." Since cases containing acanthamoeba usually have bacterial or fungal co-contamination, or both,³ we reasoned that a general reduction in the rate of microbial contamination, as occurs with regular replacement of cases,⁴ may reduce the likelihood of acanthamoeba contamination. Obviously, the use of disinfectant with a cysticidal effect is preferable.

With regard to the crude minimum relative risk of 2.63 ($P=0.004$) calculated for Acuvue compared with other brands of daily wear disposable lenses, we accept that the introduction of five (not 47) brands of disposable lenses (according to our definition²) by the time that the controls were collected may have led to a slight overestimate. Given the high proportion (84%) of Vistakon lenses used by the controls using disposable lenses, however, this is unlikely to have been appreciable. To confirm this we have recalculated the crude relative risk for patients presenting in the final year of the study: among users of daily wear disposable lenses ($n=46$) Acuvue lenses were still associated with a significant excess risk (crude minimum relative risk 1.16 ($P=0.044$)).

Our epidemiological data suggest that there may be a causal relation between certain disposable lenses and acanthamoeba keratitis. Although there is no laboratory evidence of this, available data are limited. We agree that further investigation is required, both in the laboratory and in a nationwide population based incidence study.

CHERRY F RADFORD
Research optometrist
JOHN K G DART
Consultant ophthalmologist

Moorfields Eye Hospital,
London EC1V 2PD

DARWIN C MINASSIAN
Senior lecturer in epidemiology

Department of Preventive Ophthalmology,
Institute of Ophthalmology,
London EC1V 9EL

- 1 Risk factors for acanthamoeba keratitis [letters]. *BMJ* 1995;311:808. (23 September.)
- 2 Radford CF, Bacon AS, Dart JKG, Minassian DC. Risk factors for acanthamoeba keratitis in contact lens wearers: a case-control study. *BMJ* 1995;310:1567-70. (17 June.)
- 3 Bottone EJ, Madayag RM, Qureshi MN. Acanthamoeba keratitis: synergy between amoebic and bacterial cocontaminants in contact lens care systems as a prelude to infection. *J Clin Microbiol* 1992;30:2447-50.
- 4 Devonshire P, Munro FA, Abernathy C, Clark BJ. Microbial contamination of contact lens cases in the west of Scotland. *Br J Ophthalmol* 1993;77:41-5.

Placebo controlled trials of ondansetron for postoperative nausea and vomiting

EDITOR,—Rebecca L Aspinall and Neville W Goodman conclude that patients who participated in placebo controlled studies investigating the role of ondansetron in the management of postoperative nausea and vomiting were denied access to effective treatment.¹ This is not correct: no patient was denied access to effective treatment, and the health and wellbeing of patients are of overriding importance in the design of all studies sponsored by GlaxoWellcome. Any patient who developed nausea and vomiting could request and receive rescue treatment at any time. In addition, all of the studies discussed were subject to local ethical approval by independent review boards or

ethics committees, and all the patients were fully informed about the implications of the studies and volunteered to participate.

Aspinall and Goodman concede that the use of antiemetics for prophylaxis against postoperative nausea and vomiting is not accepted by all anaesthetists. There are many causes of postoperative nausea and vomiting, and placebo controlled studies can provide valuable information on the relative influence that the various factors may have on its occurrence.² In the studies to investigate the efficacy of ondansetron in established postoperative nausea and vomiting the number of patients who received placebo was kept to a minimum. Indeed, the true number of patients included in the studies cited by Aspinall and Goodman was 313 and not "over 400" as the quoted figures contain some duplicate reports.

There will always be debate about the need for and desirability of placebo controlled studies irrespective of the disease or treatment under investigation. GlaxoWellcome, however, will continue to work to high clinical standards to ensure the development of safe and effective medicines that meet the needs of patients, clinicians, and regulators.

JAMES B D PALMER
Director

Group Medical Operations,
GlaxoWellcome,
Greenford,
Middlesex UB6 0HE

CHRIS G HAIGH
Head, anaesthesiology and antiemesis

TREVOR G GIBBS
Director

CHARLOTTE M TAYLOR
Medical strategy manager

International Medical Affairs,
GlaxoWellcome,
Uxbridge,
Middlesex UB11 1BU

- 1 Aspinall RL, Goodman NW. Denial of effective treatment and poor quality of clinical information in placebo controlled trials of ondansetron for postoperative nausea and vomiting: a review of published studies. *BMJ* 1995;311:844-6. (30 September.)
- 2 Palazzo MGA, Strunin L. Anaesthesia and emesis. I. Etiology. *Canadian Anaesthetists Society Journal* 1984;31:178-87.

Deprivation payments should be based on enumeration districts

EDITOR,—In 1981 the Saffron estate, a district of council housing afflicted by poverty, made up one half of an electoral ward. The other half of the ward was a comfortable suburb outside the Saffron Group Practice's area. The Jarman index for the ward was 22. Although nearly 5000 of our patients lived in enumeration districts with Jarman indices over 30 or 40, the practice received no deprivation payments until July 1995. In 1983 the ward boundaries were redrawn. The estate now forms part of a new ward, and the Jarman score is no longer heavily diluted by the prosperous suburb.

It is ironic that, although deprivation on the Saffron estate, as measured by the Jarman index, has improved, the family health services authority now makes deprivation payments to the practice worth £29 000 a year. Decisions made by the Electoral Boundary Commission result in the allocation or denial of huge resources to primary care. A further revision of boundaries or small demographic changes could suddenly deprive our practice of over 15% of its capitation income. We cannot make long term financial plans in the face of such uncertainty.

I welcome the conclusion by Tim Crayford and colleagues that deprivation payments should be based on enumeration districts in future.¹ The practice would then be able to decide how to use these extra resources. As well as asking the Department of Health to make this change now I would like the philosophy behind these payments to be clarified. I believe that they should be regarded as a supplementary capitation fee to allow practices in deprived areas with high workloads to maintain