smaller lists but enjoy similar incomes to those of practices in more prosperous areas. There is a danger that health planners may see the payments as the resources needed to provide additional staff and better premises. The provision of extra staff to meet greater health needs and adequate cost rent funding of high land and building costs are separate issues. Unless the Department of Health affirms its belief in the value of small lists in deprived areas the serious problem of recruitment of doctors to work in these areas will rapidly become a crisis.

ADRIAN HASTINGS
General practitioner

Saffron Group Practice,
Leicester LE2 6UL

1 Crayford T, Shanks J, Bajekal M, Langford S. Analysis from inner London of deprivation payments based on enumeration districts rather than wards. *BMJ* 1995;311:787-8. (23 September.)

Public discussion of rationing decisions is essential

EDITOR,—David J Hunter offers an eloquent defence of "muddling through elegantly" in the cause of health care rationing or priority setting.¹ Yet in counselling against a national debate on priority setting he confuses different aspects of the priority setting process. Decisions by clinicians on whether and how to treat individual patients do not need to be open to public scrutiny and are best made implicitly. The same does not apply to decision making at the level of health authorities or government. It is here that both transparency and public debate are essential. As Rudolf Klein comments in his editorial in the same issue, other countries have shown how this can be done.²

To argue against public discussion is to run the risk of paternalism of the worst kind. Decisions on priorities for communities and whole populations involve value judgments that are too important to be left to professionals. It is all the more important, therefore, that these decisions involve as wide a set of interests as possible. The outcome may not always satisfy the experts, but this is always a risk in a democratic process.

CHRIS HAM
Director

University of Birmingham,
Health Services Management Centre,
Birmingham B15 2RT

1 Hunter DJ. Rationing: the case for "muddling through elegantly." *BMJ* 1995;311:811. (23 September.)

2 Klein R. Priorities and rationing: pragmatism or principles? *BMJ* 1995;311:761-2. (23 September.)

Community care and general practice

Focus should be put on barriers that hinder progress

EDITOR,—In their editorial on community care and general practice David Kingdon and Sue Sumners state that the changes in community care in 1993 "seem to have had a smooth take off".¹ This depends on the perspective from which the issues are being explored.

In a study to explore the impact of the reforms of community care on general practice we contacted general practitioners from areas in the former North East Thames region who had previously responded to a questionnaire; 90% reported changing the patterns of their input to the groups of people eligible for community care.² Most of the general practitioners and representatives of social services whom we interviewed believed that clients were not being offered an increased choice in their care regimens; this was central to the aims of the

original white paper.³ Moreover, two thirds of the general practitioners thought that the needs of people with chronic mental illness were either being met badly or were not being met at all. Lastly, almost two thirds of the general practitioners rated their relationship with their local social services as neutral, bad, or non-existent.

Surely, therefore, the continuing challenge of community care is how to enhance joint working to meet patients' increasing needs within this context of poor relations between the core providers of service in the community. Perhaps face to face contact and a joint commitment to closer, more effective working would be a start, as has been suggested in a pilot project in London.^{2,4}

Recently the King's Fund has stated that general practitioners ought to be engaged in community care.⁵ On the basis of our questionnaire data and interviews with general practitioners we believe that this should be amended to "re-engaged," since many general practitioners felt ill prepared before full implementation of the reforms and yet are expected to help augment care in people's homes.

Schemes involving fundholding and joint commissioning need evaluation, but the basic requirement for health and social practitioners to work together is an essential precursor to this. The jury is still out on the overall success of the reforms—necessarily, since they form a 10 year plan. It is surely now prudent, however, to stop talking of a smooth take off and focus more on the barriers that hinder progress and devise strategies to overcome them.

SURINDER SINGH MARGARET LLOYD
Lecturer in general practice Reader in general practice
SARAH WEBB
Researcher

Royal Free Hospital School of Medicine,
Department of General Practice and Primary Care,
London NW3 2PF

1 Kingdon D, Sumners S. Community care and general practice. *BMJ* 1995;311:823-4. (30 September.)

2 Lloyd M, Webb S, Singh S. *General practitioners and the community care reforms*. London: North Thames Regional Health Authority, 1995.

3 Department of Health. *Caring for people: community care in the next decade and beyond*. London: HMSO, 1989. (Cm 849.)

4 Tarbuck H. *Health and Social services development project*. London: City and East London Family Health Services Authority and London Borough of Tower Hamlet Social Services, 1995.

5 Robinson J. *Community care development programme: consultation document, a response from the King's Fund*. London: King's Fund, 1995.

Multidisciplinary, proactive care is needed

EDITOR,—David Kingdon and Sue Sumners highlight the potential of general practitioners to enhance the medical care of some of the most vulnerable members of society.¹ Our experience is that community care cannot be delivered by the primary health care team in isolation: it requires new working relationships with hospital consultants, community nurses, the local authority, and the independent sector. In Fitznells Manor Surgery, which is a fundholding practice with four partners and over 11 000 patients, the process has been activated by the establishment of a 13 person, multidisciplinary proactive care team representative of these liaisons. The team organises the care of vulnerable patients (mainly elderly people) and meets every two months to review key cases, ensuring that all patients in the cohort are monitored continuously. Quality of care is enhanced by assigning to each patient a key worker, who communicates the patient's progress to the team.

Kingdon and Sumners raise an important concern in general practice—namely, that community care places an increased burden on overworked general practitioners. Our proactive care team has provided considerable support for, and greatly reduced the work of, practice staff caring for vulnerable people.

Team meetings enhance awareness of the needs of patients and their carers. The cohesion generated suggests a commonality of purpose that avoids the potential counterproductivity that can arise when different services are involved and reduces the risk of mixed messages to the patient. In addition, meetings provide a forum where trade offs can be made and pragmatic options for each patient identified. The range of services that each discipline offers becomes known, and overreliance on an underresourced profession is avoided. At a time when the market may turn health professionals into competitors the proactive care team encourages people to work together.

As Kingdon and Sumners state, general practitioners are in an ideal position to develop innovative systems for providing community care, but we suggest they cannot do it alone. Seamless quality care requires the harmonisation of the wide range of disciplines needed to provide holistic care. When this is accomplished effectively, creative solutions to difficult problems may emerge that are beyond an individual professional's domain.

ANDREW SINGLETON
Medical anthropologist

Division of General Practice,
St George's Hospital Medical School,
London SW17 0RE

RICHARD MARTIN
General practitioner

SUE MITCHELL
General practitioner

Fitznells Manor Surgery,
Ewell,
Surrey KT17 1TF

1 Kingdon D, Sumners S. Community care and general practice. *BMJ* 1995;311:823-4

Emergency contraception

Care must be taken to ascertain that woman is not already pregnant

EDITOR,—While I acknowledge that every effort should be made to reduce the rate of unwanted pregnancies in Britain, I do not believe that making emergency contraception available over the counter is the right way forward. My main concern, which Jeanette Cayley alludes to only briefly in her editorial,¹ is about women taking contraception when already pregnant. For a general practitioner prescribing emergency contraception, one of the most important aspects of the consultation is to ascertain that the woman is not already pregnant. As Cayley acknowledges, little is known about the effect of emergency contraception if pregnancy is already established. I suspect that some women who think that they may be pregnant may be tempted to take it to try to induce an early abortion.

Surely the way forward is to promote the availability of contraception in general and of emergency contraception in particular, as reducing medical control over it may lead to abuse with unknown consequences.

MARK DINWOODIE
General practitioner

Fair Park Health Centre,
Bath BA1 6EA

1 Cayley J. Emergency contraception. *BMJ* 1995;311:762-3. (23 September.)

Use of the term is erroneous

EDITOR,—It is vital to be accurate when using words and phrases. Jeanette Cayley's editorial is entitled "emergency contraception,"¹ but contraception means "against conception"—that which prevents conception occurring. The object of using the postcoital contraceptive pill is to prevent nidation and further development of the conceptus and to procure an early chemical abortion. One cannot prevent what has already occurred, and