hernia and usually shorter for bilateral or recurrent hernia.

The authors' data on return to work are depressing, with long median times in both the group that had laparoscopic repair (22 days) and the group that had open repair (28 days), and may reflect undue caution on the part of either the patients or their general practitioners. In my experience, with clear and emphatic advice that it is safe to do so sedentary workers return to work in one week and those who do more physical work in two.

Two sentences in the opening paragraph of the section on the outcome measures suggest an initial bias against laparoscopic repair. It seems highly optimistic to have assumed a 0.5% complication rate for open repair (with no supporting data) and highly pessimistic with regard to laparoscopic repair to choose a sample size that would show only a complication rate of 18% or greater.

This study falls into the trap of comparing inexperienced laparoscopic hernia repair with experienced open hernia repair.

ROGER W MOTSON
Consultant surgeon

Colchester General Hospital,
Colchester,
Essex CO4 5JL

1 Lawrence K, McWhinnie D, Goodwin A, Doll H, Gordon A, Gray A, *et al.* Randomised controlled trial of laparoscopic versus open repair of inguinal hernia: early results. *BMJ* 1995;311:981-5. (14 October.)

Lichtenstein technique should be used

EDITOR,—Since the early 1960s a technique of inguinal hernia repair, the Lichtenstein repair, has been available that precludes the use and inherent risks of general anaesthesia. The true Lichtenstein technique is tailor made to suit most patients regardless of their anaesthetic risk, is cost effective, and gives immediate postoperative mobility and reduced postoperative morbidity and pain; patients return to work or normal function after on average 2 to 10 days. In addition, the Lichtenstein repair shows long term recurrence rates of 0.13% at five years.¹ Given these factors, it is alarming that in Kate Lawrence and colleagues' study of only 125 men, in all of whom general anaesthesia was used, the Maloney darn was taken as the current standard surgical technique,² with, we presume, standard recurrence rates of 10-15% at three years.³

The statement that open herniorrhaphy rarely causes severe pain or morbidity is at variance with popular belief and the authors' results, which show that 40-50% of patients who had open repair had not returned to normal function by four weeks. Indeed, when standard techniques are used it is common for patients to return to normal function only after six weeks.⁴

The absence of local data surely reflects auditing techniques and is not an excuse to assume a short term complication rate of 0.5%. It is disconcerting that 1% of patients crossed from laparoscopic to open repair. Moreover, 12% of patients who had laparoscopic repair suffered important early complications, with 5% requiring admission.

Despite these findings the authors propose to undertake longer term, larger studies of laparoscopic techniques, hoping that complication rates decrease as part of a learning curve. They conclude that sample sizes of thousands are needed to detect differences in recurrence rates. Such sample sizes have already been studied: the short and long term benefits of the Lichtenstein technique of hernia repair to patients are conclusive.⁵

As first wave fundholders, we offer Lichtenstein repairs to patients in a primary health care setting. A clinical audit and an audit of patients' perceptions are being undertaken, and we have had a 100% response rate to our anonymised questionnaires. Results for one year will be available in June. In our opinion, further assessment of laparoscopic

herniorrhaphy is unnecessary and the use of archaic standard repairs inappropriate when modern, safer, and more effective techniques are readily available—now in a primary health care setting.

DAVID PELTA

Senior partner

PHILIP LAFFERTY

Surgeon in primary health care

Queensway Surgery,
Southend on Sea,
Essex SS1 2AB

- 1 Amid PK, Shulman AG, Lichtenstein IL. Critical scrutiny of the open "tension-free" hernioplasty. *Am J Surg* 1993;165:369-71.
- 2 Lawrence K, McWhinnie D, Goodwin A, Doll H, Gordon A, Gray A, *et al.* Randomised controlled trial of laparoscopic versus open repair of inguinal hernia: early results. *BMJ* 1995;311:981-5. (14 October.)
- 3 Halverson K, McVay CB. Inguinal and femoral hernioplasty: a 22 year study of the author's methods. *Arch Surg* 1970;101:127-35.
- 4 Shulman AG, Amid P, Lichtenstein I. Returning to work after herniorrhaphy. *BMJ* 1994;309:216-7.
- 5 Amid P, Shulman AG, Lichtenstein I. A critical comparison of laparoscopic hernia repair with Lichtenstein tension-free hernioplasty. *Med J Aust* 1994;161:239-40.

Hernia repair should be individualised to the patient

EDITOR,—The value of Kate Lawrence and colleagues' study comparing laparoscopic with open repair of inguinal hernia lies in the outcome measures chosen and the analysis of cost effectiveness.¹ The results and conclusions drawn are far more meaningful than those drawn from any other comparison between open and laparoscopic hernia repair. Although in their randomised trials of laparoscopic versus open repair Payne *et al* and Millikan *et al* showed an increased cost for laparoscopic repair of \$500 (£330) and \$4500 (£3000) respectively, the economic analysis was not nearly as detailed—hence the wide difference in estimated costs.^{2,3}

Other important data in the authors' study are the demographic details of the patients, including the type of job undertaken and social class. A recent study in Seattle, United States, compared two groups of patients, one of which received workers' compensation while the other had commercial insurance. For the patients with commercial insurance the median duration of postoperative pain was 7.5 days and the median number of days off work was 8.5 days. In contrast, the patients with workers' compensation had pain for 27 days postoperatively and did not return to work for 36.5 days.⁴ Thus selection biases may affect results, and motivation affects clinical outcome. Perceived pain and the ability to return to activity are influenced more by socioeconomic factors than by the procedure or the anatomy involved.

However severe postoperative pain may be, patients should be given sufficient analgesia to permit rehabilitation and recovery. Balanced analgesia, with a combination of non-steroidal anti-inflammatory drugs, opioids, and local anaesthesia (as given in Lawrence and colleagues' study), acts at different points on pain pathways and so maximises analgesia and minimises side effects. Thus pain should not be a major factor hindering recovery in any patient.

Lawrence and colleagues' study began in 1992, before the popularisation of the Lichtenstein operation in Britain. Since then two non-randomised studies in Britain have indicated potential benefits of this procedure compared with other open techniques, such as the Maloney darn.^{5,6} The Lichtenstein operation can reduce costs further since it can be performed under local anaesthesia by a junior surgeon on a day case basis in suitable patients.

Finally, Rutkow has suggested five outcome measures that should be incorporated into any evaluation of hernia surgery: technical difficulty, complication rates, overall rehabilitation, recur-

rence rates, and socioeconomic factors (notably cost benefit and cost effectiveness). To some degree hernia repair needs to be individualised to various clinical, social, and economic situations.

A N KINGSNORTH

Reader in surgery

Department of Surgery,
Faculty of Medicine,
University of Liverpool,
Liverpool L69 3BX

- 1 Lawrence K, McWhinnie D, Goodwin A, Doll H, Gordon A, Gray A, *et al.* Randomised controlled trial of laparoscopic versus open repair of inguinal hernia: early results. *BMJ* 1995;311:981-5. (14 October.)
- 2 Payne JH, Grininger LM, Izawa MT, Podoll EF, Lindahl PJ, Balfour HH. Laparoscopic or open inguinal herniorrhaphy: a randomised prospective trial. *Arch Surg* 1994;129:973-81.
- 3 Millikan KW, Kosik ML, Doolas A. A prospective comparison of transabdominal preperitoneal laparoscopic hernia repair versus traditional open hernia repair in a university setting. *Surg Laparosc Endosc* 1994;4:247-53.
- 4 Salcedo-Wasicek CM, Thirby RC. Postoperative course after inguinal herniorrhaphy: a case-controlled comparison of patients receiving workers' compensation vs patients with commercial insurance. *Arch Surg* 1995;130:29-32.
- 5 Davies N, Thomas M, McIlroy B, Kingsnorth AN. Early results with the Lichtenstein tension-free hernia repair. *Br J Surg* 1994;81:1478-9.
- 6 Kark AK, Kurzer M, Waters KJ. Tension-free mesh hernia repair: review of 1098 cases using local anaesthesia in a day unit. *Ann R Coll Surg Engl* 1995;77:299-304.
- 7 Rutkow IM. The recurrence rate in hernia surgery: how important is it? *Arch Surg* 1995;130:575-7.

Provision of palliative care is suboptimal

EDITOR,—As a palliative physician, I found it upsetting to read Betty Bridger's account of the medical care given to her husband during the last few days of his life.¹ The care was inappropriate and inadequate and resulted in an agonising and painful death, which Bridger cannot eradicate from her mind.

Bridger mentions two means by which such a situation might be averted in the future. She urges that families of terminally ill patients with cancer should be given telephone numbers for emergency care that is available round the clock and that surgeries should have a list of patients who are seriously ill and are at risk. Several lessons can be learnt from this case.

Firstly, patients with a serious illness can deteriorate rapidly, thus confounding any well planned strategy of treatment. Secondly, late referral to palliative care services is all too common and may be due to denial on the part of doctors or relatives. The author says that she had not thought it necessary to explore palliative treatment and help as her husband had managed so well up to that weekend. In retrospect it is easy to say what could have been done better, but it should have been the responsibility of at least one of the doctors or other professionals to ensure that the possibility of palliative care was explored earlier. This would have helped both Bridger and her husband to come to terms with his illness and to make plans for such a situation as arose, and it would have helped Bridger to prepare for her bereavement. Thirdly, the education of all doctors in the basics of palliative care, including communication and listening skills, is still far short of optimal. Fourthly, it is regrettable that patients with chronic illnesses may need to be admitted after hours to hospitals with which they are not familiar. In this case it led to the patient not being admitted at all.

Finally, as doctors, we should recognise when we are out of our depth and should not be afraid to admit this and ask for help.

MICHAEL CUSHEN
Palliative physician

St Elizabeth Hospice,
Ipswich IP3 8LX

- 1 Bridger B. "Quiet consummation have." *BMJ* 1995;311:882. (30 September.)