are) are the only measure available. Moreover, croup scores are useful in hospital clinical practice. As most croup presents outside normal working hours, croup scores are useful to junior paediatricians as a reminder of assessment criteria and as a measure of severity.

M Cronin and R Diedericks express concern that I did not refer to nebulised adrenaline. As the title of my editorial made clear, my primary aim was to draw attention to the role of corticosteroids in croup. The one trial comparing nebulised adrenaline with nebulised corticosteroids showed broadly similar effects, although, as it is reported only in abstract form, interpretation must be limited (D Fitzgerald *et al*, meeting of Australian Thoracic Society, March 1995). Although nebulised adrenaline clearly has a role in the management of croup, I believe that it has no role in mild to moderate croup and (with rare exceptions) should not be used in the community. The role of nebulised adrenaline is to avert or delay the need for intubation, and consequently it should be used only under controlled conditions. Relapse after the use of nebulised adrenaline is well described—indeed, in the study that Cronin and Kiedericks cite, by Kristjansson *et al*, over a third of the children receiving adrenaline relapsed, and the authors recommended that children receiving adrenaline should be observed for two hours. I believe that children who require nebulised adrenaline should be observed where facilities are available for urgent intubation should the need arise.

IOLO DOULL

Senior registrar in paediatric intensive care

Great Ormond Street Hospital for Children NHS Trust,
London WC1N 3JH

1 Jacobs S, Shortland G, Warner J, Dearden A, Gataure PS, Tarpey J. Validation of a croup score and its use in triaging children with croup. *Anaesthesia* 1994;49:903-6.

Deaths of children in house fires

EDITOR,—Ian Roberts's editorial on the deaths of children in house fires omits to mention what is overwhelmingly the most important factor in fatal residential fires—namely, cigarettes.¹ "Smokers' materials" are the source of ignition in nearly half of the house fires in Britain that result in a death. Matches are not included in the government's definition of this term, and in fatal residential fires the term essentially refers to cigarettes. Any discussion of deaths among children because of house fires ought therefore to mention cigarette smoking by members of the victims' households.

In 1994, at a national conference on fatal house fires in Scotland, Professor Timothy Squires noted that "over 40% to almost 50%" of such fires were caused by cigarettes. Now, against the background of an overall decrease in the rate of deaths in fires in the two ensuing years, that percentage has risen to 54% in Scotland.

Recently a scientist who used to be the vice president in charge of research and development for a major cigarette manufacturer in the United States went on public record as saying that the tobacco industry has known "for at least 30 years" how to produce what are popularly called "fire safe cigarettes" (ordinary cigarettes with a drastically reduced propensity to result in a fire).²

Because of requests from the Association for Nonsmokers' Rights, in July 1994 the government released data on the number of children (aged 16 and under) killed in fires caused by smokers' materials in the United Kingdom in 1988-92. The statistical tables (provided by the Home Office's fire statistics section) showed that, on average, 24 children a year died in house fires caused by smokers' materials. Over 90% of such deaths could be avoided if fire safe cigarettes were sold.³

There are many other reasons, too, why cigarette smoking should be taken into consideration when

physical harm to children is being discussed. The chewing and swallowing of cigarettes by toddlers accounts for about 15% of unintentional poisonings of children in Greece. In two thirds of the cases studied both parents were smokers, and in 93% of cases at least one parent was a smoker.⁴ Add to this the effects of smoking during pregnancy and of passive smoking by children⁵ (as well as the effect of smoking on the budgetary constraints of families) and it becomes clear that adults' cigarette smoking should be one of the first factors to be considered by scientists discussing the physical harm done to children in our society.

PHILLIP WHIDDEN
Publications editor

Association for Nonsmokers' Rights,
Edinburgh EH7 4BU

- 1 Roberts I. Deaths of children in house fires. *BMJ* 1995;311:1381-2. (25 November.)
- 2 Wigand J. Affidavit of Jeffrey Wigand in the case of Kearney vs Philip Morris, Inc; civil action No 92-11079-K, 21 July 1995. MA: US District Court for the District of Massachusetts, 1995.
- 3 McGuire A. There's death on the block, there's hope in Congress. *J Public Health Policy* 1987;8:451-4.
- 4 Petridou E, Polychronopoulou A, Kouri N, Karpathios T, Trichopoulos D. Childhood poisonings from ingestion of cigarettes. *Lancet* 1995;346:1296.
- 5 Royal College of Physicians. *Smoking and the young*. London: Royal College of Physicians, 1992.

High levels of polyunsaturated fat may inhibit cancer growth

EDITOR,—Mahmoud Zureik and colleagues report a striking reduction in cancers of all types in association with high levels of polyunsaturated fat in blood cholesterol esters.¹ They reject a causal hypothesis, partly because the inverse association of cancer with polyunsaturated fat is not site specific.

This rejection of a causal hypothesis may be premature. In 1956 Sinclair suggested that high levels of polyunsaturated fat might protect against a range of cancers.² In the past decade investigators from many laboratories have reported that concentrations of polyunsaturated fatty acids that do not harm normal cells inhibit the growth of, or kill, over 40 human cancer cell lines.^{3,4} Thus a plausible biological basis exists for an inhibitory effect of high levels of polyunsaturates on cancer growth.

DAVID F HORROBIN
Chief executive

Scotia Holdings,
Peasmarsh,
Guildford,
Surrey GU3 1NA

- 1 Zureik M, Ducimetière P, Warner J-M, Orssaud G. Fatty acid proportions in cholesterol esters and risk of premature death from cancer in middle aged French men. *BMJ* 1995;311:1251-4. (11 November.)
- 2 Sinclair HM. Deficiency of essential fatty acids and atherosclerosis, etc. *Lancet* 1956;ii:381-3.
- 3 Dippenaar N, Booyens J, Fabri D, Katzef E. The reversibility of cancer: evidence that malignancy in melanoma cells is gamma-linolenic acid deficiency dependent. *S Afr Med J* 1982;62:505-9.
- 4 Begin ME, Ellis G, Das UN, Horrobin DF. Differential killing of human carcinoma cells supplemented with n-3 and n-6 polyunsaturated fatty acids. *J Natl Cancer Inst* 1986;77:1053-62.
- 5 Jiang WG, Hiscox S, Hallett MB, Horrobin DF, Mansel RE, Puntis MCA. Regulation of the expression of E-cadherin on human cancer cells by gamma-linolenic acid (GLA). *Cancer Res* 1995;55:5042-8.

Acoustic neuroma may impair vision

EDITOR,—A Wright and R Bradford's review of the management of acoustic neuroma omits the importance of corneal health and the visual consequences of ignoring early signs of corneal disease.¹ Patients with acoustic neuroma may have three factors leading to corneal compromise: decreased

corneal sensation preoperatively, postoperative facial palsy with associated poor lid closure, and decreased vasomotor tone to the lacrimal gland after seventh nerve palsy. Thus they often have an unprotected, dry eye devoid of sensation.

We have recently reviewed the records of patients who have had conventional surgery (excluding stereotactic radiosurgery) for acoustic neuroma in Leeds and Sheffield. Of 132 patients, 62 had documented involvement of the fifth nerve preoperatively. Eighty five had facial palsy in the postoperative period. Fifty three patients required definitive lid surgery (lateral tarsorrhaphy, central tarsorrhaphy, or medial canthoplasty). Thirteen patients developed corneal vascularisation, stromal opacities, or frank ulceration. A further three had visual impairment as a result of optic atrophy.

The anterior segment disease can be prevented by several prophylactic measures. Firstly, air may be injected into the subcutaneous tissues of the upper lid to produce a mechanical ptosis, which will last for a few days until it is clear whether facial nerve function is preserved. Secondly, preoperative tarsorrhaphy with histoacryl glue (superglue) will provide 10-14 days' cover. This should not be done by inexperienced doctors since glue can seep backwards into the interpalpebral fissure, form a plaque, and abrade the cornea. Superglue can also be reapplied postoperatively if there is a transient facial paresis that seems to be resolving. Thirdly, botulinum toxin can be used to produce a ptosis that will usually last for about three months. This is not risk free since the Bell's phenomenon may be abolished and permanent vertical diplopia may occur.

We recommend early (even preoperative) referral to an ophthalmologist and the use of air or superglue in the immediate perioperative period until the patient is fit enough for full slit lamp examination. Ocular lubricants should be used freely and frequently. Patients should be told preoperatively that surgery for an acoustic neuroma can potentially impair vision in the ipsilateral eye.

NEIL K ROGERS
Registrar

Department of Ophthalmology,
Royal Hallamshire Hospital,
Sheffield S10 2JF

CHRISTOPHER S BRAND
University lecturer

Department of Ophthalmology and Orthoptics,
Royal Hallamshire Hospital

- 1 Wright A, Bradford R. Management of acoustic neuroma. *BMJ* 1995;311:1141-4. (28 October.)

Plasma cholesterol response to dietary saturated fat

Conclusions are based on circular arguments

EDITOR,—Charlotte Cox and colleagues report a double crossover trial looking at individual variation in the response of plasma cholesterol concentrations to dietary saturated fat.¹ Responsiveness to diet was calculated for each subject as the difference between total cholesterol concentration during two types of diet. Consistent responders were then defined as those in whom the difference between the two differences was within one standard deviation of the mean for all participants; the remainder were defined as variable responders. Not surprisingly, the authors found subjects who fulfilled their definition, which led them to conclude that, indeed, a group of consistent responders exists. Their tautological and circular arguments then continue.

Firstly, the authors note a "striking" correlation in the changes between the first and second crossover periods. We would have expected this: it merely confirms the definition of consistency in response. The authors then postulate the existence