discounting was not appropriate. The paper does, however, provide the necessary data for relating these costs to future benefits using any selected discount rate.

NICHOLAS J WALD
Professor
PHILIP MURPHY
Computer programmer

PHILIPPA MAJOR
Research assistant
CAROL PARKES
Lecturer

CHRIS FROST
Statistician

Cancer Research Campaign Cancer Screening Research Group,
Wolfson Institute of Preventive Medicine,
St Bartholomew's Hospital Medical College,
London EC1M 6BQ

JOY TOWNSEND
Senior scientist

Medical Research Council Epidemiology and Medical Care Unit,
Wolfson Institute of Preventive Medicine

Low weight gain in infancy and suicide in adult life

The evidence doesn't support the theory

EDITOR,—David Barker and colleagues suggest that suicide in adulthood may be a direct consequence of low weight during the first year of life.¹ This deterministic theory of development is based on the idea of fetal programming of neuroendocrine function, which is a common cause of poor weight gain in infancy and of depression, leading to suicide, in later life. This is in contrast with modern studies of psychosocial risk, which suggest that development is a non-linear, probabilistic process—and is not based on linear, deterministic principles.²

The evidence presented in the paper provides very little support for their theory. Firstly, no association was found between birth weight and the risk of suicide. Presumably, growth during fetal life would be expected to be affected by a putative abnormality of endocrine function arising in utero. The association between weight and suicide was seen only at age 1 year, which suggests that, for some reason, these infants displayed a relative failure to thrive. A possible cause of poor weight gain may have been maternal depression—which was not mentioned in the paper. Simply recording the method of feeding (breast or bottle) would not have detected subtle difficulties in feeding and problems in mother-child attachment. Murray *et al* showed that postnatal depression in women increases the vulnerability of male children to the adverse risks of low social class.³ Controlling for social class did weaken the relation between weight gain and the risk of suicide. If this study was being conducted today, a variety of possible psychosocial confounders such as maternal depression and the quality of the mother-infant relationship would need to be measured before a relation between infant weight gain and subsequent psychiatric disorder could be suggested.

Secondly, the association between abnormal endocrine function and depression does not prove that endocrine dysfunction is a primary cause of depression. In fact, considerable evidence now points to the possibility that endocrine dysfunction is secondary to psychosocial stress and increased concentrations of cortisol. Inadequate psychosocial stimulation may itself be a cause of abnormal endocrine function in infancy, as well as increasing the risk of reduced nutritional intake. If there is a primary causal link between endocrine dysfunction and depression or suicide, an increased risk of depression or suicide in those individuals with short stature due to primary growth hormone deficiency would be expected.

Finally, to say that each kilogram decrease in weight gain results in an increased risk of suicide of 45% in males (an odds ratio of 1.45) is potentially misleading, as it seems to inflate the importance of what is a very small risk in absolute terms. Given the low prevalence of suicide in the study

population (0.27%), for each kilogram decrease in weight gain the attributable risk of suicide in males increased by only 0.12%.

CHRIS HOLLIS
Clinical lecturer

Department of Child and Adolescent Psychiatry,
Institute of Psychiatry,
London SE5 8AF

- 1 Barker DJP, Osmond C, Rodin I, Fall CDH, Winter PD. Low weight gain in infancy and suicide in adult life. *BMJ* 1995;311:1203. (4 November.)
- 2 Rutter M. Epidemiological approaches to developmental psychopathology. *Arch Gen Psychiatry* 1988;45:486-95.
- 3 Murray L. The impact of post-natal depression on infant development. *J Child Psychol Psychiatry* 1992;33:543-62.

Authors' reply

EDITOR,—Chris Hollis is incorrect in several respects. To suggest that prenatal programming of key endocrine axes is linked to later disease is not deterministic.¹ Persisting changes in endocrine function arising in utero do not necessarily affect birth weight—for example, babies with growth hormone deficiency have normal birth weight but fail to grow postnatally.² In our study, adjustment for social class did not importantly weaken the association between suicide and weight gain in infancy, since the average weight at 1 year of people who committed suicide was 0.4 kg less, before and after adjustment, than that of people who did not. Whether expressed as a relative or an absolute risk, a 1 kg decrease in weight at 1 year is associated with a 45% increase in the risk of suicide, and the significance of this finding is not diminished because suicide is an uncommon cause of death.

We are currently studying a sample of men and women in Hertfordshire, linking their depression scores with fetal and infant growth, taking account of their current psychosocial status and their postnatal development, recorded by health visitors at the time.

D J P BARKER
Director
C OSMOND
Statistician

I RODIN
Registrar
C H D FALL
Epidemiologist
P D WINTER
Computing manager

MRC Environmental Epidemiology Unit,
Southampton General Hospital,
Southampton SO16 6YD

- 1 Wulff HR. Mothers, babies and disease in later life. *Lancet* 1995;345:309.
- 2 Gluckman PD, Gunn AJ, Wray A, Cutfield WS, Chatelain PG, Guilbaud O, *et al*. Congenital idiopathic growth hormone deficiency is associated with prenatal and early postnatal growth failure. *J Pediatr* 1992;121:920-3.

Steroids in the management of croup

Croup scores are rarely used in general practice

EDITOR,—Iolo Doull's recommendations about the role of steroids in the management of croup are controversial and deserve comment.¹ An earlier publication that reviewed essentially the same reports was much more circumspect in its recommendations about nebulised steroids.²

There is now good evidence for a biological effect of oral, intravenous, and nebulised steroids in varying degrees of croup, but the evidence for a clinically important effect of nebulised steroids is less convincing. Studies relying on improvements in the croup score (which assigns numbers to clinical signs) as the main outcome measure^{3,4} can be criticised on the basis that the validity, reliability, and responsiveness of the score (in all its various forms) have not been evaluated. Hence, while these studies have shown a significant difference between groups given nebulised steroids and those given placebo, the extent to which these

changes in the score reflect clinically important improvements in an individual child is unclear. Croup scores are rarely used in clinical practice.

Doull suggests that croup severe enough to warrant intervention could be treated at home with oral or nebulised steroids by general practitioners, and recommends that the child should be reviewed two to four hours later. This may be reasonable as an emergency measure while urgent admission to hospital is being arranged but in isolation could be dangerous.

Finally, Landau and Geelhoed remind us that 50-80% of children with croup who receive placebo in trials show considerable spontaneous improvement.² Any new management strategies, particularly for mild croup, need to be evaluated in the knowledge that most children recover without active treatment.

P I MACFARLANE
Consultant paediatrician
S SURI
Registrar in paediatrics

Department of Child Health,
Rotherham District General Hospital,
Rotherham,
South Yorkshire S60 2UD

- 1 Doull I. Corticosteroids in the management of croup. *BMJ* 1995;311:1244. (11 November.)
- 2 Landau LI, Geelhoed GC. Aerosolized steroids for croup. *N Engl J Med* 1994;331:322-3.
- 3 Husby S, Agertoft L, Mortensen S, Pedersen S. Treatment of croup with nebulised steroid (budesonide): a double blind, placebo controlled study. *Arch Dis Child* 1993;68:352-5.
- 4 Klassen TP, Feldman ME, Watters LK, Sutcliffe T, Rowe PC. Nebulised budesonide for children with mild to moderate croup. *N Engl J Med* 1994;331:285-9.

Nebulised adrenaline is also useful

EDITOR,—We are concerned that Iolo Doull's editorial¹ seems to disregard the importance of nebulised adrenaline in children with croup and that some general practitioners or even paediatricians may consequently consider steroids alone to be adequate management for those with moderate or severe airway obstruction. The safety and efficacy of nebulised adrenaline with and without oral steroids have been shown.^{2,3} The effect of nebulised adrenaline may be relatively short lived, but it usually provides prompt relief of airway obstruction and in our experience has often prevented the need for intubation, especially early after admission.

Clearly, a controlled clinical trial is needed to evaluate the effect of nebulised budesonide compared with that of nebulised adrenaline. Both are effective forms of treatment but differ in their mode of action and their place in the management of croup.

M CRONIN
Consultant paediatrician
R DIEDERICKS
Consultant paediatrician

Epsom General Hospital,
Epsom,
Surrey KT19 8PB

- 1 Doull I. Corticosteroids in the management of croup. *BMJ* 1995;311:1244. (11 November.)
- 2 Ledwith CA, Shea LM, Mauro RD. Safety and efficacy of nebulized racemic epinephrine in conjunction with oral dexamethasone and mist in the outpatient treatment of croup. *Ann Emerg Med* 1995;25:331-7.
- 3 Kristjansson S, Berg-Kelly K, Winso E. Inhalation of racemic adrenaline in the treatment of mild and moderately severe croup. *Acta Paediatr* 1994;83:1156-60.

Author's reply

EDITOR,—P I Macfarlane and S Suri question both the validation of croup scores and the scores' usefulness. They do not, however, mention what they consider to be a more appropriate measure of croup. The Syracuse croup score has been validated on the unit to admit a patient to an intensive therapy unit.¹ In the absence of a more objective measure, croup scores (flawed as they