replace a defective gene with a normal one may be thwarted by the knowledge that the single gene disease affecting a particular patient may not have been caused solely by the malfunctioning of that single gene.

Secondly, genetic complexity has economic implications. Insurance companies in the United States are already using altered genotype as an indication of a pre-existing condition and excluding people from health insurance coverage on that basis.⁶ This policy is illogical as well as unfair, as many people with the altered genotype will never become ill.

Finally, an appreciation of genetic complexity has implications for the understanding of multifactorial human behavioural traits such as intelligence, addiction, and criminality. Critics of biological determinist theories of human behaviour have long argued that, even assuming that some human behaviours have a genetic basis, interactions among genes and between genes and the environment must be important in their development. The complexities underlying "simple" single gene diseases like Huntington's disease and cystic fibrosis suggest that the genetic complexities underlying human behaviour are even greater than the critics of biological determinism imagined. Ten years ago most people believed that the presence of the Huntington's disease or cystic fibrosis genotype guaranteed the presence of the disease. Now that this belief has been shaken, how much less likely that multifactorial traits, with their myriad additional

complexities, are inevitable outcomes of particular genotypes.

Recent advances in human genetics have been spectacular. However, overinterpretation of these advances has reinforced a simplistic model of the genetic determination of traits. This model has encouraged premature application of genetics to medicine, the unjustified use of genetic information by insurance companies and employers, and the misinterpretation of genetic results by policy analysts in their discussion of social problems. Scientists and journalists should be cautious and circumspect when discussing genetic research and its implications.

JOSEPH S ALPER
Professor and director

Department of Chemistry and Center for
Genetics and Public Policy,
University of Massachusetts-Boston,
Boston, MA 02125,
USA

- 1 Wolf U. The genetic contribution to the phenotype. *Hum Genet* 1995;95:127-48.
- 2 Gusella JF, MacDonald ME. Huntington's disease and repeating trinucleotides. *N Engl J Med* 1994;330:1450-1.
- 3 Kremer B, Goldberg P, Andrew SE, Theilmann J, Telenius H, Zeisler J, et al. A worldwide study of the Huntington's disease mutation—the sensitivity and specificity of measuring CAG repeats. *N Engl J Med* 1994;330:1401-6.
- 4 Walsh MJ, Tsui L-C, Boat TF, Beaudet AL. Cystic fibrosis. In: Scriver CR, Beaudet AL, Sly WS, Valle D, eds. *The metabolic and molecular bases of inherited diseases*. New York: McGraw-Hill, 1995:3799-876.
- 5 Kieseewetter S, Macek M, Davis C, Curnstin SM, Chu C-S, Graham C, et al. A mutation in CFTR produces different phenotypes depending on chromosomal background. *Nat Genet* 1993;5:274-8.
- 6 Billings PR, Kohn MA, de Cuevas M, Beckwith J, Alper JS, Natowicz MR. Discrimination as a consequence of genetic testing. *Am J Hum Genet* 1992;50:476-82.

Data protection, health care, and the new European directive

Extension of data protection to paper records is logical but problematic

Last summer a directive on data protection was agreed by the European Union's Council of Ministers and is now in force in all member states.¹ Although it confirms the general principles already established by the United Kingdom's Data Protection Act of 1984,² it has particular implications for health care information. It provides for the collection and processing of sensitive personal information for health care purposes and of epidemiological, public health, and scientific data for research. Its biggest impact, however, will be its extension of data protection to manually held records containing personal information.

The directive has to be implemented for computer based data systems within three years. The United Kingdom's effective Data Protection Act enforcement programme has ensured that health care computer based systems which have achieved compliance in Britain should have little difficulty complying with the directive. Like the existing act, the directive specifies that data must be: obtained and processed fairly and lawfully; held and processed only for specified, explicit and legitimate purposes; adequate, relevant, and not excessive; accurate, complete, and, where necessary, up to date; held no longer than necessary; accessible to the subject, who has rights to have the information corrected or erased; and kept secure and confidential.

Unlike the Data Protection Act, however, the directive provides for compensation for damages and establishes a right to privacy. Details of legal sanctions, civil or criminal, will be left to individual member states, and this needs to be legislated for by the British government; until it does so the Data Protection Act remains in force.

The big change is that all patient health care records, whether held on computer or manually, will be subject to the directive. Fortunately, member states have 12 years to

implement the provisions for manual systems, though "further active processing" of manual records should conform before then. Personal rights to access and correct information are established now.

These changes apply not only to hospitals but also to paper records in general practice. Patient records will have to adhere to the basic principles of data protection. There can be no enforcement until laws are passed; but British civil cases probably would be greatly influenced now. At present many patient records breach the data protection principles: many are inadequate, irrelevant, and excessive, and organisations fail to ensure their accuracy, completeness, and currency. Although the legitimacy of collecting patient data for health care seems firmly established, it would be wise for organisations to determine explicitly their purpose well in advance of legislative implementation of the directive. Likewise, it should not be hard to justify keeping patient records for a long time, but the justification should be explicit and prepared in advance of complaints. Use of the data for historical, statistical, or scientific purposes is allowed so long as privacy is safeguarded.

A major problem for manual record systems will be the requirement to implement technical and organisational measures to protect personal information against accidental or unlawful loss³: manual systems are much harder to protect from these than computer systems. Given the many millions of manual patient records held and the poor condition of many of them even 12 years may be a short time to achieve compliance with the directive. The directive addresses, though it does not completely resolve, contentious data protection issues such as the use of clinical data for research and the collection of personally sensitive yet clinically important information.⁴⁻⁶ Nevertheless, it does seem relatively

amenable to the needs for such information within health care. The data protection registrar has welcomed the directive, and there is no reason to suspect that its implementation in Britain will be anything but speedy and wholehearted.⁷

The Data Protection Act has deeply affected the attitudes and behaviour of organisations and people concerned in the computer storage and processing of personal data over the past 10 years. Fortunately, the act mainly enforced what was already good professional practice, and the overall effect has been to strengthen the progress of computer based information systems. The directive's extension of data protection to manual record systems is logical, and its effect on organisations holding manual patient records is likely to be at least as profound as that of the Data Protection Act and, in

the long run, equally beneficial. Getting to that point, however, may prove to be painful and expensive.

M F SMITH

Professor of health informatics

Keele University,
Keele ST5 5AY

- 1 On the protection of individuals with regard to the processing of personal data and on the free movement of such data. Brussels: European Union, The Council, 1995 (amended June 1995).
- 2 Data protection act 1984: the guideline series and guidance notes (guidelines 1-8). Wilmslow, Cheshire: Office of the Data Protection Registrar, 1992.
- 3 Setting the records straight. London: Audit Commission, 1995.
- 4 NHS contract minimum data sets. Wilmslow, Cheshire: Data Protection Registrar, 1993.
- 5 Lyng L. European directive on confidential data: a threat to epidemiology. *BMJ* 1994;308:490.
- 6 Westrin CG, Nilsson T. The ethics of data utilisation: a comparison between epidemiology and journalism. *BMJ* 1994;308:522-3.
- 7 European Union data protection directive. *Insights: practical guidance on data protection*. Wilmslow, Cheshire: Data Protection Registrar, 1995.

Thrombosis, phlebitis, and varicose veins

Varicose veins don't confer significant risk of deep vein thrombosis

Varicose veins have been placed high on the list of risk factors for deep vein thrombosis both in British and European guidelines.^{1,2} But warnings of a high risk of thrombosis for people with varicose veins are based on limited evidence³⁻⁷ and raise some important questions for those advising and treating them. Firstly, what is the risk of deep vein thrombosis in patients with varicose veins, either in their normal daily lives or when other risk factors are present? Secondly, what is the risk of deep vein thrombosis in patients having surgery for varicose veins, and what prophylaxis should be used?

The evidence for an association between varicose veins and deep vein thrombosis is based solely on the results of studies on cohorts of patients undergoing abdominal or pelvic surgery.³⁻⁷ These documented varicose veins as an independent risk factor for thrombosis. However, there is no evidence that varicose veins on their own cause deep vein thrombosis: duplex ultrasound examination is now commonly used to investigate varicose veins and does not reveal "incidental" deep vein thromboses.⁸ Fears that varicose veins pose a significantly increased risk of spontaneous thrombosis in women taking the contraceptive pill or hormone replacement therapy are also unsubstantiated. (These reassurances apply only to patients with primary varicose veins and not to the small minority whose varicose veins are the result of previous deep vein thrombosis, for whom the risks of further thrombosis are higher.)

Is there a special risk of deep vein thrombosis for patients with varicose veins who develop superficial thrombophlebitis? Some studies have shown deep vein thromboses in limbs with phlebitis,^{8,9} but these thromboses have mostly occurred in circumstances of special risk, such as after an operation or illness⁹⁻¹¹ and in those with history of deep vein thrombosis.^{10,12} Older patients and those with bilateral phlebitis may also be at increased risk.¹⁰ Phlebitis in legs without varicose veins is more likely to be associated with deep vein thrombosis.¹⁰ Duplex scanning of the deep veins is probably worth while in these higher risk patients. A special risk of deep vein thrombosis also exists if phlebitis progresses proximally up the long saphenous vein in the thigh,^{11,13,14} and some surgeons advise emergency ligation of the saphenofemoral junction to prevent extension of thrombosis into the femoral vein and possible pulmonary embolism.^{11,13} By contrast, localised phlebitis of varicose veins in fit young people is unlikely to be associated with deep vein thrombosis, and suggestions that they should all have their deep veins scanned are unjustified.^{8,9}

A special dilemma exists for surgeons operating on varicose veins, because there is no good evidence to show a special risk of deep vein thrombosis. Nevertheless, the existence of guidelines¹² that list varicose veins as an important risk factor for deep vein thrombosis, means that there is a medicolegal threat if a patient develops venous thromboembolism after varicose vein surgery and "demonstrable prophylaxis" was not used. Exactly what constitutes demonstrable prophylaxis is open to debate. Subcutaneous heparin is of proven benefit in other areas of surgery, but may worsen postoperative bruising and haematoma,¹⁵ which are the most common and troublesome sequelae of varicose vein operations. In response to a recent questionnaire, only 29% of 289 vascular surgeons considered varicose veins a special risk factor for deep vein thrombosis, and only 12% used heparin routinely for varicose vein surgery.¹⁶ Some used heparin selectively in obese patients, elderly patients, or those with recurrent varicose veins, and most did so only for patients with a history of deep vein thrombosis and those on the contraceptive pill. Other proved methods of prophylaxis¹⁵ such as antiembolism stockings and intermittent compression are impractical for use on the affected leg during varicose vein operations. Most surgeons recommend antiembolism stockings and early mobilisation after operation.¹⁶

Reliable data on the incidence of deep vein thrombosis after varicose vein surgery are needed. Until these are available, thromboembolic prophylaxis will be determined more by medicolegal considerations than by clinical judgment. There is good evidence for an increased risk of deep vein thrombosis at the time of major abdominal or pelvic surgery, but at other times people with varicose veins are unlikely to be at special risk.

BRUCE CAMPBELL

Consultant in vascular and general surgery

Royal Devon and Exeter Hospital,
Exeter EX2 5DW

- 1 Thromboembolic Risk Factors (THRIFT) Consensus Group. Risk of and prophylaxis for venous thromboembolism in hospital patients. *BMJ* 1992;305:567-74.
- 2 European Consensus Statement. *Prevention of venous thromboembolism*. London: Med-Orion, 1992.
- 3 Kakkar VV, Howe CT, Nicolaides AN, Renney JTG, Clarke MB. Deep vein thrombosis of the leg: is there a "high risk" group? *Am J Surg* 1970;120:527-30.
- 4 Nicolaides AN, Irving D. Clinical factors and the risk of deep vein thrombosis. In: Nicolaides AN, ed. *Thromboembolism: aetiology, advances in prevention and management*. Lancaster: Medical and Technical Publishing, 1975:193-204.
- 5 Crandon AJ, Peel KR, Anderson JA, Thompson V, McNicol GP. Postoperative deep vein thrombosis: identifying high risk patients. *BMJ* 1980;281:343-4.
- 6 Lowe GDO, Osborne DH, McArdle BM, Smith A, Carter DC, Forbes CD, et al. Prediction and selective prophylaxis of venous thrombosis in elective gastrointestinal surgery. *Lancet* 1982;i:109-12.