istration is focusing on this in its "health in the next millennium" work programme, as is the Medical Research Council in its Gambia project, and the Liverpool School of Tropical Medicine now incorporates non-communicable diseases into its diploma courses. We suggest that the specialty of tropical medicine should shift towards a broader specialty of "health in developing countries," with the multidisciplinary inputs that the authors recommend, rather than be subsumed into infectious diseases.

D McLARTY
Professor of medicine

Muhimbili Medical Centre,
PO Box 65243,
Dar es Salaam,
Tanzania

K G M M ALBERTI
Professor of medicine

N UNWIN
Lecturer in public health medicine

Human Diabetes and Metabolic Research Centre,
Medical School,
University of Newcastle upon Tyne,
Newcastle upon Tyne NE2 4HH

1 De Cock KM, Lucas SB, Mabey D, Parry E. Tropical medicine for the 21st century. *BMJ* 1995;311:860-2. (30 September.)

Genitourinary medicine should remain a separate specialty

EDITOR,—While we agree with the general thrust of Kevin M De Cock and colleagues' thesis—that tropical medicine should be subsumed within the specialty of infectious diseases—we do not agree with their assertion that "the uniquely British concept of a separate career structure for specialists in sexually transmitted diseases, divorced from the broader subject of infectious diseases, is ill adapted to a future within Europe and the broader world."¹ Britain—with 236 genitourinary medicine clinics countrywide, all run by appropriately trained physicians—now has one of the lowest prevalences of sexually transmitted diseases in western Europe (bettered only by Sweden).

The existence of this infrastructure has also enabled genitourinary medicine physicians to provide most of the care for people infected with HIV. Departments of genitourinary medicine are well placed to do this, with their policy of open access, strict code of confidentiality, and access to health promotion facilities. The training of genitourinary medicine physicians has been modified to take account of this new area of health provision, but we acknowledge that a broader experience of infectious diseases would be useful.

Many genitourinary medicine clinics now provide additional services for family planning, psychosexual counselling (including for erectile dysfunction), and colposcopy—that is, a complete sexual health service. Colleagues overseas, where genitourinary medicine is not a single specialty, envy the benefits of a single discipline; they do not urge us to join with others.

J S BINGHAM
Consultant physician

St Thomas's Hospital,
Guy's and St Thomas's Hospital Trust,
London SE1 7EH

1 De Cock KM, Lucas SB, Mabey D, Parry E. Tropical medicine for the 21st century. *BMJ* 1995;311:860-2. (30 September.)

Clinical care in resource poor countries is often provided by people outside the medical establishment

EDITOR,—We believe that the second topic in Kevin M De Cock and colleagues' analysis of the future role of tropical medicine in Northern countries—namely, the role and future of clinical services in resource poor countries—needs more debate.¹ The authors note that the World Bank's *World Development Report 1993* reaffirms the

importance of good clinical care.² They fail to emphasise that effective provision of the "essential clinical package" involves people outside medicine.

Firstly, in large areas of resource poor countries clinical care is provided by nurses and paramedical workers, who often work independently of physicians. Secondly, good clinical care includes nursing care and effective communication and counselling, especially for chronic diseases and conditions that require substantial behavioural change to prevent their recurrence. Skills to develop communities are also required if services are to have widespread impact. Training in these skills is rarely emphasised in Northern medical institutions. Thirdly, the development of clinical skills is wasted if resources and support systems are lacking or mismanaged. Management information systems are needed to monitor inputs, and surveillance systems are needed to monitor activities and diseases; these systems are usually not regarded as part of "tropical medicine." Fourthly, activities such as family planning and treatment of common childhood diseases, included in the "essential clinical services" package, could be (and often are) performed through channels outside the medical establishment.

Clinical services are, nevertheless, an important component of international health, and the development of a cadre of infectious disease specialists would benefit countries in the North and South alike. As De Cock and colleagues point out, experience in resource poor settings is valuable for clinicians specialising in infectious diseases in Europe; yet Northern institutions regard sending doctors to hospitals in tropical countries as part of overseas aid. Periods in developing countries may more appropriately be regarded as part of a person's training, which should contribute to accreditation but could be paid for in the same way that we expect those from resource poor countries to pay fees for training in the North.

Training in tropical medicine should reflect the changes in clinical priorities highlighted by De Cock and colleagues, emphasise management and communication skills, and give increased emphasis to training and supporting nurses and paramedics. Northern institutions and donors should work with colleagues from resource poor countries to determine how collaborative programmes, based in both the South and the North, can improve the global control of major diseases. Such programmes should recognise the extent to which the North benefits from the experience of the South.

FELICITY CUTTS
Senior lecturer in epidemiology
ANNE MILLS
Professor of health economics and policy

London School of Hygiene and Tropical Medicine,
London WC1E 7HT

1 De Cock KM, Lucas SB, Mabey D, Parry E. Tropical medicine for the 21st century. *BMJ* 1995;311:860-2. (30 September.)
2 World Bank. *World development report 1993. Investing in health*. New York: Oxford University Press, 1993.

Videotaping of general practice consultations

EDITOR,—Jacqueline E Bain and Neil S D Mackay¹ suggest that the consent rate of 91% in our study of videotaped consultations in general practice² is evidence of coercion and state that consent rates of 4-10% are to be expected when coercion is removed. In neither paper that they quote in support of these figures were patients invited to be videotaped. In one study the authors asked patients to speculate how they might feel if so invited.³ In the other study patients were given leaflets inviting them to volunteer.⁴ Those patients who did not care one way or the other were included in the 90% who were claimed to have rejected videotaping. The suggestion that this

study produced a consent rate of 10% is not true: people cannot consent to something unless they have been asked. These points were made in 1987 in response to the original article.^{5,6}

We believe that many patients agree to videotaping out of a sense of altruism and an idea that they may be helping to train better doctors. Clearly, in some circles this could be seen as coercion. Bain and Mackay seem to consider it to be coercive merely to invite a patient to allow his or her consultation to be videoed. There has been a long history in Britain of patients agreeing to contribute to medical education at both undergraduate and postgraduate levels—for example, by being seen by medical students or acting as cases in examinations.

Since coercion in this context is impossible to define or measure we concentrated on outcome measures on the basis that it was important to determine whether videotaping had an adverse effect on the consultation. If 80% of our sample of patients did feel unhappy about the videotaping it is astonishing that the patients who were videoed reported the same satisfaction as those who were not videoed. Our patients did not produce an "acquiescence response set," as is shown by our results, and our statistics showed that it was extremely unlikely that we had missed a difference in satisfaction between the two groups.

L M CAMPBELL
Assistant adviser (assessment)

T S MURRAY
Regional adviser

West of Scotland Postgraduate Medical
Education Board,
Glasgow G12 9LX

F M SULLIVAN
Senior lecturer

Department of General Practice,
University of Glasgow,
Glasgow

- 1 Bain JE, Mackay NSD. Videotaping of general practice consultations. *BMJ* 1995;311:952. (7 October.)
- 2 Campbell LM, Sullivan FM, Murray TS. Videotaping of general practice consultations: effect on patient satisfaction. *BMJ* 1995;311:236. (22 July.)
- 3 Bain JE, Mackay NSD. Patients' assessment of trainee general practitioners. *Med Educ* 1995;29:91-6.
- 4 Servant JB, Mathieson JAB. Video recording in general practice: the patients do mind. *J R Coll Gen Pract* 1986;36:555-6.
- 5 Boardman AP, Craig TKJ. Video recording in general practice. *J R Coll Gen Pract* 1987;37:180-1.
- 6 Mackay HAF. Video recording in general practice. *J R Coll Gen Pract* 1987;37:181.

Chemotherapy in non-small cell lung cancer

Large trial will reduce uncertainty

EDITOR,—The recent meta-analysis of the role of chemotherapy in non-small cell lung cancer suggests that treatments containing cisplatin improved survival.¹ The effect of chemotherapy seemed to be consistent across all subgroups (stage of disease, age, sex, performance status, and tumour histology). Because of the heterogeneity of the treatments used in the various trials, however, it was not possible to recommend a particular chemotherapy regimen or to be certain of the size of the benefit. Most importantly, it was not possible to state reliably whether a short course of a widely used cisplatin based treatment would improve survival, or by how much. The economic aspects and quality of life were also not addressed, for obvious reasons.

Considerable doubt therefore remains over the usefulness of combination chemotherapy in non-small cell lung cancer. To answer these questions we have designed a large randomised trial to compare chemotherapy with no chemotherapy in each of the four settings studied in the meta-analysis (surgery, surgery plus radiotherapy, radical radiotherapy, and supportive care). The recommended chemotherapy regimen is three

cycles of either mitomycin, ifosfamide, and cisplatin or mitomycin, vinblastine, and cisplatin.

Nine thousand patients will be required for the trial to give a reliable and clinically relevant result. Potentially all patients with non-small cell lung cancer will be eligible. Similar large trials have been conducted in breast and ovarian cancer, with considerable success. The "big lung trial" will be the first large scale randomised trial to ask this basic but important question about the treatment of this major disease.

On behalf of the steering committee of the big lung trial

R J STEPHENS

Statistician

M K B PARMAR

Statistician

MRC Cancer Trials Office,
Cambridge CB2 2BW

R L SOUAMI

Professor of clinical oncology

Department of Oncology,
University College London Medical School,
Middlesex Hospital,
London W1N 8AA

S SPIRO

Clinical director of medicine

University College London Hospitals,
Middlesex Hospital

1 Non-small Cell Lung Cancer Collaborative Group. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. *BMJ* 1995;311:899-909. (7 October.)

Quality of life was ignored in meta-analysis

EDITOR,—The Non-small Cell Lung Cancer Collaborative Group's meta-analysis suggests that chemotherapy regimens that contain cisplatin may improve survival in patients with non-small cell lung cancer.¹ It was a disappointment to find that, with a median increase in survival of 1.5 months in the group given supportive care, no data are presented on the impact of chemotherapy on patients' quality of life. Surely the time has come when no studies should be funded unless they include an assessment of quality of life, and the Non-small Cell Lung Cancer Collaborative Group should include such an assessment in future meta-analyses.

IRENE HIGGINSON
Senior lecturer

London School of Hygiene and
Tropical Medicine,
London WC1E 7HT

1 Non-small Cell Lung Cancer Collaborative Group. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised control trials. *BMJ* 1995;311:899-909. (7 October.)

It may not be ethical to ask patients to participate in future trials

EDITOR,—Meta-analysis suggests that chemotherapy may have a role in the treatment of non-small cell lung cancer.¹ The authors note that the place of such treatment is uncertain, given the toxicity of platinum based combinations and the small survival advantage. An accompanying editorial suggests that, as the number of people with lung cancer worldwide is large, even a small individual chance of benefit is worth while in terms of population health gain.²

In the developing world, where the epidemic of lung cancer is just beginning,³ health care resources are insufficient for any treatment to be offered to most patients. In England and Wales, if 80% of registered lung cancers are of non-small cell type and radical primary treatment with platinum based chemotherapy was given to all patients aged under 70 with localised and locally advanced disease (estimated at 40% of the total number), there would be an extra 199 survivors at two and five years, who might be potentially cured. If platinum based chemotherapy was also given to 80% of those aged under 70 with advanced disease (the

remaining 20% being considered unfit for active treatment) then 681 more patients might be alive at the end of one year than would otherwise be the case; but for most of these death is only postponed.

For this survival gain to be achieved, 12 492 patients would have to receive pulsed treatment with emetic drugs as an inpatient for up to six months, though some would also experience relief of their symptoms or psychological benefit. Patients with the best prognosis with "standard" treatment may be harmed by the addition of optimal, platinum based chemotherapy (95% confidence intervals for mortality hazard ratios 0.74 to 1.02 and 0.79 to 1.11, respectively (table II in the meta-analysis). The authors conclude with a plea for future trials to be fewer in number and larger in size; hopefully, these will also include outcomes other than survival, as treatment for most patients is palliative rather than curative. A decade ago, however, most experts in lung cancer who responded to a postal questionnaire said that they would decline randomisation into trials of chemotherapy.⁴ Unless the meta-analysis substantially alters professional opinion will it be ethical to ask patients to participate in such trials?

MAGGIE HARDING

Senior registrar in public health medicine

Health Care Evaluation Unit,
Department of Public Health Sciences,
St George's Hospital Medical School,
London SW17 0RE

- 1 Non-small Cell Lung Cancer Collaborative Group. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. *BMJ* 1995;311:899-909. (7 October.)
- 2 Carbone DP, Minna JD. Chemotherapy for non-small cell lung cancer. *BMJ* 1995;311:889-90. (7 October.)
- 3 Peto R. Smoking and death: the past 40 years and the next 40. *BMJ* 1994;309:937-8.
- 4 Mackillop WJ, Ward GK, O'Sullivan B. The use of expert surrogates to evaluate clinical trials in non-small cell lung cancer. *Br J Cancer* 1986;54:661-7.

Allocating census data to general practice populations

EDITOR,—We wish to reply to the two letters¹ commenting on our paper on allocating census data to a general practice population.² Paul Aveyard proposes using the method of Bland and Altman rather than correlation to measure agreement between the number of patients aged over 65 obtained from the family health services authority's register and that calculated from the postcodes of patients and the proportion in their census area. Ronan A Lyons and colleagues suggest that better results could have been obtained in their area, where they achieve a high proportion of matches between postcodes and census areas.

We presented the correlation coefficients since they are what most others in this field (for example, Majeed *et al*)³ have used to justify the use of census data, but we then went on to show, by using a graph and χ^2 tests, how misleading a high correlation can be. The technique of Bland and Altman is not appropriate here since it implicitly assumes that neither proportion has higher validity than the other and so would use the average as the best estimate of the true proportion. In this case, the family health services authority's register, whatever its shortcomings, must be regarded as the most accurate available source of information.

Lyons and colleagues express concern about the number of unmatched postcodes in our study. While this concern may be valid, one of our aims was to investigate how easy or difficult it was to match postcodes in registers of patients with their enumeration district using the postcode to enumeration district directory released with the 1991 census. The number of missing postcodes is an indication of the difficulty that exists. We were looking for a process that could be used straightforwardly and systematically nationally.

Both letters express concern at the use of sample surveys as an alternative way of obtaining data on practice populations. We are aware of the problems and envisaged instead a requirement to collect more details from patients at registration and the collection of more information from practices on a regular basis, in the way that data on chronic disease management are becoming available. The low income scheme index, described by Lloyd *et al*,⁴ is another example of data collected about the practice rather than estimated from another source.

Lyons and colleagues express concern about the transfer of resources from more deprived to less deprived areas. If they examine the figure in our paper they will find that this is exactly what the method that they defend achieves since practices at the extremes have predicted values that are closer to the mean than the true value.

We concede that the decision about whether the estimation of practice characteristics from census variables is "good enough" is ultimately subjective. We believe, however, that the discrepancies that we found with estimating such a gross characteristic as the proportion of elderly people are sufficient to cast grave doubt on the use of characteristics such as permanent sickness, which, because of small numbers, are much more difficult to estimate.

GLEN SCRIVENER

Social geographer

DAVID C LLOYD

Applied research statistician

University of Leeds,
Leeds LS2 9NZ

- 1 Allocating census data to general practice populations [letters]. *BMJ* 1995;311:875-6. (30 September.)
- 2 Scrivener G, Lloyd DCEF. Allocating census data to general practice populations: implications for study of prescribing variation at practice level. *BMJ* 1995;311:163-5. (15 July.)
- 3 Majeed FA, Cook DG, Polonicki J, Griffiths J, Stones C. Sociodemographic variables for general practices: use of census data. *BMJ* 1995;310:1373-4.
- 4 Lloyd DCEF, Harris CM, Clucas DW. Low income scheme index: a new deprivation scale based on prescribing in general practice. *BMJ* 1995;310:165-9.

Learning from primary care in developing countries

Innovations in developing countries have had worldwide relevance

EDITOR,—Paul Johnstone and Isobel McConnan point out that the health service in Britain can learn many lessons from health services in poorer countries,¹ and I agree. However, the push for community based rather than hospital services has been difficult both to initiate and to sustain. In the authors' example of Tanzania, although in the early 1970s the expenditure in hospitals may have been reduced to less than 20% of the health budget, this was difficult to maintain and the proportion has crept up again (G J Ebrahim, personal communication).

The health services in most developing countries are a mixture of traditional-indigenous, private-commercial at many levels, and government provided. Most British people who have worked in developing countries in recent years have worked with charitable non-government organisations, which have the greatest opportunity for innovation and flexibility.

Johnstone and McConnan refer to principles, and I would like to illustrate these with a few specific examples. It was in a non-government organisation in Nigeria that individual child growth and health records retained by the family were pioneered.² Now no reputable fundholding practice or child health trust in Britain would be without these records. It was in Bengal and Bangladesh that oral rehydration therapy for diarrhoea with dehydration was developed into a practical treatment.³ Now it is a basic recom-