Redesigning the journal: having your say

Tony Delamothe, Richard Smith

Our quarterly surveys suggest that a sizeable proportion of readers believe that the BMJ's layout now looks tired and out of date, and we think that the time has come to redesign it. Our last redesign (eight years ago) was rated a success, and we are keen to introduce only those changes that readers think are improvements.

So far we have sought readers' views on the journal's design by using a questionnaire survey and a series of focus groups. On the basis of the findings, we drew up an initial brief (box) and asked three designers (short-listed from 20) to suggest changes to the cover, a research paper, and a section that they thought would especially benefit from a redesign. (They all chose the news.)

A, B, or C?

On the following pages are their designs, labelled A, B, and C. They are one quarter actual size and do not represent a return to the days of miniprint. Even if you can read the words (and in one case they are in Latin) don't get distracted by what they say—on this occasion, we are interested in your opinion of the "look" of the pages. We care about content too, but improving the content is not the purpose of this redesign.

Remember that what you dislike about the designs is as important to us as what you like. Please address your comments to the editor. We will use them to help us choose the ultimate designer, who will be asked to take the comments into account.

As you will see, the greatest differences emerged in the suggestions for the cover—reflecting the contradiction set out in the design brief. This is that readers find the current cover both usefully comprehensive and too crowded. Readers should be reassured that if we opted for a cover that omitted a full listing of contents then this would appear elsewhere in the journal.

Ask the customer

This is not the first time we have consulted readers on how we should change. Readers may remember that this time last year we sought their advice regarding the journal's coverage of obituaries.¹ Responses to a questionnaire (which yielded a response rate of 60%) and more than a 100 letters to the editor² helped us to change this section, we think for the better.

Our Internet site provides for electronic feedback, and the changes that we have made³ and are planning to make to what we provide there have been at our readers' suggestion. We think that consulting our readers over the journal's redesign marks the next phase in our closer collaboration with them.

These designs, and the opportunity to record comments on line, are available on the BMJ's web pages at <http://www.bmj.com/bmj/>

- 1 Crossan L, Smith R. Growing pressure on BMJ's obituaries. *BMJ* 1995;310:5-6.
- 2 Correspondence. Obituaries. *BMJ* 1995;310:660-2.
- 3 Delamothe T. Hospital jobs on the Internet. *BMJ* 1995;311:966.

Brief for the redesign of the BMJ

Aims of the redesign

The current perception of the journal is that it looks dense, dull, and somewhat forbidding. The aim of the redesign is to liven up its appearance while maintaining its authority. A successful redesign would increase reading by current readers and would attract new readers to the journal. The redesign is explicitly about the appearance of the journal, not its content.

Specific requests

- Maintain A4 page size and maintain current order of articles
- Bear in mind the difficulty of photocopying articles from bound volumes
- Address problem of "too many words to the page" by altering type face, type size, leading, amount of white space, or combination of these
- Improve signposting
- Explore more use of colour, especially spot colour (eg cyan)
- Improve data presentation
- Cover:
 - Drop the snake
 - Differentiate four different editions

Because they make the journal instantly recognisable we would be reluctant to jettison both the BMJ logo with the distinctive dropped J and the cyan cover entirely (although there is room for manoeuvre)

How much of the content goes on the front cover (and how much goes elsewhere) and in what detail is open to negotiation. The current perception is that the cover is too crowded but usefully comprehensive. Somewhere in the journal all the details that appear on the front cover and first page will have to appear.

Background

The BMJ's mission is "to publish rigorous accessible information to help doctors improve their practice and to influence the international debate on health" and to do so cost effectively. It sees its functions as informing, educating, amusing, and entertaining its readership. The journal is meant to appeal to all sections of the medical profession, at all levels of experience—from newly qualified doctors to retired doctors. Four editions of the journal are published each week: Clinical Research, General Practice, International, Compact. They have identical editorial copy but differ in their advertisement content.

BMJ, London WC1H 9JR
 Tony Delamothe, deputy
 editor
 Richard Smith, editor

BMJ 1996;312:232-5

Clinical Research

British Medical Association
 Tavistock Square, London, WC1H 9EP
 Published weekly
 ISSN 0959-8134
 ASTM Code
 BMDJAE 511 (7000) 775-540 (1995)

BMJ

British Medical Journal
 No. 708 Volume 511
 Saturday 23 September 1995

Priorities and rationing: pragmatism or principles?

A self administered parental questionnaire sent to the parents on the day of examination in the 1991 survey was used to collect information on birth weight and gestational. Editorials p.761

Cholesterol and mortality: do men and women differ?

Subjects born in the two study areas, including birth weight and gestational age between maternal recall and birth record information, analyses of birth weight and blood pressure. Papers p.779

ABC of medical computing: computers in general practice

Hospital birth record information was collected on subjects born in the study areas, including birth weight. Education p.800

News

Royal college calls for national priorities setting

Jane Smith, BMJ

The Royal College of Physicians is calling for a national council to help determine priorities in the health service (see also p.761). In launching the college's report, *Setting Priorities in the NHS: A Framework for Decision Making*, the President, Sir Leslie Turnbull, emphasised that choices had to be made within the NHS since not everything that was possible was affordable.

The report argues that equity for the whole population is more important than freedom of individual choice but says that the criteria for setting priorities must be open. Sir Leslie said that there was confusion about who was taking decisions on priorities. "There need to be more open systems by which priorities can be made and much greater involvement by the public," he said.

This weeks report makes a strong push by the medical establishment for openness and explicitness in the setting

of the priorities. The proposed National Council for Health Care Priorities would review the methods used in determining priorities, monitoring how they are set and evaluating their implications. The council should, says the report, also aim to involve and educate the public, profession, and government. Sir Leslie emphasised that it would not be the council's job to determine what treat-

ments health authorities should and should not provide.

An example of the sorts of issues the council should consider was given at the press launch by Professor John Grimley Evans, professor of geriatric medicine in Oxford. Both patients and professionals were baffled, he said, by different experts expounding their own approaches to criteria for setting priorities. These varied from the quality adjusted life years of economists, through the epidemiologists' active life expectancy, to newer techniques that explore patient's personal values. "A national advisory council could bring these different approaches together and compare and contrast them," he said. Specialist units, with their reliance on complicated referral procedures, could well be threatened by current prioritising, says the report. It sees them as victims of priorities being set by cost considerations and argues that they need protection not only because of the care they offer but for research and development and training of health professionals. Two specific recommendations are that regional postgraduate deans should have a stronger role in coordinating educational priorities, to newer techniques that explore patient's personal values. The college aims to enlist broad support for its national council and is talking to the Department of Health and other organisations such as the Patients Association and the National.

Taxation alone cannot save the NHS, says report

Claudia Court, BMJ

The report argues that equity for the whole population is more important than freedom of individual choice but says that the criteria for setting priorities must be open. There was confusion about who was taking decisions on priorities. "There need to be more open systems by which priorities can be made and much greater involvement by the public," he said. This weeks report makes a strong push by the

medical establishment for openness and explicitness in the setting of the priorities. The proposed National Council for Health Care Priorities would review the methods used in determining priorities, monitoring how they are set and evaluating their implications. The council should, says the report, also aim to involve and educate the public, profession, and government. Sir Leslie empha-

Papers

Birth weight and blood pressure: cross sectional and longitudinal relations in childhood

Peter Whincup, Derek Cook, Olia Papacosta, M Walker

Abstract

Objective—To examine cross sectional and longitudinal relations between birth weight and blood pressure in childhood.

Design—Cross sectional study of primary school children aged 9-11 years, with analysis in relation to previous measurements at 5-7 years in a sub-group.

Setting—20 primary schools in Guilford and Carlisle. **Subjects**—1511 children measured at 9-11 years (response rate 79%), including 549 who had been measured at 5-7 years.

Main outcome measures—Blood pressure at 9-11 years, change in blood pressure between 5-7 and 9-11 years, birth weight (based on maternal recall), and placental weight (based on birth records).

Results—At 9-11 years birth weight was inversely related both to systolic blood pressure (regression coefficient -2.80 mm Hg/kg; 95% confidence interval -3.84 to -1.76) and to diastolic blood pressure (regression coefficient -1.42 mm Hg/kg; -2.14 to -0.70) once current height and body mass index were taken into account. Placental weight was inversely related to blood pressure after adjustment for current height and body mass index but placental ratio (placental weight to birth weight) was unrelated to blood pressure. Between 5-7 and 9-11 years systolic blood pressure rose more rapidly in children of lower birth weight (regression coefficient -1.17 mm Hg/kg; -3.35 to 0.070). This effect seemed to be stronger in girls.

Conclusions—Birth weight rather than placental ratio is the early life factor most importantly related to blood pressure in childhood. The results support the possibility of "amplification" of the relation between birth weight and blood pressure, particularly in girls.

University Department of Public Health
 Royal Free Hospital School of Medicine, London NW3 2PF
 Peter Whincup, senior lecturer in clinical epidemiology
 Olia Papacosta, research statistician
 M Walker, research administrator
 Department of Public Health Sciences
 St George's Hospital Medical School, London NW3 2PF
 Derek Cook, senior lecturer in epidemiology
 Correspondence to: Dr Whincup
 BMJ 1995; 311: 773-6

Introduction

Barker and colleagues argued that early life factors, acting particularly in utero, may have a strong influence on subsequent cardiovascular risk.¹ An inverse relation between birth weight and subsequent blood pressure (lower birth weight being associated with higher subsequent mean blood pressure) has been reported in adults²⁻⁴ and children,⁵⁻¹⁴ though not consistently in adolescents.¹⁵ Earlier reports also suggested that a large placenta, particularly in the presence of low birth weight, was related to subsequent high blood pressure and the risk of hypertension.¹⁶ Recent reports have suggested that the birth weight and blood pressure relation becomes stronger with age ("amplification")¹⁷ and that this may be an example of "programming".¹⁸ However, the evidence for amplification is based almost entirely on comparison of cross sectional studies with subjects of different ages and times of birth rather than on longitudinal measurements.¹⁹

We studied the relation between birth weight and blood pressure in children. In cross sectional analyses at 9-11 years of age we examined the relations between birth weight and blood pressure and the influence of placental weight. In longitudinal analyses based on children measured at both 5-7 and 9-11 years of age we examined the development of the birth weight and blood pressure relation with age.

Subjects and methods

We present data based on two surveys four years apart (1987 and 1991) in Guilford and Carlisle. The sampling and the methods used have been described elsewhere.^{20,21} Both surveys were carried out in the same sample of 10 primary schools in each town. The second survey included all children from the earlier survey still attending the schools together with similar numbers of children in the same classes. In both surveys measurements were made by a team of trained nurse observers, who visited schools in the two areas in sequence.

Blood pressure was measured after five minutes' rest with the child seated and the arm supported at chest level. Three blood pressure measurements were made one minute apart on the right arm with the Dinamap 18465C oscillometric blood pressure recorder (Critikon Inc, United States).²² All analyses were based on the means of the three readings. At 9-11 years all measurements were made with the adult size cuff (cuff bladder 22 cm x 12 cm); at 5-7 years all measurements

Papers

were standardised to the child size cuff (cuff bladder 15 cm x 9 cm), as described.²³ These cuff sizes ensured that the minimum cuff bladder width to arm circumference ratio of 40% recommended by the American Heart Association²⁴ was met for 98% of the study population at age 5-7 years. Ethnic group was assessed on the basis of the child's appearance.

A self administered parental questionnaire sent to the parents on the day of examination in the 1991 survey was used to collect information on birth weight and gestational age (assessed on the basis of recall of the number of weeks the birth was early or late). With the permission of mothers, hospital birth record information was collected on subjects born in the two study areas, including birth weight and gestational age between maternal recall and birth record information, analyses of birth weight and blood pressure at 9-11 years were based on maternal recall data.

A total of 924 children aged 5-7 years (84% response rate) took part in the 1987 survey; 1511 children aged 9-11 years (79% response rate) took part in the 1991 survey. All but eight children were of European origin; exclusion of the others had no effect on the results. Among the 1511 children measured in 1991, questionnaires with full birth details were returned for 1280 (81%). Of these, 996 children were born in the study towns; maternal consent to retrieval of birth records was obtained for 995 children. Complete birth records were obtained for 838 (84%) of these children, 473 of whom were Carlisle subjects with data on placental weight. The 1991 survey included 549 children of 699 (79%) who took part in the 1987 survey and were still attending the survey schools. The longitudinal relations between birth weight and blood pressure were examined in 523 of these 549 subjects for whom maternal recall information on birth weight was available.

Statistical analysis—All analyses were carried out with the statistical analysis software (SAS) package.²⁵ Birth weight, placental weight, age, and measures of current body size—height and body mass index (weight (kg) divided by height (m)²)—were either treated as continuous variables or ranked and grouped in equal thirds or fifths. Standard linear regression techniques were used throughout and all analyses standardised for the effect of the study area. Changes in rank by using

Fig 2. Systolic blood pressure centiles between 6 and 10 years of age. Centiles are based on data from 549 children studied longitudinally, measured at mean ages of 6.2 and 9.9 years and are standardised for sex, height, body mass index. (See text for options (a) and (b).)

standard deviation scores. These were calculated separately for blood pressure at 5-7 years and 9-11 years, standardised for age, sex, height, and body mass index at each age. Changes in standard deviation scores were then regressed on birth weight to examine the relation between birth weight and change in blood pressure rank between 5-7 years and 9-11 years.

Results

Birth weight and current body size (age 9-11)

Birth weight showed little or no relation with blood pressure in a model adjusted only for age and sex (fig 1). However, after additional adjustment for height and body mass index (correlation coefficients with birth weight -0.19 , -0.10 respectively) birth weight showed a smooth graded inverse relation with both systolic and diastolic pressure (fig 1). The corresponding regression coefficients defining the increase in blood pressure for a 1 kg fall in birth weight were -2.80 mm Hg (95% confidence interval -3.84 to -1.76 ; $P<0.0001$) for diastolic pressure. The relations between birth weight and blood pressure were very similar in preterm and term infants, in boys and girls, and in children in the top and bottom thirds of the distribution of height and body mass index. They were also independent of parity, social class, maternal blood pressure, and maternal smoking in pregnancy.

Birth weight and placental weight

Relations between birth weight, placental weight, and blood pressure were examined in 473 Carlisle children with data on these variables. Placental weight was strongly correlated with birth weight ($r=0.45$). Like birth weight, it showed a smooth inverse relation with blood pressure, particularly by systolic pressure, once current height and body mass index were taken into account. For a 100 g fall in placental weight systolic pressure rose by -0.84 mm Hg (95% confidence interval -1.50 to -0.18 ; $P=0.012$). However, including birth weight in the regression model greatly attenuated the relation between placental weight and blood pressure. Moreover, the placental ratio (placental weight to birth weight) showed no relation either with systolic pressure ($P=0.89$) or with diastolic pressure ($P=0.57$).

When the mean systolic blood pressure for each third of placental weight and birth weight was examined (table 1) the highest mean blood pressure was observed in the group in the lowest third of both birth weight and placental weight distributions; subjects with low birth weight and high placental weight did not have a particularly high mean blood pressure. Similar findings were recorded for diastolic pressure (data not shown). There was no evidence of interaction between birth weight and placental weight, either for systolic ($P=0.56$) or diastolic

Fig 1. Birth weight and blood pressure at 9-11 years, showing effect of adjustment for current body size. Mean blood pressure and 95% confidence intervals (bars) are shown for each fifth of birth weight. — Adjusted for age and sex only. — Adjusted for age, sex, height and body mass index.

BMJ

BRITISH MEDICAL JOURNAL
NUMBER 7008 VOLUME 311
SATURDAY 23 SEPTEMBER 1995

Editorial

761 Priorities and rationing: pragmatism or principles?

Rudolf Klein

764 Postoperative shivering: the influence of body temperature

Angus W A Crossley

766 Assessing the quality of care

How T O Davies, Iain K Crombie

Paper

773 Birth weight and blood pressure: cross sectional and longitudinal relations in children

Peter Whincup, Derek Cook, Ole Papacosta

783 Waiting list for dynamics and the impact of earmarked funding

John Newton, Jane Henderson, Michael J Goldacre

Education & Debate

780 Recent Advance: Interventional neuroradiology

Wendy Taylor, Georges Rodesch

796 A purchaser perspective of managing new drugs: interferon beta as a case study

T Wally, S Barton

NEWS

The NHS needs to go to its priorities right, says PCP

Royal College Calls For National Priorities Setting

Baraker and colleagues argued that early life factors, the acting of particularly in utero, may have strong influence on subsequent cardiovascular risk. An inverse relation between birth weight and the of subsequent blood pressure (lower birth weight being associated with higher subsequent mean it is in blood pressure) has been reported

consistently in adolescents. Earlier reports also suggested that a large placenta in between particularly in the presence of low birth weight, was related to subsequent a highest blood pressure and the risk of the hypertension. Recent reports have suggested that the birth weight and blood pressure relation becomes stronger with age (amplification) and that

this may be an good example of the "programming". However, all the evidence for amplification is based almost entirely on comparison of cross sectional studies with subjects of different ages and times of birth rather than on longitudinal measurements. We studied of the relation between the birth weight and each blood cell and pressure in children. In cross

sectional analyses at 9-11 years of age we examined the very relations between birth weight and blood pressure and the influence of placental weight. In longitudinal analyses based on children measured at both 5-7 and 9-11 years of age we examined the development of a the birth weight and blood pressure relation with age.

We present data based on two surveys four years of apart (1987 and 1991) in Guildford and Carlisle. The sampling and methods used have been detailed elsewhere. Both the surveys were carried out in the same sample of 10 primary schools in each town. The of second survey included all children from the earlier survey still attending of the schools together with similar numbers of children in the same classes. In both surveys measurements were made by a team of the trained nurse observers, who visited schools in the two areas in sequence. Blood pressure was measured after five halves minutes' rest with the child seated and the arm supported at chest level. Three blood pressure measurements were made one minute apart on the right arm with the Dinamap 1846SX oscillometric blood pressure recorder (Criticum Inc, United States). All analyses were based on the means of the three readings. At 9-11 years all measurements were made with the adult size cuff (cuff bladder 22 cm x 12 cm) at 5-7 years all measurements were standardised to the child

BMJ VOLUME 311 23 September 1995

769

PAPER

Birth weight and blood pressure: cross sectional and longitudinal relations in childhood

Peter Whincup, Derek Cook, Ole Papacosta, M Walker

ABSTRACT

Objective: To examine cross sectional and longitudinal relations between birth weight and blood pressure in childhood. **Design:** Cross sectional study of primary school children aged 9-11 years, with analysis in relation to previous measurements at 5-7 years in a subgroup. **Setting:** 20 primary schools in Guildford and Carlisle.

Subjects: 1511 children measured at 9-11 years (response rate 79%), including 549 who had been measured at 5-7 years.

Main outcome measures: Blood pressure at 9-11 years, change in blood pressure between 5-7 and 9-11 years, birth weight (based on maternal recall), and placental weight (based on birth records).

Results: At 9-11 years birth weight was inversely related both to systolic blood pressure (regression coefficient -2.80 mm Hg/kg; 95% confidence interval -3.84 to -1.76) and to diastolic blood pressure (regression coefficient -1.42 mm Hg/kg; -2.14 to -0.70) once current height and body mass index were taken into account. Placental weight was inversely related to blood pressure after adjustment for current height and body mass index but placental ratio (placental weight to birth weight) was unrelated to blood pressure. Between 5-7 and 9-11 years systolic blood pressure rose more rapidly in children of lower birth weight (regression coefficient -1.71 mm Hg/kg; -3.35 to -0.07). This effect seemed to be stronger in girls.

Conclusions: Birth weight rather than placental ratio is the early life factor most important related to blood pressure in childhood. The results support the possibility of "amplification" of the relation between birth weight and blood pressure, particularly in girls.

INTRODUCTION

Baraker and colleagues argued that early life factors, acting particularly in utero, may have a strong influence on subsequent cardiovascular risk. An inverse relation between birth weight and subsequent blood pressure

(lower birth weight being associated with higher subsequent mean blood pressure) has been reported in adults¹ and children^{2,3} though not consistently in adolescents.^{4,5} Earlier reports also suggested that a large placenta, particularly in the presence of low birth weight, was related to subsequent high blood pressure and the risk of hypertension.⁶ Recent reports have suggested that the birth weight and blood pressure relation becomes stronger with age (amplification)⁷ and that this may be an example of "programming".⁸ However, the evidence for amplification is based almost entirely on comparison of cross sectional studies with subjects of different ages and times of birth rather than on longitudinal measurements.⁹

We studied the relation between birth weight and blood pressure in children. In cross sectional analyses at 9-11 years of age we examined the relations between birth weight and blood pressure and the influence of placental weight. In longitudinal analyses based on children measured at both 5-7 and 9-11 years of age we examined the development of the birth weight and blood pressure relation with age.

SUBJECTS AND METHODS

We present data based on two surveys four years apart (1987 and 1991) in Guildford and Carlisle. The sampling and methods used have been detailed elsewhere.^{10,11} Both surveys were carried out in the same sample of 10 primary schools in each town. The second survey included all children from the earlier survey still attending the schools together with similar numbers of children in the same classes. In both surveys measurements were made by a team of trained nurse observers, who visited schools in the two areas in sequence.

Blood pressure was measured after five minutes' rest with the child seated and the arm supported at chest level. Three blood pressure measurements were made one minute apart on the right arm with the Dinamap 1846SX oscillometric blood pressure recorder (Criticum Inc, United States).¹² All analyses were based on the means of the three readings. At 9-11 years all measurements were made with the adult size cuff (cuff bladder 22 cm x 12 cm) at

5-7 years all measurements were standardised to the child size cuff (cuff bladder 15 cm x 9 cm), as described.¹³ These cuffsizes ensured that the minimum cuff bladder width to arm circumference ratio of 40% recommended by the American Heart Association was met for 98% of the study population at age 9-11 years and for 99% of the population at age 5-7 years. Ethnic group was assessed on the basis of the child's appearance.

A self administered parental questionnaire sent to the parents on the day of examination in the 1991 survey was used to collect information on birth weight and gestational age (assessed on the basis of recall of the number of weeks the birth was early or late). With the permission of mothers, hospital birth record information was collected on subjects born in the two study areas, including birth weight, gestational age based on date of last menstrual period, and (available in Carlisle only) placental weight. Because there was close agreement in birth weight and gestational age between maternal recall and birth record information, analyses of birth weight and blood pressure at 9-11 years were based on maternal recall data.

A total of 924 children aged 5-7 years (84% response rate) took part in the 1987 survey, 1511 children aged 9-11 years (79% response rate) took part in the 1991 survey. All but eight children were of European origin; exclusion of the others had no effect on the results. Among the 1511 children measured in 1991, questionnaires with full birth details were returned for 1230 (81%). Of these, 996 children were born in the study towns; maternal consent to retrieval of birth records was obtained for 995 children. Complete birth records were obtained for 838 (84%) of these children, 473 of whom were Carlisle subjects with data on placental weight. The 1991 survey included 549 children of 699 (79%)

who took part in the 1987 survey and were still attending the survey schools. The longitudinal relations between birth weight and blood pressure were examined in 523 of these 549 subjects for whom maternal recall information on birth weight was available.

Statistical analysis: All analyses were carried out with the statistical analysis software (SAS)¹⁴ packaged. Birth weight, placental weight, age, and measures of current body size—height and body mass index (weight (kg) divided by height (m)²)—were either treated as continuous variables or ranked and grouped in equal thirds or fifths. Standard linear regression techniques were used throughout and all analyses standardised for the effect of study area. Changes in blood pressure between 5-7 and 9-11 years were examined both as absolute changes and as changes in rank by using standard deviation scores. These were calculated separately for blood pressure at 5-7 years and 9-11 years, standardised for age, sex, height, and body mass index at each age. Changes in standard deviation scores were then regressed on birth weight to examine the relation between birth weight and change in blood pressure rank between 5-7 and 9-11 years.

RESULTS

BIRTH WEIGHT AND CURRENT BODY SIZE (AGE 9-11)

Birth weight showed little or no relation with blood pressure in a model adjusted only for age and sex (fig 1). However, after additional adjustment for height and body mass index (correlation coefficients with birth weight $r=0.19$, $r=0.10$ respectively) birth weight showed a smooth graded inverse relation with both systolic and diastolic pressure (fig 1). The corresponding regression coefficients defining the increase in blood pressure for a 1 kg fall in birth weight were -2.80 mm Hg (95% confidence interval -3.84 to -1.76, $P<0.0001$) for systolic pressure and -1.42 mm Hg (-2.14 to 0.70; $P<0.0001$) for diastolic pressure. The relations between birth weight and blood pressure were very similar in preterm and term infants, in boys and girls, and in children in the top and bottom thirds of the distribution of height and body mass index. They were also independent of parity, social class, maternal blood pressure, and maternal smoking in pregnancy.

1 Baker DJR, et al. The fetal and infant origins of adult disease. London: BMJ Publishing Group, 1992.

2 Baker DJR, Osmond C, Golding J, Kell SJ, Madsen H, et al. Growth in utero, blood pressure, childhood and adult life, and mortality from cardiovascular disease. BMJ 1990;301:254-7.

3 Baker DJR, Osmond C, Golding J, Osmond S. Fetal and placental size and risk of hypertension in adult life. BMJ 1990;301:254-7.

4 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

5 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

6 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

7 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

8 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

9 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

10 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

11 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

12 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

13 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

14 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

15 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

16 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

17 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

18 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

19 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

20 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

21 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

22 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

23 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

24 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

25 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

26 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

27 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

28 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

29 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

30 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

31 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

32 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

33 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

34 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

35 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

36 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

37 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

38 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

39 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

40 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

41 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

42 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

43 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

44 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

45 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

46 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

47 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

48 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

49 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

50 Lissner G, Björk M, Lapid L. Low birth weight and risk of high blood pressure in adulthood. BMJ 1990;301:254-7.

BMJ VOLUME 311 23 September 1995

773

773

BMJ VOLUME 311 23 September 1995

BRITISH MEDICAL JOURNAL

Editorials

- 763 Angiotensin converting enzyme genotypes and disease
RASHA N. TAYLOR
- 764 Postoperative shivering: the influence of body temperature
ANGELA M. A. CROOK
- 765 Managing cleft lip and palate
TONY MARSH, PETER WARD-BENNETT

Commentary

- 761 Priorities and rationing: pragmatism or principles?
RASHA N. TAYLOR
- 762 Emergency contraception
JANEY C. LAYTON
- 766 Assessing the quality of care
HOW T. O. DAVIES, JANE R. GARDNER

Papers

- 773 Birth weight and blood pressure: cross sectional and longitudinal relations in childhood
HOW T. O. DAVIES, PETER WARD-BENNETT
- 776 FcγR2 polymorphism and risk of atopy in a general population sample
M. A. HALL, A. S. JAMES, J. A. FINE, G. RANK, J. M. THOMPSON, P. A. STOKES, W. S. M. COCHRAN
- 779 Total cholesterol concentration and mortality at a relatively young age: do men and women differ?
JANEY C. LAYTON, HOW T. O. DAVIES
- 783 Waiting list dynamics and the impact of earmarked funding
JOHN N. NEWBURN, JANE HENDERSON, MICHAEL J. COLEMAN
- 785 Acquired immunodeficiency without HIV infection: epidemiology and clinical outcome in Italy
C. RINALDI, P. PROSPERINI, F. ACTIS DONATO, THE ITALIAN STUDY GROUP ON NON-HIV AIDS
- 789 Recent Advances: Interventional neuroradiology
WENDY TAYLOR, GORDON BOWLER
- 793 Detecting differences in quality of care: the sensitivity of measures of process and outcome in treating acute myocardial infarction
JANEY C. LAYTON, HOW T. O. DAVIES

Features

- 787 Analysis from inner London of deprivation payments based on enumeration districts rather than wards
TIM CRABTREE, JOHN SHANKS, MADHURI RAJAGAL, STEPHEN LANGRISH
- 796 A purchasers perspective of managing new drugs: interferon beta as a case study
T. WALSH, J. BARNES
- 800 ABC of Medical Computing: Computers in general practice
ANDREW WILLIAMS, NICHOLAS LEE, ANDREW BROWN

BRITISH MEDICAL ASSOCIATION • BRITISH MEDICAL JOURNAL • WILEY • ISSN 0954-6820 • A51M COUNCIL: 88/004/95 (C) 1995 BMJ. 0954-6820/95

POLITICS

Royal college calls for national priorities setting

O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

Gallia est okmnis diuisa in partes tres, quarum unam incolent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appellantur. Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

VOLUME 312 23 SEPTEMBER 1995

De sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni.

intermittam. Eterna vobisque tan tum ergo semper fecit et hotum toyum eusque persuaderet. Quis post mortem et vitam suis et eis volent videre. Gallia est okmnis diuisa in partes tres, quarum unam incolent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appellantur. Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

VOLUME 312 23 SEPTEMBER 1995

rem fedes quod uocare dixisse eu ego dubite fumo iodio si cum sedat, necuo comodo me conuenire possit. Atqui si quisquam et si facili hic est, tamen non dubito quin me male oderit. Haec et eiusmodi multa. Sed ad propositum quicquid erit non modo magnum sed etiam pa rum scribes equidem nihil in ter mittam. Eterna vobisque tan tum ergo semper fecit et ho tum toyum eusque per vos detur. Quis post mortem et vit am suis et eis volent videre.

Gallia est okmnis diuisa in partes tres, quarum unam in colent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appel lan tur. Hi omnes lingua instituit legibus inter de different. Gal los ab Aquitanis Garunna flu men, a Belgis Ma trona et Seq uana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi volentem videre. Gallia est okmnis diuisa in partes tres, quarum unam incolent Belgae. ■

Global changes in climate will affect health

O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

W allia est okmnis diuisa in partes tres, quarum unam incolent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appellantur. Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

VOLUME 312 23 SEPTEMBER 1995

PAPERS

niam illum desiderat, namo sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

EXPECTAREM FEDES QUOD

Ruocare dixisse eu ego dubite fumo iodio si cum sedat, necuo comodo me conuenire possit. Atqui si quisquam et si facili hic est, tamen non dubito quin me male oderit. Haec et eiusmodi multa. Sed ad propositum quicquid erit non modo magnum sed etiam pa rum scribes equidem nihil in termittam. Eterna vobisque tantum ergo semper fecit et hotum toyum eusque persuaderet. Quis post mortem et vitam suis et eis volent videre. Gallia est okmnis diuisa in partes tres, quarum unam incolent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appellantur. Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

De sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

VOLUME 312 23 SEPTEMBER 1995

Birth weight and blood pressure: cross sectional and longitudinal relations in childhood

PETER WARD-BENNETT, DEBRA COOK, OLGA PAPADIMITRI, M. WALKER

ABSTRACT

OBJECTIVE Atqui si quisquam et si facili hic est, tamen non dubito quin me male oderit. Haec et eiusmodi multa. Sed ad propositum quicquid erit non modo magnum sed etiam pa rum scribes equidem nihil in termittam. Eterna vobisque tantum ergo semper fecit et hotum toyum eusque persuaderet. Quis post mortem et vitam suis et eis volent videre.

DESIGN Atqui si quisquam et Gallia est okmnis diuisa in partes tres, quarum unam incolent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appellantur. Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

SETTING Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

MAIN OUTCOME MEASURES Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

RESULTS Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

CONCLUSION Atqui si quisquam et Gallia est okmnis diuisa in partes tres, quarum unam incolent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appellantur. Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

Gallia est okmnis diuisa in partes tres, quarum unam incolent Belgae, aliam Aquitani, tertiam, quis iporum lingua Celtae, nostra galli appellantur. Hi omnes lingua instituit legibus inter de different. Gallos ab Aquitanis Garunna flumen, a Belgis Matrona et Sequana diuidit. Reverti ad illu: de quo tecu mane nihil ditius explicare te non posse. Enim si ille tibi tali ingenio exitu non reperiebat. Muc reperi? Quid quaeris Perisse omnia aiebat: quod aut scio an ita sit uerim ille gaudens et scicabre. O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

O prudente oppii qui nihilinimus il lum desiderat, sed loquitor nihil queng bonum offendat. Sed haec hactenus. Tu quaro quicquid noui multa autem ex pecto scribere, ne pigre in his. De km bog sexto fati ne certum, maxime au te de Bruto nostro de quo quidem illo, ad quem diueri solitum dicere magni refert hic quid uelit, sed quicquid ne au te de Bruto nostro de quo quidem pigre in his. De km bog sexto fati ne certum quicquid noui.

VOLUME 312 23 SEPTEMBER 1995

VOLUME 312 23 SEPTEMBER 1995