a substantial proportion of women who lost their virginity under the age of 16 consider in retrospect that this was too young.^{4,15} Little is known about the effect of early sexual intercourse on subsequent psychosexual functioning. The issue of whether a young person is ready to embark on a sexual relationship remains a complex one. The scale of morbidity currently associated with under age sex is sobering and suggests that, for many teenagers, sexual activity is far from appropriate.

SUE STUART-SMITH

Registrar in child and adolescent psychiatry

Child Mental Health,
High Wycombe,
Bucks HP13 6PQ

¹ Department of Health. *Family planning clinic services 1993-4. Summary information from form KT31* England. London: DoH, 1995.

- 2 Knight B. *Legal aspects of medical practice*. Edinburgh: Churchill Livingstone, 1982.
- 3 Loudon N. *Handbook of family planning*. Edinburgh: Churchill Livingstone, 1991.
- 4 Wellings K, Field J, Johnson AM, Wadsworth J. *Sexual behaviour in Britain*. London: Penguin, 1994.
- 5 Farrow JA, ed. *The medical clinics of North America: adolescent medicine*. Vol 74 (No 5). Philadelphia: W B Saunders, 1990.
- 6 Mellanby A, Phelps F, Tripp JH. Teenagers, sex, and risk taking. *BMJ* 1993;307:25.
- 7 Durbin M, DiClemente RJ, Siegel D, Krasnovsky F, Lazarus N, Camacho T. Factors associated with multiple sex partners among junior high school students. *Journal of Adolescent Health Care* 1993;14:202-7.
- 8 Zabin LS, Hardy JB, Smith EA, Hirsch MB. Substance abuse and its relation to sexual activity among inner city adolescents. *Journal of Adolescent Health Care* 1986;7:320-31.
- 9 Zabin LS. The association between smoking and sexual behaviour among teens in US contraceptive clinics. *Am J Public Health* 1984;74:261-3.
- 10 Gordon DE. Formal operational thinking: the role of cognitive developmental processes in adolescent decision making about pregnancy and contraception. *Am J Orthopsychiatry* 1990;60:346-56.
- 11 Anastasiow NJ. *The adolescent parent*. London: Brookes, 1982.
- 12 Adcock AG, Nagy S, Simpson JA. Selected risk factors in adolescent suicide attempts. *Adolescence* 1991;26:817-28.
- 13 Denman S, Gillies P, Wilson S, Wijewardene K. Sex education in schools: an overview with recommendations. *Public Health* 1994;108:251-6.
- 14 Essex B. *Doctors dilemmas decisions*. London: BMJ Publishing, 1994:113.
- 15 Curtis HA, Lawrence CJ, Tripp JH. Teenage sexual intercourse and pregnancy. *Arch Dis Child* 1988;63:373-9.

Being creative about rationing

An innovatory theory helps clearer thinking

Two of the biggest questions in health care are "How much money should we spend?" and "What should we spend it on?" We don't have adequate methods for answering either question, but Ronald Dworkin, professor of law in both Oxford and New York and one of the world's leading thinkers on medical ethics, has developed a theory that provides a means to begin to answer both questions. Last week he presented his "prudent insurance principle" at a meeting in London organised by the Institute for Public Policy Research.

Current practice with health care expenditure is based—more implicitly than explicitly—on what Professor Dworkin calls "the insulation model." This postulates that health care is fundamentally different from other goods, that equality of access is essential for health care because it is so different, and that when something can be done acutely to avoid death it should be done ("the rescue principle"). In the United States this principle has led to such inappropriate actions as huge sums being spent to separate Siamese twins joined at the heart even though one will inevitably die and the other has only a tiny chance of surviving and no chance of living a normal life. In Britain large sums of money have been spent to give the teenager Jaymee Bowen further treatment for her leukaemia even though the discomfort to her is high, the costs substantial, and the chances of her long term, symptom free survival small. The model is, said Professor Dworkin, neither "sensible nor sane": no society would spend all its resources on health care rather than on education, housing, or job creation; and nor would any individual. Anybody who devoted all of his or her resources to avoiding any chance of death would live a grossly diminished life.

Professor Dworkin also dismisses the idea that increased efficiency can avoid the problem of having to make hard choices about what can be afforded. When he examines this model he discovers that it necessitates excluding "low yield" procedures like breast screening in women under 50. But there is all the difference between "low yield" and "no yield" procedures, and people who advocate this model are really accepting the need for rationing. Resources might be allocated, some say, "on the basis of need with an eye to cost effectiveness." Professor Dworkin argues that this is a benign but unhelpful theory: it does not help answer the questions in practical terms. Because no ready theory is to hand and because the questions depend on deep cultural values where

nobody is more expert than anybody else another argument is to leave it all to politics and politicians. But this will lead to decisions based on what will win votes rather than on what is rational.

Something better is needed. Professor Dworkin asked his audience to imagine a world with five characteristics: wealth is justly distributed, meaning for Professor Dworkin that people have roughly the same amount; information is available to all on all aspects of medicine and its effectiveness; people make decisions rationally; parents put their children's interests on the same level as their own; and nobody knows anything about genetic, cultural, or social predispositions to disease. In this imaginary world the government would abstain from providing health care, and people would be left to make their own decisions on which insurance policies to buy. Importantly, people would be obliged to decide what health problems to insure against and what intensity of treatment they should be covered for. Professor Dworkin claims that in such a world the amount and distribution of spending on health care would be just. People would decide rationally on how much to spend on health care as opposed to, say, education or housing, and people would make their own decisions on whether to buy a policy that gave them access to heart surgery until they were 75 years old or a much more expensive policy that gave them access until they were 90.

Doctors, who are practical people distrustful of theories, may not see any possible use for Professor Dworkin's theory. But economists, whose whole discipline is built on theories with equally fantastic assumptions, may be interested. Professor Dworkin argues that the theory can be useful for many decisions. Few people, he says, would buy a health insurance policy that offered them life saving treatment if they fell into a persistent vegetative state, and yet about 10 000 people in the United States are currently kept alive in such a state. Nor does he think there would be many takers for a policy that offered life saving treatment in the late stages of dementia. And, if they were faced with choosing between a policy that would deny them life saving treatment when they were known to be within four months of death and another much more expensive policy that offered all possible treatments, Professor Dworkin thinks that most people would opt for the first. In fact 40% of Medicaid expenditure is on people

in the last four months of their lives (not all of whom, of course, are expected to be dead within the four months).

Professor Dworkin thinks that his model also makes it clear that age would be a factor in most people's decision making. A policy that denied life saving treatment over the age of, say, 85 would surely be preferred over a much more expensive one that allowed it—partly because people might not reach such an age and more because they would have to forego substantial benefits during their lives in order to pay for the policy. And even if they paid for a policy that allowed treatment to an older age there would be some point at which they desisted.

At the other end of the range, most people would buy policies that paid for mending a broken leg or treating children with meningitis. There would probably be three categories of provision: some that almost everybody agreed were necessary; some that almost everybody agreed were unnecessary; and some in the middle, where people would choose differently. In the United States this might mean that people choose between different packages (as happens

increasingly with what is offered by American employers), or in Britain it might mean that the state provides a basic package and then people make private provision thereafter.

Professor Dworkin suggested three ways in which his model might be used in the real world: for "talking to people" through debates in the media, focus groups, and the like; for exploring insurance markets "unsoiled" by inequalities in wealth—like negotiations between large employers and unions over packages of health care; and for political mechanisms like that employed in Oregon. Those attending last week's seminar quizzed Professor Dworkin on his model, but nobody dissented from the notion that the idea provided a rich way to explore the issue of rationing. And we have few, if any, other such creative aids to thought with this most difficult of questions.

RICHARD SMITH
Editor

BMJ,
London WC1H 9JR

Adjuvant treatment for colorectal cancer

No more room for nihilism

It is now over five years since the publication of two randomised trials showing that fluorouracil and levamisole reduced the relapse rate by 40% and mortality by a third in patients with node positive colorectal cancer.^{1,2} The final report of these data was recently published, showing 168 deaths in control patients versus only 121 in patients who received adjuvant treatment ($P < 0.005$).³ Despite these data, a recent survey of oncologists and surgeons in Britain showed that most oncologists offered adjuvant chemotherapy to appropriate patients while many surgeons remained sceptical (R Grey, personal communication).

The combination of fluorouracil and folinic acid has proved superior to fluorouracil alone in patients with advanced colorectal cancer.⁴ The Canadian and European consortium trial (IMPACT) compared adjuvant treatment with high dose fluorouracil and folinic acid with no treatment in nearly 1500 patients; it showed a 22% reduction in mortality, predominantly in patients with Dukes's type C colon cancer.⁵ A similar but much smaller Italian study showed a 39% reduction in mortality in patients with Dukes's type C cancer.⁶ An ongoing intergroup study showed a significant difference in recurrence free and overall survival of patients given fluorouracil with low dose folinic acid compared with patients receiving no adjuvant treatment at three years⁷ and five years (M J O'Connell, personal communication).

America's National Surgical Adjuvant Breast And Bowel Project compared outcome in patients given fluorouracil and high dose folinic acid with that in control patients given semustine, vincristine, and fluorouracil (MOF),⁸ a regimen that had previously been shown to offer a survival advantage in colorectal cancer.⁹ After three years, 23% of the trial patients had died compared with 32% of the controls.

The National Surgical Adjuvant Breast and Bowel Project recently conducted a study (protocol C-04) comparing fluorouracil and folinic acid with fluorouracil and levamisole and with the combination of all three. The group receiving fluorouracil and folinic acid had a borderline significant advantage in terms of survival and disease free survival compared with the group receiving fluorouracil and levamisole (N Wolmark, personal communication).

Another approach has been to use portal vein infusions of fluorouracil immediately after surgery. A meta-analysis of nine randomised trials showed a 13% reduction in risk of death for patients with Dukes's type B or C cancer.¹⁰ The largest trial of portal vein infusion was conducted by the National Surgical Adjuvant Breast And Bowel Project.¹¹ An updated analysis at five years showed a reduction in the incidence of liver metastasis and a reduction in death at five years from 29% in the control arm to 23% in the treatment arm ($P = 0.03$) (N Wolmark, personal communication).

Trials in patients with rectal cancer have found similar effects. Two small randomised trials found a significant survival advantage from fluorouracil and semustine in patients with rectal cancer compared with control patients receiving no chemotherapy.^{1 2 13}

Chemotherapy with fluorouracil and levamisole or fluorouracil and folinic acid is not without toxicity, but this usually consists only of mild nausea, mucositis, diarrhoea, and lethargy. Most of these toxicities are mild and can be controlled with modern antiemetic and antidiarrhoeal drugs.

Adjuvant chemotherapy has been established as part of standard treatment for most patients with Dukes's type C colorectal cancer, but its role in patients with Dukes's type B tumours remains uncertain; most trials have been too small to detect a survival advantage as the number of deaths has been much lower. The appropriate treatment for a rising carcino-embryonic antigen after surgery is unknown, as is the role of adjuvant therapy after hemihepatectomy for liver metastasis.

There are now two major ongoing trials in Britain that are designed to answer several important questions, both under the aegis of the United Kingdom Coordinating Committee for Cancer Research. The QUASAR trial looks at the role of fluorouracil and folinic acid, in both high and low doses with or without levamisole, and the AXIS trial should provide the definitive answer on the role of portal vein infusion. These trials have been designed in such a manner that there can be few patients who would not be eligible for one of them.

A nihilistic attitude to the treatment of colorectal cancer is no longer appropriate. We can debate the size of the survival benefit from adjuvant chemotherapy and the balance of