will be made available to vendors of general practitioner and hospital software for incorporation in their systems, whether on DOS or other platforms.

Copies of the *BNF* are purchased by the Department of Health in bulk and then provided free to doctors and pharmacists working for the NHS. No such arrangement exists at present for the electronic version, which is available only through the publishers (BMA and the Royal Pharmaceutical Society of Great Britain).

The stand alone *eBNF* runs adequately on personal computers with a 386 processor, though a more powerful processor and more memory will increase its speed. The stand alone *eBNF* is run from the hard disk. Most new non-clinical software that clinicians may want to use runs under Windows; users are therefore likely to replace their systems with newer hardware that will run such software.

STELLA DUTTON
Former director of publications

Royal Pharmaceutical Society of Great Britain,
London SE1 7JN

*Stella Dutton is now executive director of the BMJ Publishing Group—EDITOR

Program calculates doses for neonates

EDITOR,—The value of computer based prescribing¹ should be immense, and the potential value is even greater for babies than adults. Drug injections for adults often use a half to a whole standard ampoule, but sick or premature neonates frequently require only a tiny proportion of the phial, which increases the scope for accidental and possibly lethal overdose² caused by medical or nursing error.³ A sequence of calculations may be required to obtain the correct dose: the weight of the drug in milligrams or micrograms must be chosen according to the baby's weight and then adjusted according to factors such as age, gestation, or renal function. The displacement volume of dissolved powder or crystals may be relevant when small proportions of a phial are required.⁴ Once the weight of the dose is known, the correct volume of the solution must be chosen and the dose diluted to obtain a manageable volume, bearing in mind that a 0.02 ml injection cannot be practically given and that in multiple drug treatment minimal volumes must be used to avoid fluid overdose. Each of these stages adds scope for error, however accurate or user friendly the compendium or formulary, in spite of the use of an electronic calculator.

At the 1995 annual scientific meeting of the British Association for Perinatal Medicine I presented a computer program (BABYDRUG) that carries out all these stages automatically. It now incorporates 31 drugs and runs off Microsoft Windows (version 3.1 or later). The program calls up relevant personal and medical factors and gives screen and printer output of the dosage as well as the preparation of the final injection to the above criteria. The formulation can be exactly and easily tailored for the individual baby in a way that would be cumbersome or oversimplified by traditional calculation. Manufacturers' instructions are used where possible, and extra drugs will be added and improvements to the schedules made by further consultation with potential users. A database for presenting qualitative information about drugs and saving prescriptions for audit is being developed. Transfer at present is by a single floppy disk, and this is not a commercial venture.

J M DAVIES
Consultant paediatrician

Special Care Baby Unit,
Grimsby Maternity Hospital,
Grimsby, South Yorkshire DN3 1NW

1 Wyatt J, Walton R. Computer based prescribing. *BMJ* 1995;311:1181-2. (4 November.)

2 Ferner RE, Whittington RM. Coroner's cases of death due to errors in prescribing or giving medicines or due to adverse drug reactions. *J R Soc Med* 1994;87:145-8.

3 Rolfe S, Harper NJN. Ability of hospital doctors to calculate safe doses. *BMJ* 1995;310:1173-4.

4 Mulholland P. Displacement volumes of powder injections. *Pharmaceutical Journal* 1993;251:14-5.

Guidelines for GPs are being tested

EDITOR,—December saw the launch of the electronic version of the *British National Formulary* with its full text indications, side effects, etc.¹ January saw the start of a unique national research and development programme of computerised prescribing guidelines which will be synergistic to the electronic *BNF*. PRODIGY (prescribing rationally with decision support in general practice study) is the first programme of its kind in the United Kingdom to test the concept of computerised prescribing guidelines in a live environment. PRODIGY will provide computerised advice to general practitioners during the consultation on the different drug choices and advice leaflets that can be offered to patients.

PRODIGY has been developed from a Dutch program of computerised guidelines (PRESCRIPTOR),² collated and referenced by staff at the Medical Advisor Support Centre in Liverpool. The English guidelines have been reviewed by an expert panel with representatives of several professional bodies (General Medical Services Committee, Royal College of General Practitioners, Royal College of Physicians, and Royal Pharmaceutical Society of Great Britain), who have made over 400 amendments. The guidelines will be further developed within PRODIGY from suggestions from focus groups of general practitioners selected from the 150 volunteer practices participating and from other observational data from the study. Towards the end of phase 1 of PRODIGY, specifications for future continual development and change control methods will be drawn up from the experiences and views of users, thereby ensuring that the system developed is professionally driven.

The software implementation of these guidelines has been developed in the Sowerby Unit from an initial user requirement developed and published by the NHS Executive and will undergo an iterative development during several phases of PRODIGY. The development of this specification is exciting in that for the first time at a national level the user requirements for the computer system are being developed using qualitative and quantitative methods. The software requirement is being fine tuned by a collaboration of the five major suppliers of general practice software in England and the Sowerby Unit. If the project proves successful consideration will be given to including the software requirement specification in the requirement for accreditation³ for general practice computer systems.

The overall aim of phase 1 of PRODIGY is to test the concept and the acceptability of prescribing decision support for general practitioners.

IAN PURVES
Director

Sowerby Unit for Primary Care Informatics,
Department of Primary Health Care,
University of Newcastle,
Newcastle upon Tyne NE2 4AA

1 Wyatt J, Walton R. Computer based prescribing. *BMJ* 1995;311:1181-2. (4 November.)

2 Ter Wee RJM, van der Kleijn E, Breninkmeijer RF, Holmberg N. Development of an electronic prescription processing option; an aid for general practice. *Br J Gen Pract* 1991;41:151-4.

3 Information Management Group. *General medical practice computer systems: requirements for accreditation. Version 3* 1995/96. Leeds: NHS Executive, 1995.

Database is linked to Read codes

EDITOR,—Jeremy Wyatt and Robert Walton say that the compilers of the *British National Formulary* have begun the task of restructuring

their data for use as a drugs database for computers.¹ I sympathise because it is an enormous and quite daunting task to turn a book into a database.

The team that tackled the task of putting together knowledge about drugs for our World Standard Drug Database had the great advantage of being able to structure it for use on computers right from the start. Besides me (a general practitioner who spent six years working with James Read on the Read codes) the team consisted of Dr Ivan Stockley (a pharmacologist with special interest in drug interactions), Dr David Beckles (a mathematician and systems analyst), and Mr Colin Duncton (formerly a hospital chief pharmacist).

Our database is purpose built and written for integration with or to "bolt on" to existing Read coded clinical systems in hospital or general practice so that, when prescribing, users have full drug information at their fingertips. This information includes drug ingredient based indications; contraindications; side effects; and drug use in pregnancy, lactation, liver disease, and renal disease. Each product is linked to dosage, legal information, and other product specific information.

The database is pitched at a level of information comparable with the *British National Formulary* and the *Datasheet Compendium* so that it is crisp, to the point, and very user friendly. Read code terms are used throughout the 30 000 products, which have been coded for ingredients and strength. Our thesaurus, which contains all words, terms, and messages in the prescribing knowledge domain, is fully coded and classified. It includes dosage regimens; defined daily dosages; drug form terms (tablet, suppository, etc); Read codes (for indications, contraindications, and side effects); all drug ingredients (world list); pharmacological actions; interaction messages; and messages relating to use of drug ingredients in pregnancy, lactation, renal failure, and liver disease. It also has tables to show the cheapest product for any ingredient(s) and alternative products with similar pharmacological actions.

The drug interaction alerting module targets specific individually named drug ingredient pairs, thereby avoiding all the "noise" and false interactions generated when drugs are categorised into groups. Each interaction message that is "flagged" says what is likely to happen and offers a recommendation on how to handle it.

We have produced software called SafeScript-Interactive (available from Doctors Independent Network Coding Ltd at the address below) to demonstrate the World Standard Drug Database. We believe that the database will help to reduce errors in prescribing by flagging timely information and advice to clinicians during prescribing. As the initiator and editor I have been able to ensure that the database is tailor made for prescribers like myself. I use it every day in my surgery when prescribing, and I can't imagine how I used to manage without it.

ROGER WEEKS
General practitioner

2 Deanhill Road,
London SW14 7DF

1 Wyatt J, Walton R. Computer based prescribing. *BMJ* 1995;311:1181-2. (4 November.)

Patient's consent to abortion is usually necessary even in emergencies

EDITOR,—In Clare Dyer's report on a gynaecologist who was acquitted of performing an illegal abortion after he carried out a hysterectomy while suspecting that the patient was pregnant she states that "the emergency provisions of the 1967 Abortion Act . . . allow a doctor to terminate a pregnancy without the patient's consent if her physical or mental health is thought to be endangered by the