Published in final edited form as:

Tetrahedron. 2008 March 10; 64(11): 2537–2552.

Pd(OAc)₂-catalyzed domino reactions of 1,2-dihaloarenes and 2-haloaryl arenesulfonates with Grignard reagents: efficient synthesis of substituted fluorenes

Cheng-Guo Dong and Qiao-Sheng Hu *

Department of Chemistry, College of Staten Island and the Graduate Center of the City University of New York, Staten Island, New York 10314

Abstract

Pd(OAc)₂-catalyzed domino reactions of 1,2-dihalobenzenes and 2-haloaryl arenesulfonates with hindered Grignard reagents to form substituted fluorenes, which are believed to occur through palladium associated aryne intermediates, is described. Such palladium associated aryne reaction pathway was found to be favored by omitting the use of phosphine and *N*-heterocyclic carbene ligands for palladium catalysts and with better leaving groups. Our study suggested that Pd(leaving group) X associated arynes should be formed first and the *sp*³ C-H activation preferentially occurred at benzylic C-(1°)H bonds. The work described here provides a high yield, one-step access to substituted fluorenes from readily available 1,2-dihalobenzenes and 2-haloaryl arenesulfonates and hindered Grignard reagents, and this substituted fluorene-making method may find applications in the preparation of substituted fluorene-containing molecules including polymers.

Keywords

1,2-dihalobenzenes; domino reactions; Grignard reagents; 2-haloaryl arenesulfonates; palladium; substituted fluorenes

Introduction

Over the past decades, palladium-catalyzed cross-coupling reactions, e. g., the Suzuki coupling, the Kumada coupling, the Stille coupling, the Sonogashira coupling, and the amination of aryl halides with amines, etc., have become powerful tools for organic synthesis.¹ They have been extensively employed to synthesize a wide variety of organic compounds ranging from small organic molecules to macromolecules.^{1,2} Extensive study established that there are three key elementary steps in the catalytic cycles of Pd(0)-catalyzed cross-coupling reactions:¹ oxidative addition of Pd(0) with an aryl halide to form Pd(II) complex; transmetalation of the Pd(II) complex with an organometallic reagent to form an diorganopalladium(II) complex; and reductive elimination of the diorganopalladium(II) complex to form the cross-coupling product and regenerate the Pd(0) catalyst (Figure 1). Based on our understanding of the general mechanism depicted in Figure 1, we envisioned that the individual elementary steps in each catalytic cycle might be controlled. Such a control, especially combined with other bond forming processes, might provide us excellent opportunities to develop new reactions/

Fax: 1-718-982-3910; E-mail: qiaohu@mail.csi.cuny.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

processes and thus could make the already powerful transition metal-catalyzed cross-coupling reactions even more powerful for organic synthesis. Toward this end, we have recently documented Pd(0)/*t*-Bu₃P-catalyzed Suzuki cross-coupling of dihaloarenes with arylboronic acids,³ a process that relies on the control of oxidative addition step. We have also reported Pd(0)/*t*-Bu₃P-catalyzed reaction of 2-bromobiphenyl with hindered Grignard reagents to form substituted fluorenes (Scheme 1),^{4–6} a cyclization process that combines the control of the transmetalation step with *sp*³ C-H activation (Scheme 1).^{7–9} We have further incorporated a cross-coupling step into the latter type of reaction and developed a new type of tandem reaction that combines the cross-coupling strategy and cyclization via *sp*³ C-H activation strategy (Scheme 2).^{4,10,11} With Pd(0)/*t*-Bu₃P as catalyst, substituted fluorenes could be easily prepared in excellent yields from 1,2-dibromobenzenes and 1-bromo-2-iodobenzene.

Although Pd(0)/*t*-Bu₃P has been demonstrated to be a highly efficient catalyst system for the tandem reaction of 1,2-dibromobenzenes and 1-bromo-2-iodobenzene to form substituted fluorenes, we reasoned that further improvement of this new substituted fluorene making process could be possible. For example, the air-sensitivity of *t*-Bu₃P makes the handling of the catalyst very stringent. In addition and more importantly, Pd(0)/*t*-Bu₃P catalyst system was ineffective for substrates other than 1,2-dibromobenzenes and 1-bromo-2-iodobenzene, e. g., 1-chloro-2-halobenzenes. To expand the substrate scope and explore more practical catalyst systems, we envisioned a pathway other than the cross-coupling-cyclization (via *sp*³ C-H activation) strategy, namely palladium associated aryne forming pathway. We investigated factors that could influence the reaction to proceed via the palladium associated aryne forming pathway including the ligand effect and leaving group effect. We also investigated the substrates other than 1,2-dibromobenzenes and 1-iodo-2-bromobenzene, i. e., 1-chloro-2-halobenzenes, 1-bromo-2-fluorobenzene and 2-haloaryl tosylates and benzenesulfonates for the reaction. Herein, our detailed study is reported.¹²

Results and Discussions

1. Palladium associated arynes as intermediates for the reaction of 1,2-dihalobenzenes with Grignard reagents: ligand effect and leaving group effect

Our mechanistic study in Pd(0)/*t*-Bu₃P catalyst system suggested that the key obstacle to employ 1-chloro-2-halobenzenes as substrates for the reaction is the reluctance of the C-Cl bond in **I** to undergo oxidative addition with Pd(0) catalyst (Scheme 3).⁴ We thus envisaged that if no oxidative addition of C-Cl bond with Pd(0) was involved in the formation of ArPd(II)Cl (**II**), these substrates could then be suitable substrates for the new type of annulative tandem reactions. Recognizing that halo groups are known to function as leaving groups, we reasoned that this could be possible: as depicted in Scheme 4, *ortho*-leaving group (LG) bearing *o*-aryl(LG)palladium(II) halides (**IV**), in addition to undergoing transmetalation with organometallic reagents to form **I** (Scheme 4, Path Aa), could undergo β-LG-elimination to form palladium associated arynes (**V**) (Scheme 4, Path Ab). The generated Pd(II)(LG)X associated arynes could then undergo transmetalation followed by carbopalladation to form intermediate **II**, which could further undergo other transformations (Scheme 4, Path Ab). Therefore, through palladium associated aryne intermediates, intermediate **II** could be formed without undergoing the oxidative addition of C-LG bond with Pd(0), thus making it possible to employ 1-chloro-2-halobenzenes as suitable substrates for the domino reaction.

Arynes are very reactive and have recently been demonstrated as useful substrates for palladium-catalyzed carbon-carbon bond forming reactions.^{13–17} Careful examination of aryne chemistry showed that the aryne generation strategy in reported reactions, in which arynes were generated *in situ* from expensive *o*-trimethylsilylaryl triflates, often does not work well with the use of catalytic amount of palladium because short-lived arynes needed to search for inherently unstable palladium intermediates for reactions to occur. Consequently, large

excess of *o*-trimethylsilylaryl triflates were often required to achieve good yields. The pathway to generate arynes depicted in Scheme 4 (Path Ab) could employ widely available *o*-halo(LG) benzenes including 1,2-dihaloarenes as substrates with the assurance of every generated aryne associated with a palladium. More importantly, in this strategy, because the formation of **II** does not involve the oxidative addition of C-leaving group bond, e. g., C-Cl bond, with Pd(0) species, the overall reactivity of *o*-halo(LG)benzenes would be governed by the reactivity of C-halo bonds that undergo the initial oxidative addition with Pd(0) catalysts. Thus, substrates that bear leaving groups other than Br and I groups might also be suitable substrates for tandem/domino reactions. Therefore, exploration of this palladium associated aryne generation strategy could be fundamentally interesting and synthetically useful.

As depicted in Scheme 4, the direct transmetalation of *o*-aryl(LG)palladium(II) halides (**IV**) to form cross-coupling product **I** (Scheme 4, Path Aa) would compete with the formation of Pd(II)(LG)X associated benzyne (Path Ab). Therefore, factors that could influence the transmetalation and/or the β -leaving group elimination of *o*-aryl(LG)Pd(II) halides, e. g., the steric hindrance and nucleophilicity of organometallic reagents, LG leaving ability, ligand effect, basicity of the base and reaction temperature, etc., were expected to affect the generation of Pd associated arynes. We reasoned that for a given type of *o*-aryl(LG)Pd(II) halides with the same LG and a given type of organometallic reagents such as Grignard reagents at a certain reaction temperature, the influential factors could be narrowed down to steric hindrance and ligand effect. Since increasing the steric hindrance of Grignard reagents has been established to slow down the transmetalation process, we thus began our study by examining the ligand influence. The reaction of 1-bromo-2-chlorobenzene with bulky 2-mesitylmagnesium bromide was employed as the model reaction. As the initial oxidative addition was believed to occur at the C-Br bond^{18,19} and 2-chloro-2',4',6'-trimethylbiphenyl (**Ia**, LG = Cl) was found to be inert under Pd(OAc)₂/Grignard reagent condition (Scheme 5), we expected that the domino reaction product 2,4-dimethylfluorene would be expected if the reaction proceeded via benzyne intermediate (Path Ab). The reaction product would be 2-chloro-2',4',6'-trimethylbiphenyl if the reaction occurred via transmetalation followed by reductive elimination (Path Aa). We screened a number of ligands and palladium source and our results are listed in Table 1. We found that with monodentate PPh₃, PCy₃, *t*-Bu₃P, a *N*-heterocyclic carbene (NHC),²⁰ or bidentate DPPE, DPPB and BINAP as ligands, 2-chlorobiaryl was obtained as the major product (Table 1, entries 1–8), suggesting the reactions proceeded predominately through Path Aa. However, 2,4-dimethylfluorene was observed as the major product with PdCl₂, Pd₂(dba)₃, Pd(OAc)₂, Pd(PhCN)₂Cl₂ or Pd/C (10% weight) as catalysts (Table 1, entries 9–13), implying that the reaction proceeded predominately through Path Ab and Pd-associated benzyne were involved.

The ligand effect on reaction pathways was further confirmed by the reaction results of 1-bromo-2-chlorobenzene with less sterically hindered *p*-tolylmagnesium bromide. We expected that 4,4''-dimethyl-(1, 1', 2', 1'')-terphenyl (**VII**) would be the product if the reaction occurs via palladium associated aryne pathway (Path Ab) and 2-chloro-4'-methylbiphenyl (**Ia**, LG = Cl) would be the product via transmetalation followed by reductive elimination (Path Aa) (Scheme 4). Our results showed that with *t*-Bu₃P as ligand, 4,4''-dimethyl-(1, 1', 2', 1'')-terphenyl and 2-chloro-4'-methylbiphenyl were formed in a ratio of 7: 93 (Based on ¹H NMR), suggesting Path Aa should be the major pathway. However, with only Pd(OAc)₂ as the catalyst, 4,4''-dimethyl-(1, 1', 2', 1'')-terphenyl was obtained as the major product (Scheme 6). These results further suggested that the absence of *t*-butylphosphine favored the reaction occur via palladium associated aryne intermediates. Comparing these results with the reaction result of 1-bromo-2-chlorobenzene with 2-mesitylmagnesium magnesium (Table 1, entry 11) further showed that increasing the steric hindrance of Grignard reagents would slow down the transmetalation process and favor the reaction to proceed via the aryne pathway.

To gauge the leaving group effect on the reaction pathways, we chose 2-mesitylmagnesium bromide as the reagent and studied 1,2-dihalobenzenes with different leaving groups. We first examined 1-bromo-2-halobenzenes (halo = F, Cl, Br) as the substrates and found that the better the leaving group, the higher the ratio of **III:I**, suggesting benzyne intermediates formed easier with better leaving group. (Table 2, entries 1–4). With 1-fluoro-2-halobenzenes as substrates, as 2-fluoro-2',4',6'-trimethylbiphenyl (**IIb**) was found to be also inert under Pd(OAc)₂/Grignard reagent condition (Scheme 5),²¹ formation of 2,4-dimethylfluorene would be expected if the reaction proceeded via benzyne intermediate (Path Ab). With a bromo group as the leaving group, the reaction occurred smoothly at room temperature with exclusive formation of 2,4-dimethylfluorene (for the establishment of palladium associated benzyne pathway, see the section that describes the reactions of 1,2-dibromobenzenes with hindered Grignard reagents (*vide infra*)). For comparison purpose, room temperature reaction of 1-bromo-2-chlorobenzene with 2-mesitylmagnesium bromide was also tested, a conversion of 61% was observed. As the oxidative addition of C-Br bond with Pd(0) for 1-bromo-2-chlorobenzene is expected to occur at the same rate as or higher than that of 1,2-dibromobenzene because of the more electron-withdrawing nature of Cl group than Br group, the lower conversion for 1-bromo-2-chlorobenzene suggested that the benzyne formation occurred slower with a Cl leaving group than that with a Br leaving group.

We have also carried out the reactions of 1-chloro-2-fluorobenzene and 1, 2-dichlorobenzene with 2-mesitylmagnesium bromide. Our results showed that higher ratio of **III:I** was observed for 1,2-dichlorobenzene than that for 1-chloro-2-fluorobenzene (Table 2, entries 5 and 6), and thus further confirmed that benzyne intermediates formed easier with better leaving groups. The higher conversion for 1-chloro-2-fluorobenzene than that of 1,2-dichlorobenzene might be explained by thinking of the C-Cl bond in 1-chloro-2-fluorobenzene being more activated than that in 1,2-dichlorobenzene because a fluoro group is more electron-withdrawing than a Cl group.

2. Pd(OAc)₂-catalyzed annulative reactions of 1-chloro-2-haloarenes and 1-bromo-2-fluorobenzene with hindered Grignard reagents via palladium associated arynes

After establishing Pd(OAc)₂, Pd₂(dba)₃ and Pd(PhCN)₂Cl₂ as excellent catalysts for 1-bromo-2-chlorobenzene to undergo annulative reaction with 2-mesitylmagnesium bromide, we next examined other 1-chloro-2-halobenzenes and hindered Grignard reagents. Our results are listed in Table 3. We found that all tested 1-chloro-2-halobenzenes including 1,2-dichlorobenzene gave good to excellent yields of substituted fluorene products. For 1,2-dichlorobenzene, higher reaction temperature and longer reaction time is needed to achieve good yield, likely because the initial oxidative addition of C-Cl bond with Pd(0) is more difficulty than C-Br bonds. We have also employed 1-bromo-2-fluorobenzene as substrate and a good yield of 2,4-dimethylfluorene was obtained (Table 3, entry 11).

The palladium associated benzyne intermediates could be formed via several possible pathways, e.g., transmetalation followed by benzyne formation (Path A), transmetalation and benzyne formation occurred simultaneously (Path B), or benzyne formation first followed by transmetalation (Path C) (Scheme 7). We reasoned that the results with unsymmetrical 1-halo-2-chlorobenzenes as substrates could be used to differentiate these pathways as different ratio of isomeric substituted fluorenes would be obtained for these pathways. Both Path A and B would give a mixture of fluorene products in a ratio related to oxidative addition rate difference between C-X bond and C-Cl bond. The third pathway, Path C, is expected to give the ratio of products related to the transmetalation rates between Pd-Cl and Pd-X. Previous study on the reaction of 1-bromo-4-chlorobenzene with phenylboronic acid showed that the oxidative addition of Pd(0) occurred almost exclusively at C-Br bond (≥97%).¹⁹ Path A or B would yield the two isomeric fluorenes in a ratio of 97:3 or higher for unsymmetrical 1-

bromo-2-chlorobenzenes. Our results with 3-bromo-4-chlorotoluene and 3-bromo-4-chloroanisole as substrates showed that the two isomeric fluorenes were formed in about 4:1 ratio (Table 3, entries 6–8),²² suggesting that the benzyne intermediates formed first followed by transmetalation. Since C-I bond has also been established to have higher oxidative addition rate than C-Br bond,^{1,20} *o*-chloriodobenzenes would yield two isomeric fluorenes in a ratio close to 100:0 if the reaction proceeded via Path A or B. Our result with 3-chloro-4-iodotoluene as the substrate showed that the two isomeric fluorenes were formed in about 85:15. This observation was consistent with that the palladium associated benzyne intermediates were formed via Path C. Comparison of results of 3-bromo-4-chlorotoluene with that of 3-chloro-4-iodotoluene indicated that the first C-C bond formed mainly at the more reactive C-Br or C-I bond, suggesting that the formed benzyne species (**VI**) didn't undergo coordination flipping (dissociation followed by recoordination in a reversed way). Otherwise, substituted fluorenes with the same ratio would be observed for 3-bromo-4-chlorotoluene and 3-chloro-4-iodotoluene.

The observation of the formation of two isomeric fluorenes in a similar ratio (4~5:1) for 3-bromo-4-chlorotoluene, 3-bromo-4-chloroanisole and 3-chloro-4-iodotoluene (Table 3, entries 6–9) suggested that the transmetalation rates for Pd-Br and Pd-I bonds with hindered Grignard reagents were comparable to each other.

3. Pd(OAc)₂-catalyzed annulative reactions of 1,2-dibromoarenes and 1-bromo-2-iodobenzene with hindered Grignard reagents

Our study on the leaving group ability effect suggested that 1,2-dibromobenzenes should also be excellent substrates for the Pd(OAc)₂-catalyzed annulative reaction. We have thus also employed 1,2-dibromobenzenes as substrates for the reaction and our results are listed in Table 4. We found with Pd(OAc)₂ as catalyst, the reaction could proceed at room temperature for 1,2-dibromobenzene and 3,4-dibromotoluene, and excellent yields of substituted fluorenes were obtained. Unsymmetrical 3,4-dibromotoluene gave two isomeric fluorenes in 42: 58 ratio, suggesting that the reactivity of two C-Br bonds is close to each other. For 1,2-dibromo-4,5-dimethylbenzene, heating to 60 °C was necessary to achieve high yields, likely because of slower initial oxidative addition of C-Br with Pd(0) species. We have also tested 1-bromo-2-iodobenzene as substrate and found it was also an excellent one for the reaction. Our study showed that Pd(OAc)₂ was a highly efficient, operationally convenient catalyst for this reaction.

However, no cyclization product was observed with 2,3-dibromopyridine as the substrate (Table 4, entry 9).

To establish that the Pd(OAc)₂-catalyzed annulative reactions of 1,2-dibromobenzenes with hindered Grignard reagents did not proceed via cross-coupling followed by C-H activation pathway, we adopted the same strategy we used for Pd(0)/*t*-Bu₃P catalyst system: to detect the existence of the intermediate **I** by interrupting the reactions before their completion by using limited amount of Grignard reagent. 2-Bromo-2',4',6'-trimethylbiphenyl (**I**) was found to be inert under the reaction condition (Scheme 8) and have been established to be detectable by GC-MS.⁴ We thus reasoned it could be detected if it was formed in the reaction as the intermediate. We have thus carried out the reaction of 1,2-dibromobenzene with 0.5 equivalent of 2-mesitylmagnesium bromide. Our result showed that no **I** was observed with Pd(OAc)₂ as catalyst (Table 5, entry 1). In contrary, 7% of **I** was observed with Pd(0)/*t*-Bu₃P as catalyst (Table 5, entry 2). The reaction of 3,4-dibromotoluene with 0.5 equivalent of 2-mesitylmagnesium bromide gave similar results: **I** was not detected with Pd(OAc)₂ as catalysts, and 12% **I** observed with Pd(0)/*t*-Bu₃P as catalyst (Table 5, entries 3, 4). In combination with our results that no reaction was observed when mixing **I** with 2-mesitylmagnesium bromide in the presence of Pd(OAc)₂ (Scheme 8), The absence of **I** strongly suggest that the pathway

for the Pd(OAc)₂-catalyzed reaction of 1,2-dibromobenzenes as well as 1-bromo-2-iodobenzene with bulky Grignard reagents should be benzyne formation pathway, rather than the cross-coupling followed by C-H activation pathway.

To understand whether other types of carbon-hydrogen bonds (nonbenzylic C-(1°)H bonds, benzylic C-(2°)H bonds, and benzylic C-(3°)H bonds) could also be activated under our condition, we tested 2-ethyl-6-methylphenylmagnesium bromide and 2-isopropyl-6-methylphenylmagnesium bromide for the domino reaction. We found that the reaction with sterically more hindered Grignard reagents underwent slower (Table 6, entries 1–3). Such a slower reaction could be improved by raising the reaction temperature (Table 6, entries 3 and 4). We also found that the *sp*³ C-H activation occurred at the benzylic methyl groups, suggesting that benzylic C-(1°)H bonds preferentially be activated over nonbenzylic C-(1°)H bonds (nonbenzylic methyl group), benzylic C-(2°)H bonds and C-(3°)H bonds (Table 4, entry 2, Table 6, entries 1, 2). This observation was further confirmed by the fact that no reaction was observed for 2,6-diethylphenylmagnesium bromide (Table 6, entry 4).

4. Pd(OAc)₂-catalyzed annulative reactions of hindered Grignard reagents with 2-haloaryl arenesulfonates

Based on the hypothesis that the reactions of 1,2-dihalobenzenes proceeded with hindered Grignard reagents could proceed via palladium associated aryne intermediates, we found by omitting the use of phosphine and NHC ligands for palladium catalysts, a broad range of 1,2-dihalobenzenes including 1,2-dibromobenzenes, 1-bromo-2-iodobenzene, 1-chloro-2-halobenzenes, and 1-bromo-2-fluorobenzene could be suitable substrates for the preparation of substituted fluorenes. Our palladium associated aryne hypothesis suggested that other types of *o*-halo(LG)arenes, such as 2-haloaryl tosylates (Ts) and benzenesulfonates (Bs) which contain very inert C-OTs or C-OBs bonds, should also be suitable substrates for the domino reaction because the oxidative addition of Pd(0) with the very inert COTs or C-OBs bond would not be involved and the OTs/OBs group would only serve as a leaving group. We thus tested 2-haloaryl tosylates and benzenesulfonates, which are readily available from 2-halophenols,²³ as substrates for the domino reactions (Table 7). We found that with Pd(OAc)₂ as catalyst, good to excellent yields of substituted fluorenes were obtained for 2-iodoaryl tosylates/benzenesulfonates and 2-bromoaryl tosylates/benzenesulfonates (Table 7, entries 1–13). However, 2-chloro-4-methylphenyl tosylate was found to be a poor substrate (Table 7, entry 14). Comparing the reaction results with 2-halo-4-methylphenyl tosylates as substrates (Table 7, entries 6, 7 and 14) revealed that 2-iodo-4-methylphenyl tosylate was a better substrate than 2-bromo-4-methylphenyl tosylate and 2-chloro-4-methylphenyl tosylate was virtually inactive. As the only difference for these substrates was the halo group and the chloro group was more electron-withdrawing than the Br or I group, our results suggested that the initial oxidative addition should occur at the C-X bond, rather than at the C-OTs bond. Otherwise, 2-chloro-4-methylphenyl tosylate would be the better substrate than 2-bromo-4-methylphenyl tosylate and 2-iodo-4-methylphenyl tosylate. This thinking was further confirmed by the fact that no reaction was observed for pyrocatechol ditosylate (Table 7, entry 15). Thus, the reluctance of the C-Cl bond in 2-chloro-4-methylphenyl tosylate to undergo the initial oxidative addition with Pd(0) species excluded it as a suitable substrate for the domino reaction. The observation of two isomers, rather than only one, in similar ratio for 2-bromophenyl tosylates/benzenesulfonates, 2-iodo-4-methylphenyl tosylate and 1-bromo-2-naphthyl tosylate/benzenesulfonate (Table 7, entries 6–13) further suggested that the transmetalation occurred with comparable transmetalation rates for Pd-Br and Pd-I bonds.

Aryl tosylates have been demonstrated to undergo cross-couplings with *para*-substituted phenylmagnesium bromides and *ortho*-tolymagnesium bromide when catalyzed by Pd(0)/Josiphos ligand.²⁴ We have carried out the reaction of phenyl tosylate with 2-

mesitylmagnesium bromide in the presence of 5% Pd(OAc)₂ at 60 °C for 20 h. We found that the cross-coupling product was formed in less than 5% yield, suggesting Pd(OAc)₂ cannot catalyze the cross-coupling of aryl tosylates with hindered Grignard reagents efficiently (Scheme 9). We also found that 2-(2', 4', 6'-trimethyl)-4-methylphenyl tosylate, intermediate that would be formed if the domino reaction proceeded via cross-coupling-oxidative addition-*sp*³ C-H activation mechanism, was unable to be converted to 2,4,6-trimethylfluorene under Pd(OAc)₂/Grignard reagent or Pd(OAc)₂/MgCl₂/Grignard reagent condition (Scheme 10). We have further carried out the Pd₂(dba)₃/*t*-Bu₃P-catalyzed reaction of 2-bromo-4-methylphenyl tosylate with 2-mesitylmagnesium bromide and 2-(2', 4', 6'-trimethyl)-4-methylphenyl tosylate was isolated in 82% yield with no cyclized fluorene being observed (Scheme 11). These results strongly suggested that Pd(OAc)₂-catalyzed domino reactions of 2-haloaryl arenesulfonates with hindered Grignard reagents did not proceed through the cross-coupling-oxidative addition-*sp*³ C-H activation mechanism.

As our results supported that the initial oxidative addition should not occur at the C-OTs bond, the similar ratio of the substituted fluorenes observed for unsymmetrical 2-bromoaryl tosylates/benzenesulfonates, 2-iodo-4-methylphenyl tosylate and 1-bromo-2-naphthyl tosylate/benzenesulfonate excluded the pathway that the transmetalation occur first followed by aryne formation, in which only one isomer of substituted fluorenes would be expected. Our study further suggested that the pathway to form Pd(LG)X associated arynes first followed by transmetalation should be predominant, if not the only one.²⁵

Summary

We demonstrated that simple palladium complexes such as Pd(OAc)₂ were excellent catalysts for 1,2-dihalobenzenes and 2-haloaryl arenesulfonates to undergo annulative reactions with hindered Grignard reagents to form substituted fluorenes. Our study was based on the hypothesis that two mechanistic pathways might be possible for the reaction of 1,2-dihalobenzenes with Grignard reagents, namely (a) the pathway that involves the cross-coupling of 1,2-dihalobenzenes with Grignard reagents to form 2-halobiphenyls followed by oxidative addition and *sp*³ C-H activation, and (b) the pathway that involves the oxidative addition of 1,2-dihalobenzenes to form 2-haloarylPd(II)X complexes followed by β-halo elimination to form palladium associated arynes, transmetalation, carbopalladation and *sp*³ C-H activation. We explored the ligand effect and leaving group effect on these two pathways. We found that these two pathways were highly ligand dependent and the reaction pathway involving palladium associated arynes as intermediates could be controlled by omitting the use of phosphine and NHC ligands for palladium catalysts. We also found that a better leaving halo group favored the benzyne forming pathway and the *sp*³ C-H activation preferentially occurred at benzylic C-(1°)H bonds over nonbenzylic C-(1°)H bonds (nonbenzylic methyl group), benzylic C-(2°)H bonds and benzylic C-(3°)H bonds. Our palladium associated aryne hypothesis allowed us to identify simple palladium complexes, e. g., Pd(OAc)₂, as the catalyst and a broad range of 1,2-dihalobenzenes including previously unsuitable 1-chloro-2-halobenzenes and 2-haloaryl arenesulfonates as substrates for high yield synthesis of substituted fluorenes. Our study also suggested that 2-(leaving group)arylPd(II) complexes most likely formed Pd(leaving group)X associated arynes first followed by transmetalation with Grignard reagents and carbopalladation, rather than undergo transmetalation with Grignard reagent first followed by aryne formation and carbopalladation. The work described here provides a high yield, one-step access to substituted fluorenes from readily available 1,2-dihalobenzenes and 2-haloaryl arenesulfonates and hindered Grignard reagents, and our method may find applications in the preparation of substituted fluorene-containing molecules including polymers.

Experimental section

General

NMR spectra were recorded on Varian 200 MHz or 600 MHz spectrometers. GC-MS experiments were carried out using an Agilent GC/MS instrument consisting of a 6890N series GC and a 5973 Mass Selective Detector System. All yields reported refer to isolated yields (average of two runs) unless otherwise indicated, and the product purity was estimated to be greater than 95% as determined by ^1H NMR. Compounds described in the literature were characterized by comparison of their melting points, ^1H NMR and ^{13}C NMR spectra to reported data. Some of the new compounds were also characterized by elemental analysis.

THF was dried with sodium/benzophenone. 2,6-Dimethylphenylmagnesium bromide, 2-mesitylmagnesium bromide and *p*-tolylmagnesium bromide were purchased from Aldrich Chemical Co. and used directly. $\text{Pd}_2(\text{dba})_3$ and *t*-Bu₃P were purchased from Strem and used as received. $\text{Pd}(\text{OAc})_2$ was obtained as a gift from Frontier Scientific, Inc. 3-Bromo-4-chlorotoluene,²⁶ 3-chloro-4-iodotoluene,²⁷ and 2-iodo-4-methylphenol²⁸ were prepared according to literature procedures. Aryl tosylates and benzenesulfonates were prepared by treating the corresponding phenols with *p*-toluenesulfonyl chloride or benzenesulfonyl chloride in dichloromethane containing excess of triethylamine at room temperature overnight. Bromopentamethylbenzene, pentamethyl phenylmagnesium bromide were prepared according to literature.^{29, 30} Other chemical reagents were purchased from Alfa Aesar and used directly

General procedure of the ligand effect on domino reactions of 1-bromo-2-chlorobenzene with Grignard reagents

In a glove box with N₂-atmosphere, to a mixture of 1-bromo-2-chlorobenzene (1.0 mmol), and 2.0 ml THF (in a Schlenk flask) was added palladium source (0.015 mmol) and phosphine ligands (0.06 mmol). After stirred for 5–10 minutes, Grignard reagent (2.5 ml, 1M in THF, 2.5 mmol) was added. The mixture was allowed to stir under 60°C for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate twice. The organic layer was washed with brine and the solvent was evaporated under vacuum. ^1H NMR analysis of the reaction mixture gave the conversion of the reaction, and the ratio of the products.

General procedure of the leaving group effect on domino reactions of 1,2-dihalobenzenes with Grignard reagents

In a glove box with N₂-atmosphere, to a mixture of 1,2-halobenzene (1.0 mmol), and 2.0 ml THF (in a Schlenk flask) was added $\text{Pd}(\text{OAc})_2$ (0.015 mmol). After stirred for 5–10 minutes, 2-Mesitylmagnesium bromide (2.5 ml, 1M in THF, 2.5 mmol) was added. The mixture was allowed to stir at 60°C or at room temperature for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate twice. The organic layer was washed with brine and the solvent was evaporated under vacuum. ^1H NMR analysis of the reaction mixture gave the conversion of the reaction, and the ratio of the products.

General procedure for Pd-catalyzed domino reactions of 1,2-dihalobenzenes with hindered Grignard Reagents

In a glove box with N₂-atmosphere, to a mixture of 1,2-dihalobenzenes (1.0 mmol) and 1.0 mL THF (in a Schlenk flask) was added palladium acetate (0.03 mmol) or $\text{Pd}_2(\text{dba})_3$ (0.015 mmol). After stirred for 5–10 minutes, Grignard reagent (2.5 mL, 1M in THF, 2.5 mmol) was added. The mixture was allowed to stir at room temperature or 60 °C for 20 h. After being quenched with water, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with brine and the solvent was evaporated under vacuum. Flash chromatography on silica gel (hexane: ethyl acetate = 100:0 to 90:10) gave the products.

General Procedure for Pd(OAc)₂-catalyzed domino reaction of 2-haloaryl tosylates with hindered Grignard reagents

In a glove box with N₂-atmosphere, to a mixture of 2-haloaryl tosylate (1.0 mmol), and 1.0 ml THF (in a Schlenk flask) was added palladium acetate (6.7 mg, 0.03 mmol). After stirred for 5–10 minutes, Grignard reagent (2.5 ml, 1M in THF, 2.5 mmol) was added. The mixture was allowed to stir under refluxing for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with brine and the solvent was evaporated under vacuum. Flash chromatography on silica gel (hexane: ethyl acetate = 100: 0 to 90: 10) gave the products.

2,4-Dimethylfluorene:³¹ white solid. m.p.: 66.5–67.5°C. ¹H NMR (CDCl₃, 600 MHz): δ 7.87 (d, J = 7.2 Hz, 1H), 7.53 (d, J = 7.2 Hz, 1H), 7.36 (t, J = 7.8 Hz, 1H), 7.27 (t, J = 7.2 Hz, 1H), 7.21 (s, 1H), 6.97 (s, 1H), 3.86 (s, 2H), 2.69 (s, 3H), 2.39 (s, 3H) ppm. ¹³C NMR (CDCl₃, 151 MHz): δ 143.9, 143.5, 142.7, 137.16, 136.2, 132.7, 129.9, 126.5, 125.6, 124.8, 123.1, 122.6, 36.9, 21.3, 20.9 ppm.

4-Methylfluorene:³² white solid. m.p.: 69–71°C. ¹H NMR (CDCl₃, 600 MHz): δ 7.93 (d, J = 7.8 Hz, 1H), 7.56 (d, J = 7.2 Hz, 1H), 7.39 (t, J = 7.2 Hz, 1H), 7.38 (d, J = 7.8 Hz, 1H), 7.30 (t, J = 7.2 Hz, 1H), 7.22 (t, J = 7.8 Hz, 1H), 7.15 (d, J = 7.8 Hz, 1H), 3.91 (s, 2H), 2.73 (s, 3H) ppm. ¹³C NMR (CDCl₃, 151 MHz): δ 143.7, 143.6, 142.7, 139.8, 133.0, 129.0, 126.6, 126.4, 126.0, 124.9, 123.1, 122.4, 37.1, 21.1 ppm.

1,2,3,4-Tetramethylfluorene:³³ white solid. m.p.: 121.5–122.5°C. ¹H NMR (CDCl₃, 600 MHz): δ 7.97 (d, J = 8.4 Hz, 1H), 7.53 (d, J = 7.8 Hz, 1H), 7.34 (dd, J = 8.4, 0.6 Hz, 1H), 7.25 (dd, J = 8.4, 0.6 Hz, 1H), 3.76 (s, 2H), 2.66 (s, 3H), 2.32 (s, 6H), 2.29 (s, 3H) ppm. ¹³C NMR (CDCl₃, 151 MHz): δ 143.7, 143.6, 139.9, 137.0, 133.7, 133.5, 129.4, 129.1, 126.4, 125.3, 124.7, 123.1, 36.6, 16.8, 16.5, 16.5, 16.2 ppm. Anal. calcd for C₁₇H₁₈: C, 91.84%; H, 8.16%. Found: C, 91.58%; H, 8.19%.

2, 4, 6-Trimethylfluorene/2, 4, 7-Trimethylfluorene: off-white solid. m. p.: 43–57°C. ¹H NMR (CDCl₃, 600 MHz): δ 7.74 (d, J = 7.8 Hz, 1H), 7.68 (s, 1H), 7.40 (d, J = 7.8 Hz, 1H), 7.34 (s, 1H), 7.19 (s, 1H), 7.18 (s, 1H), 7.18~7.16 (m, 2H), 7.08 (d, J = 7.2 Hz, 1H), 6.95 (s, 1H), 3.81 (s, 4H), 2.69 (s, 4H), 2.66 (s, 2H), 2.46 (s, 4H), 2.42 (s, 2H), 2.38 (s, 6H) ppm. ¹³C NMR (CDCl₃, 151 MHz): δ 144.4, 143.8, 143.7, 142.9, 140.6, 140.1, 137.2, 137.2, 136.0, 136.0, 135.7, 135.3, 132.6, 132.2, 129.8, 129.8, 127.3, 126.4, 125.6, 124.4, 123.4, 123.1, 123.1, 122.3, 36.7, 36.5, 21.8, 21.4, 21.3, 21.3, 21.0, 20.8 ppm. Anal. calcd for C₁₆H₁₆: C, 92.26%; H, 7.74%. Found: C, 91.93%; H, 7.78%.

The structure of 2, 4, 6-trimethylfluorene was established by NOE effect: Recrystallization on the mixture of 2,4,7-trimethylfluorene/2, 4, 6-trimethylfluorene by hexanes gave >90% pure 2, 4, 6-trimethylfluorene. ¹H NMR (CDCl₃, 600 MHz): δ 7.68 (s, 1H), 7.40 (d, J = 7.8 Hz, 1H), 7.19 (s, 1H), 7.08 (d, J = 7.8 Hz, 1H), 6.95 (s, 1H), 3.80 (s, 2H), 2.68 (s, 3H), 2.46 (s, 3H), 2.38 (s, 3H) ppm. ¹³C NMR (CDCl₃, 151 MHz): δ 144.4, 142.9, 140.6, 137.2, 136.1, 136.0, 132.6, 129.8, 126.4, 124.4, 123.4, 123.1, 36.5, 21.8, 21.3, 21.0 ppm. NOE effect observed when the peak at 3.80 ppm was irradiated: 3.5% for the peak with chemical shift of 7.40 ppm and 3.4% for the peak with chemical shift of 7.19 ppm.

3,5-Dimethylfluorene/2,5-dimethylfluorene: Off-white solid. m.p.: 43–62°C. ¹H NMR (CDCl₃, 600 MHz) δ 7.78 (d, J = 7.8 Hz, 1H), 7.72 (s, 1H), 7.42 (d, J = 7.8 Hz, 1H), 7.36~7.36 (m, 1H), 7.21~7.15 (m, 1H), 7.13~7.11 (m, 1H), 3.84 (s, 2H), 2.72 (s, 3H); 2.70 (s, 3H), 2.46 (s, 3H); 2.42 (s, 3H) ppm. ¹³C NMR (CDCl₃, 151 MHz): δ 144.0, 143.9, 143.4, 142.9, 140.7, 140.0, 139.8, 139.8, 136.04, 135.8, 133.0, 132.6, 128.9, 128.8, 127.4, 126.9, 126.2, 125.9, 125.6, 124.5, 123.8, 122.8, 122.4, 122.3, 36.9, 36.7, 21.8, 21.4, 21.2, 21.0 ppm.

The structure of 3,5-dimethylfluorene was established by NOE effect. Recrystallization of the mixture of 2,5-dimethylfluorene/3,5-dimethylfluorene by hexanes gave > 95% pure 3, 5-dimethylfluorene. m. p.: 77–78.5°C. ¹H NMR (CDCl₃, 600 MHz): δ 7.73 (s, 1H), 7.43 (d, J = 8.4 Hz, 1H), 7.38 (d, J = 7.8 Hz, 1H), 7.19 (t, J = 8.4 Hz, 1H), 7.13 (m, 2H), 3.85 (s, 2H), 2.73 (s, 3H), 2.47 (s, 3H) ppm. ¹³C NMR (CDCl₃, 150.855 MHz): δ 144.0, 142.9, 140.7, 139.8, 136.1, 133.0, 128.9, 126.9, 126.2, 124.5, 123.8, 122.4, 36.7, 21.8, 21.2 ppm. NOE effect observed when the peak at 3.85 (s) ppm was irradiated: 2.9% for the peak with chemical shift of 7.43 ppm and 2.5% for the peak with chemical shift of 7.38 ppm.

3-Methoxy-5-methylfluorene/2-methoxy-5-methylfluorene: obtained as a mixture with a ration of 78:22. The structure of 3-methoxy-5-methylfluorene was established by NOE effect. Recrystallization from the mixture of 3-methoxy-5-methylfluorene and 2-methoxy-5-methylfluorene in hexanes gave 3-methoxy-5-methylfluorene as a white solid. m.p.: 70–71°C. ¹H NMR (CDCl₃, 600 MHz): δ 7.49 (d, J = 2.4 Hz, 1H), 7.45 (d, J = 8.4 Hz, 1H), 7.38 (d, J = 7.2 Hz, 1H), 7.20 (t, J = 7.2 Hz, 1H), 7.14 (d, J = 7.8 Hz, 1H), 7.87 (dd, J₁ = 8.4, 2.4 Hz, 1H), 3.89 (s, 3H), 3.84 (s, 2H), 2.72 (s, 3H). ¹³C NMR (CDCl₃, 151 MHz): δ 158.8, 144.7, 143.9, 139.6, 135.8, 133.0, 128.9, 126.5, 125.1, 122.5, 111.3, 109.4, 55.6, 36.3, 21.0 ppm. NOE effect observed when the peak at δ 3.84 ppm was irradiated: 7.45 (d, 0.3%), 7.38 (d, 0.3%).

2,3,5,7-tetramethylfluorene: white solid. m. p.: 104–105°C. ¹H NMR (CDCl₃, 600 MHz) δ: 7.64 (s, 1H), 7.31 (s, 1H), 7.18 (s, 1H), 6.94 (s, 1H), 3.79 (s, 2H), 2.68 (s, 3H), 2.38 (s, 3H), 2.37 (s, 3H), 2.33 (s, 3H). ¹³C NMR (CDCl₃, 150.871 MHz) δ: 144.0, 141.2, 140.6, 137.4, 135.6, 134.5, 134.0, 132.2, 129.7, 126.0, 123.8, 123.1, 36.5, 21.3, 20.9, 20.3, 20.0 ppm.

2,3,5-Trimethylfluorene: white solid. m. p.: 86.5–87.5°C. ¹H NMR (CDCl₃, 600 MHz) δ: 7.68 (s, 1H), 7.36 (d, J = 7.2 Hz, 1H), 7.32 (s, 1H), 7.16 (t, J = 7.2 Hz, 1H), 7.11 (d, J = 7.2 Hz, 1H), 3.82 (s, 2H), 2.71 (s, 3H), 2.37 (s, 3H), 2.33 (s, 3H). ¹³C NMR (CDCl₃, 150.871 MHz) δ: 143.6, 141.4, 140.6, 140.0, 134.6, 134.5, 132.5, 128.8, 126.0, 125.8, 124.2, 122.3, 36.6, 21.1, 20.3, 20.0 ppm.

4-Ethylfluorene:³⁴ White solid. m. p.: 35 – 36°C. ¹H NMR (CDCl₃, 600 MHz) δ 7.92 (d, J = 7.8 Hz, 1H), 7.57 (d, J = 7.2 Hz, 1H), 7.41 (t, J = 7.2 Hz, 1H), 7.39 (d, J = 7.8 Hz, 1H), 7.31 (t, J = 7.2 Hz, 1H), 7.27 ~ 7.24 (m, 1H), 7.19 (d, J = 7.2 Hz, 1H), 3.92 (s, 2H), 3.12 (q, J = 7.2 Hz, 2H), 1.39 (t, J = 7.2 Hz, 3H). ¹³C NMR (CDCl₃, 150 MHz) δ 143.9, 143.7, 142.1, 139.4, 139.0, 127.1, 126.7, 126.6, 126.0, 124.9, 123.2, 122.5, 37.1, 27.2, 14.3 ppm.

4-Isopropylfluorene:³⁴ Colorless liquid. ¹H NMR (CDCl₃, 600 MHz) δ 7.97 (d, J = 7.8 Hz, 1H), 7.55 (d, J = 7.2 Hz, 1H), 7.39 (d, J = 6.0 Hz, 1H), 7.38 (t, J = 7.8 Hz, 1H), 7.33 (d, J = 7.8 Hz, 1H), 7.30 (t, J = 6.0 Hz, 2H), 3.90 (s, 2H), 3.84 (m, 1H), 1.41 (d, J = 6.6 Hz, 6H). ¹³C NMR (CDCl₃, 150 MHz) δ 144.3, 143.9, 143.8, 142.0, 138.5, 126.7, 126.6, 125.9, 124.9, 123.6, 123.0, 122.4, 37.1, 29.5, 22.8 ppm.

7,9-Dimethyl-11H-benzo[a]fluorene and 9,11-dimethyl-7H-benzo[c]fluorene: Anal. calcd. for C₁₉H₁₆: C, 93.40%; H, 6.60%. Found: C, 93.60%; H, 6.54%. The structure of 7,9-dimethyl-11H-benzo[a]fluorene was established by NOE effect: Recrystallization from the mixture of 7,9-Dimethyl-11H-benzo[a]fluorene and 9,11-Dimethyl-7H-benzo[c]fluorene by using hexanes as solvent gave 7,9-dimethyl-11H-benzo[a]fluorene as a white solid. ¹H NMR (CDCl₃, 600 MHz): δ 8.07 (d, J = 8.4 Hz, 1H), 8.00 (d, J = 8.4 Hz, 1H), 7.90 (d, J = 7.8 Hz, 1H), 7.85 (d, J = 8.4 Hz, 1H), 7.53 (t, J = 7.8 Hz, 1H), 7.45 (t, J = 7.8 Hz, 1H), 7.29 (s, 1H), 7.00 (s, 1H), 4.14 (s, 2H), 2.77 (s, 3H), 2.42 (s, 3H). ¹³C NMR (CDCl₃, 150 MHz): δ 144.0, 140.1, 139.9, 138.1, 135.9, 132.1, 131.9, 130.5, 130.1, 128.6, 127.3, 126.2, 125.1, 124.0, 123.2, 121.6, 35.6, 21.3, 20.8 ppm. NOE effect was observed when irradiated at the peak at δ 4.14 ppm: 8.00 (d, 0.5%), 7.29 (s, 0.3%). NOE effect observed when the peak at δ 2.77 ppm was

irradiated: 8.07 (d, 0.5%), 7.00 (s, 0.29%). NOE effect observed when the peak at δ 2.42 ppm was irradiated: 7.29 (d, 0.22%), 7.00 (s, 0.19%).

9,11-Dimethyl-7H-benzo[c]fluorene: off-white solid. Obtained from the mother liquid of the recrystallization of the mixture of 7,9-dimethyl-11H-benzo[a]fluorene and 9,11-dimethyl-7H-benzo[c]fluorene with >90% purity. ^1H NMR (CDCl_3 , 600 MHz) δ 8.51 (d, J = 8.4 Hz, 1H), 7.88 (d, J = 8.4 Hz, 1H), 7.76 (d, J = 7.8 Hz, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.48 (t, J = 7.8 Hz, 1H), 7.43 (t, J = 7.8 Hz, 1H), 7.21 (s, 1H), 7.07 (s, 1H), 3.90 (s, 2H), 2.82 (s, 3H), 2.39 (s, 3H). ^{13}C NMR (CDCl_3 , 150 MHz): δ 145.1, 142.3, 139.0, 138.4, 135.7, 133.7, 131.6, 131.4, 128.7, 128.6, 127.4, 126.8, 124.5, 124.5, 123.0, 122.9, 38.0, 25.1, 21.0 ppm. NOE effect observed when the peak at δ 3.90 ppm was irradiated: 7.58 (d, 0.5%), 7.21 (s, 0.3%).

7-Methyl-11H-benzo[a]fluorene and 11-methyl-7H-benzo[c]fluorene: Anal. calcd. for $\text{C}_{18}\text{H}_{14}$: C, 93.87%; H, 6.13%. Found: C, 93.69%; H, 6.01%. The structure of 7-methyl-11H-benzo[a]fluorene was established by NOE effect. Recrystallization from the mixture of 7-methyl-11H-benzo[a]fluorene and 11-methyl-7H-benzo[c]fluorene by using hexanes as solvent gave 7-methyl-11H-benzo[a]fluorene as a white solid. ^1H NMR (CDCl_3 , 600 MHz): δ 8.10 (d, J = 8.4 Hz, 1H), 8.01 (d, J = 8.4 Hz, 1H), 7.90 (d, J = 8.4 Hz, 1H), 7.86 (d, J = 8.4 Hz, 1H), 7.53 (t, J = 7.8 Hz, 1H), 7.47 (d, J = 7.2 Hz, 1H), 7.47 (t, J = 7.2 Hz, 1H), 7.22 (t, J = 7.8 Hz, 1H), 7.17 (d, J = 7.8 Hz, 1H), 4.17 (s, 2H), 2.80 (s, 3H). ^{13}C NMR (CDCl_3 , 150 MHz): δ 143.6, 140.7, 140.3, 140.0, 132.5, 132.0, 130.4, 129.2, 128.6, 127.4, 126.3, 126.0, 125.3, 124.1, 122.4, 121.8, 35.7, 21.0 ppm. NOE effect observed when the peak at δ 4.17 ppm was irradiated: 8.10 (d, 0.9%), 7.47 (s, 0.3%).

11-Methyl-7H-benzo[a]fluorene: off-white solid. Obtained from the mother liquid of the mixture of 11-methyl-7H-benzo[c]fluorene and 7-methyl-11H-benzo[a]fluorene by using hexanes as solvent with >85% purity. ^1H NMR (CDCl_3 , 600 MHz) δ 8.51 (d, J = 8.4 Hz, 1H), 7.91 (d, J = 8.4 Hz, 1H), 7.80 (d, J = 8.4 Hz, 1H), 7.62 (d, J = 8.4 Hz, 1H), 7.51 (t, J = 7.2 Hz, 1H), 7.46 (t, J = 7.2 Hz, 1H), 7.42 (d, J = 7.2 Hz, 1H), 7.26 (t, J = 8.4 Hz, 1H), 7.25 (d, J = 7.2 Hz, 1H), 3.97 (s, 2H), 2.86 (s, 3H). ^{13}C NMR (CDCl_3 , 150 MHz): δ 144.7, 142.7, 141.6, 138.3, 133.7, 132.0, 130.6, 128.8, 128.6, 128.6, 127.9, 126.8, 125.9, 124.6, 123.0, 122.0, 38.2, 25.1 ppm. NOE effect was observed when the peak at δ 3.97 ppm was irradiated: 7.62 (d, 0.3%), 7.42 (s, 0.2%).

$\text{Pd}(\text{OAc})_2$ -catalyzed reaction of 2-halo-2',4',6'-trimethylbiphenyl with 2-mesitylmagnesium bromide

In a glove box with nitrogen atmosphere, to a mixture of 2-chloro-2',4',6'-trimethylbiphenyl or 2-fluoro-2',4',6'-trimethylbiphenyl (0.1 mmol), and 0.5 ml THF (in a Schlenk flask) was added $\text{Pd}(\text{OAc})_2$ (0.7 mg, 0.003 mmol). After stirred for 5–10 minutes, Grignard reagent THF solution (0.25 ml, 1M in THF, 0.25 mmol) was added. The mixture was allowed to stir under 60°C for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with brine and the solvent was evaporated under vacuum. No reaction was observed by ^1H NMR.

$\text{Pd}(\text{OAc})_2$ -catalyzed reaction of phenyl tosylate with 2-mesitylmagnesium bromide

In a glove box with nitrogen atmosphere, to a mixture of phenyl tosylate (62mg, 0.25 mmol), and 0.5 ml THF (in a schlenk flask) was added palladium acetate (1.7 mg, 0.007 mmol). After stirred for 5–10 minutes, 2-mesitylmagnesium bromide THF solution (0.6 ml, 1M in THF, 0.6 mmol) was added. The mixture was allowed to stir under 60°C for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with brine and the solvent was evaporated under vacuum. ^1H NMR showed that the yield of the cross-coupling product was less than 5%.

Pd(OAc)₂-catalyzed reaction of 2-chloro-5-methylphenyl tosylate with 2-mesitylmagnesium bromide

In a glove box with nitrogen atmosphere, to a mixture of 2-chloro-5-methylphenyl tosylate (0.25 mmol), and 0.5 ml THF (in a Schlenk flask) was added Pd(OAc)₂ (1.7 mg, 0.0075 mmol). After stirred for 5–10 minutes, Grignard reagent (0.65 ml, 1M in THF, 0.65 mmol) was added. The mixture was allowed to stir under refluxing for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with brine and the solvent was evaporated under vacuum. No reaction was observed by ¹H NMR.

Pd₂(dba)₃/*t*-Bu₃P-catalyzed reaction of 2-bromo-4-methylphenyl tosylate with 2-mesitylmagnesium bromide

In a glove box with N₂-atmosphere, to a mixture of 2-bromo-4-methylphenyl tosylate (170.5 mg, 0.5 mmol), and 1.0 ml THF (in a Schlenk flask) was added Pd₂(dba)₃ (6.9 mg, 0.0075 mmol) and *t*-Bu₃P (6.1 mg, 0.03 mmol). After stirred for 5–10 minutes, 2-mesitylmagnesium bromide (1.25 ml, 1M in THF, 1.25 mmol) was added. The mixture was allowed to stir under refluxing for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate twice. The organic layer was washed with brine and the solvent was evaporated under vacuum. Flash chromatography on silica gel (hexane: ethyl acetate = 100: 0 to 90: 15) gave the cross-coupling product 2-(2', 4', 6'-trimethylphenyl)-4-methylphenyl tosylate in 82% yield. White solid. m.p.: 75–76°C. ¹H NMR (CDCl₃, 600 MHz): δ 7.31 (d, J = 7.8 Hz, 1H), 7.25 (d, J = 8.4 Hz, 2H), 7.15 (dd, J = 7.8, 1.8 Hz, 1H), 7.07 (d, J = 8.4 Hz, 2H), 6.94 (d, J = 1.8 Hz, 1H), 6.80 (s, 2H), 2.40 (s, 3H), 2.35(s, 3H), 2.32 (s, 3H), 1.86 (s, 6H) ppm. ¹³C NMR (CDCl₃, 151 MHz): δ 144.8, 144.3, 136.9, 136.6, 136.6, 133.7, 133.6, 133.1, 132.3, 129.2, 129.0, 127.9, 127.7, 122.6, 21.6, 21.0, 20.9, 20.3 ppm.

Palladium-catalyzed reaction of 2-(2', 4', 6'-trimethyl)-4-methylphenyl tosylate with 2-mesitylmagnesium bromide

In a glove box with N₂-atmosphere, to a mixture of 2-(2', 4', 6'-trimethyl)-4-methylphenyl-tosylate (95 mg, 0.25 mmol), and 0.5 ml THF (in a Schlenk flask) was added Pd(OAc)₂ (1.7 mg, 0.0075 mmol). After stirred for 5–10 minutes, 2-mesitylmagnesium bromide (0.65 ml, 1 M in THF, 0.65 mmol) was added. The mixture was allowed to stir under refluxing for 20 h. After quenched with water, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with brine and the solvent was evaporated under vacuum. No 2,4,6-trimethylfluorene was observed by ¹H NMR.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We gratefully thank the National Institutes of Health (GM69704) for funding. Partial support from the donors of the Petroleum Research Fund, administered by the American Chemical Society (44428-AC1), and the PSC-CUNY Research Award Programs is also gratefully acknowledged. We also thank Prof. Alec Greer at Brooklyn College of CUNY for his help on GC-MS analysis, and Frontier Scientific for its generous gifts of Pd(OAc)₂.

References

1. de Meijere, A.; Diederich, F. Metal-Catalyzed Cross-Coupling Reactions. 2. Wiley-VCH; New York: 2004. Beller, M.; Bolm, C. Transition Metals for Organic Synthesis. Wiley-VCH; New York: 1998. Collman, JP.; Hegedus, LS.; Norton, JR.; Finke, RG. Principles and Applications of Organotransition Metal Chemistry. University Science Books; Mill Valley, CA: 1987.
2. Hu, Q-S. Synthetic Methods for Step-Growth Polymers. Rogers, M.; Long, T., editors. Wiley: 2003. p. 467-526.

3. Dong CG, Hu QS. *J Am Chem Soc* 2005;127:10006–10007. [PubMed: 16011357]
4. Dong CG, Hu QS. *Angew Chem Int Ed* 2006;45:2289–2292.
5. Fluorenes are typically prepared via more than one step, for example: a) Kashulin IA, Nifant'ev IE. *J Org Chem* 2004;69:5476–5479. [PubMed: 15287801] For a recent one-step synthesis of fluorenes: Fuchibe K, Akiyama T. *J Am Chem Soc* 2006;128:1434–1435. [PubMed: 16448098] Also see Ref. 4.
6. Fluorene is the core structure of biologically active molecules, sensors and light-emitting materials. For examples: a) Rathore R, Chebny VJ, Abdelwahed SH. *J Am Chem Soc* 2005;127:8012–8013. [PubMed: 15926815] (b) Sulsky R, Robl JA, Biller SA, Harrity TW, Wetterau J, Connolly F, Jolibois K, Kunselman L. *Bioorg Med Chem Lett* 2004;14:5067–5070. [PubMed: 15380200] c) Marsitzky D, Vestberg R, Blainey P, Tang BT, Hawker CJ, Carter KR. *J Am Chem Soc* 2001;123:6965–6972. [PubMed: 11459474]
7. For recent examples on C-H bond activations: (a) Chen X, Goodhue CE, Yu JQ. *J Am Chem Soc* 2006;128:12634–12635. [PubMed: 17002342] (b) Chen X, Li JJ, Hao XS, Goodhue CE, Yu JQ. *J Am Chem Soc* 2006;128:78–79. [PubMed: 16390130] (c) Thu HY, Yu WY, Che CM. *J Am Chem Soc* 2006;128:9048–9049. [PubMed: 16834374] (d) Wang DH, Hao XS, Wu DF, Yu JQ. *Org Lett* 2006;8:3387–3390. [PubMed: 16836412] (e) Zhou C, Larock RC. *J Org Chem* 2006;71:3551–3558. [PubMed: 16626140] (f) Li Z, Li CJ. *J Am Chem Soc* 2005;127:3672–3673. [PubMed: 15771482] (g) Shi L, Tu YQ, Wang M, Zhang FM, Fan CA, Zhao YM, Xia WJ. *J Am Chem Soc* 2005;127:10836–10837. [PubMed: 16076182] (h) Shabashov D, Daugulis O. *Org Lett* 2005;7:3657–3659. [PubMed: 16092843] (i) Barder TE, Walker SD, Martinelli JR, Buchwald SL. *J Am Chem Soc* 2005;127:4685–4696. [PubMed: 15796535] (j) Desai LV, Hull KL, Sanford MS. *J Am Chem Soc* 2004;126:9542–9543. [PubMed: 15291549]
8. Pd-catalyzed cyclizations via sp^3 C-H activation are rare: Ref. 4. Also see: (a) Baudoin O, Herrbach A, Gueritte F. *Angew Chem Int Ed* 2003;42:5736–5740. (b) Suau R, Lopez-Romero JM, Rico RD. *Tetrahedron Lett* 1996;37:9357–9360. (c) Dyker G. *J Org Chem* 1993;58:6426–6428. (d) Dyker G. *Angew Chem Int Ed* 1994;33:103–105. (e) Dyker G. *Angew Chem Int Ed* 1992;31:1023–1025.
9. For examples on Pd-catalyzed tandem/domino cross-coupling-cyclization via aromatic C-H bond activation, see: (a) Pinto A, Neuville L, Retailleau P, Zhu J. *Org Lett* 2006;8:4927–4930. [PubMed: 17020338] (b) Wegner HA, Scott LT, de Meijere A. *J Org Chem* 2003;68:883–887. [PubMed: 12558411] (c) Larock RC, Tian Q. *J Org Chem* 2001;66:7372–7379. [PubMed: 11681950]
10. For recent general reviews on tandem/domino reactions: a) Tietze LF, Rackelmann N. *Pure Appl Chem* 2004;76:1967–1983. b) Nicolaou KC, Montagnon T, Snyder SA. *Chem Commun* 2003:551–564. c) Parsons PJ, Penkett CS, Shell AJ. *Chem Rev* 1996;96:195–206. [PubMed: 11848750] (d) Tietze LF. *Chem Rev* 1996;96:115–136. [PubMed: 11848746]
11. Recent examples of annulative tandem reactions based on transition metal-catalyzed addition/cross-coupling strategies: (a) Ney JE, Wolfe JP. *J Am Chem Soc* 2005;127:8644–8651. [PubMed: 15954769] (b) Huang Q, Fazio A, Dai G, Campo MA, Larock RC. *J Am Chem Soc* 2004;126:7460–7461. [PubMed: 15198591] (c) Wolfe JP, Rossi MA. *J Am Chem Soc* 2004;126:1620–1621. [PubMed: 14871078] (d) Lira R, Wolfe JP. *J Am Chem Soc* 2004;126:13906–13907. [PubMed: 15506735] (e) Ney JE, Wolfe JP. *Angew Chem Int Ed Eng* 2004;43:3605–3608. (f) Tong X, Li D, Zhang Z, Zhang X. *J Am Chem Soc* 2004;126:7601–7607. [PubMed: 15198608] (g) Cauble DF, Gipson JD, Krische MJ. *J Am Chem Soc* 2003;125:1110–1111. [PubMed: 12553780]
12. A portion of this study has been communicated: Dong CG, Hu QS. *Org Lett* 2006;8:5057–5060. [PubMed: 17048842] Also see: Hu QS. *SynLett* 2007:1331–1345.
13. Recent examples of Pd-catalyzed reactions via carbopalladation of arynes: (a) Henderson JL, Edwards AS, Greaney MF. *Org Lett*. 2007ASAP Article (b) Liu Z, Larock RC. *J Org Chem* 2007;72:223–232. [PubMed: 17194103] (c) Xie C, Zhang Y, Huang Z, Xu P. *J Org Chem* 2007;72:5431–5434. [PubMed: 17555357] (d) Bhuvaneswari S, Jeganmohan M, Cheng CH. *Org Lett* 2006;8:5581–5584. [PubMed: 17107077] (e) Henderson JL, Edwards AS, Greaney MF. *J Am Chem Soc* 2006;128:7426–7427. [PubMed: 16756281] (f) Liu Z, Zhang X, Larock RC. *J Am Chem Soc* 2005;127:15716–15717. [PubMed: 16277509] (g) Jayanth TT, Jeganmohan M, Cheng CH. *Org Lett* 2005;7:2921–2924. [PubMed: 15987170] (h) Jeganmohan M, Cheng CH. *Org Lett* 2004;6:2821–2824. [PubMed: 15281778] (i) Matsubara T. *Organometallics* 2003;22:4297–4304. (j) Yoshida H, Honda Y, Shirakawa E, Hiyama T. *Chem Commun* 2001:1880–1881. (k) Solin N, Narayan S, Szabo KJ. *J Org Chem* 2001;66:1686–1693. [PubMed: 11262114] (l) Yoshikawa E, Yamamoto Y. *Angew Chem, Int Ed* 2000;39:173–175.

14. Recent examples of Pd-catalyzed cyclotrimerization of arynes: (a) Quintana I, Boersma AJ, Pena D, Perez D, Guitian E. *Org Lett* 2006;8:3347–3349. [PubMed: 16836402] (b) Jayanth TT, Jeganmohan M, Cheng CH. *J Org Chem* 2004;69:8445–8450. [PubMed: 15549819] (c) Iglesias B, Cobas A, Perez D, Guitian E, Vollhardt KPC. *Org Lett* 2004;6:3557–3560. [PubMed: 15387547] (d) Sato Y, Tamura T, Mori M. *Angew Chem, Int Ed* 2004;43:2436–2440. (e) Pena D, Cobas A, Perez D, Guitian E, Castedo L. *Org Lett* 2003;5:1863–1866. [PubMed: 12762672] (f) Peña D, Pérez D, Guitián E, Castedo L. *Eur J Org Chem* 2003;7:1238–1243. (g) Pena D, Cobas A, Perez D, Guitian E, Castedo L. *Org Lett* 2000;2:1629–1632. [PubMed: 10841496] (h) Yoshikawa E, Radhakrishnan KV, Yamamoto Y. *J Am Chem Soc* 2000;122:7280–7286. (f) Pena D, Perez D, Guitian E, Castedo L. *J Org Chem* 2000;65:6944–6950. [PubMed: 11031014]
15. For recent examples of Pd-catalyzed carbonylation of arynes: (a) Chatani N, Kamitani A, Oshita M, Fukumoto Y, Murai S. *J Am Chem Soc* 2001;123:12686–12687. [PubMed: 11741437] (b) Yoshikawa E, Radhakrishnan KV, Yamamoto Y. *Tetrahedron Lett* 2000;41:729–732.
16. Pd(0)-coordinated benzyne complexes formed from *o*-Ph[B(OH)₂]Pd(II)XL_n have been reported: (a) Retboll M, Edwards AJ, Rae AD, Willis AC, Bennett MA, Wenger E. *J Am Chem Soc* 2002;124:8348–8360. [PubMed: 12105916] (b) Klosin J, Abboud KA, Jones WM. *Organometallics* 1996;15:2465–2468.
17. For examples of Pd(IV) palladacycles as intermediates: (a) Zhang X, Larock RC. *Org Lett* 2005;7:3973–3976. [PubMed: 16119945] (b) Cardenas DJ, Martin-Matute B, Echavarren AM. *J Am Chem Soc* 2006;128:5033–5040. [PubMed: 16608337]
18. The initial oxidative addition is believed to mainly occur at the C-Br bond in Pd(OAc)₂-catalyzed reaction of 1-chloro-2-bromobenzene with 2-mesitylmagnesium bromide based on the following experiments: Pd(OAc)₂-catalyzed reaction of 1,2-dichlorobenzene with 2-mesitylmagnesium bromide in THF at 60 °C gave low yield of 2,4-dimethylfluorene (24%, much lower than the 95% yield observed for 1-bromo-2-chlorobenzene under the same reaction condition), indicating that the initial oxidative addition at the C-Cl bond occurred very slowly. This observation suggested that with 1-bromo-2-chlorobenzene as substrate, the initial oxidative addition should mainly take place at the C-Br bond.
19. For a recent review: (a) Littke AF, Fu GC. *Angew Chem Int Ed* 2002;41:4176–4211. Also see: (b) Littke AF, Dai C, Fu GC. *J Am Chem Soc* 2000;122:4020–4028.
20. Recent reviews for N-heterocyclic carbenes as Pd(0) ligands: (a) Herrmann WA. *Angew Chem Int Ed* 2002;41:1290–1309. (b) Hillier AC, Grasa GA, Viciu MS, Lee HM, Yang C, Nolan SP. *J Organometallic Chem* 2002;653:69–82. (c) Bourissou D, Guerret O, Gabbai FP, Bertrand G. *Chem Rev* 2000;100:39–91. [PubMed: 11749234]
21. Aryl C-F bond is known to be inert for oxidative addition. For examples of Pd-catalyzed cross-coupling reaction of activated aryl fluorides, see (a) Bahmanyar S, Borer BC, Kim YM, Kurtz DM, Yu S. *Org Lett* 2005;7:1011–1014. [PubMed: 15760126] (b) Kim YM, Yu S. *J Am Chem Soc* 2003;125:1696–1697. [PubMed: 12580584]
22. Two isomeric products in close to 1:1 ratio were always observed with unsymmetrical arynes in the reported aryne generation strategy from *o*-trimethylsilylaryl triflates, see Ref. 13.
23. For examples: (a) Sharghi H, Shahsavari-Fard Z. *Helv Chim Acta* 2005;88:42–52. (b) Harada T, Chiba M, Oku A. *J Org Chem* 1999;64:8210–8213. [PubMed: 11674738]
24. (a) Limmert ME, Roy AH, Hartwig JF. *J Org Chem* 2005;70:9364–9370. [PubMed: 16268609] (b) Roy AH, Hartwig JF. *J Am Chem Soc* 2003;125:8704–8705. [PubMed: 12862447]
25. Whether the formation of arynes would be assisted by the Grignard reagents remains unclear to us at present. Bases were found to be influential on the β-elimination of (2-halo)alkenylPd(II) X complexes: (a) Organ MG, Ghasemi H, Valente C. *Tetrahedron* 2004;60:9453–9461. (b) Andreini BP, Benetti M, Carpita A, Rossi R. *Gazz Chim Ital* 1988;118:469–474.
26. Doyle MP, Siegfried B, Dellaria JF. *J Org Chem* 1977;42:2426–2431.
27. Furniss, BS.; Hannaford, AJ.; Smith, PWG.; Tatchell, AR. *Vogel's Textbook of Practical Organic Chemistry*. 5. 1989. p. P928-929.
28. Holzapfel CW, Bradley D, Williams G. *Tetrahedron* 1995;51:8555–64.
29. Miller AR, Curtin DY. *J Am Chem Soc* 1976;98:1860–1865.
30. Hawkins RT, Lennarz WJ, Snyder HR. *J Am Chem Soc* 1960;82:3053–3059.

31. Abramovitch RA, Chellathurai T, McMaster IT, Takaya T, Azogu CI, Vanderpool DP. J Org Chem 1977;42:2914–2919.
32. (a) Siedle AR, Newmark Richard A, Duerr Brook F, Leung Peter C. J Mol Catal A: Chem 2004;214:187–198. (b) Yamato T, Komine M, Matsuo K. J Chem Res Synop 1997:82–83.
33. Mojelsky TW, Strausz OP. Org Geochem 1986;9:39–45.
34. Cairns JF, Hickinbottom WJ. J Chem Soc 1962:867–70.

Figure 1.
Generalized Mechanism for Pd(0)-Catalyzed Cross-Coupling Reactions

Scheme 1.
Controlling Competition Between Direct Cross-Coupling and Cyclization via sp^3 C-H Bond Activation

Scheme 2.

Pd(0)/*t*-Bu₃P-Catalyzed Tandem Reactions of 1,2-Dihalobenzenes with Hindered Grignard Reagents

Scheme 3.

Pd(0)/*t*-Bu₃P-Catalyzed Tandem Reactions of 1,2-Dihalobenzenes with Bulky Grignard Reagents

Scheme 4.

Outline for 1-Halo-2-(leaving group)benzenes for Domino Reactions via Palladium(II)(LG)X Associated Aryne Intermediates

Scheme 5.

$\text{Pd}(\text{OAc})_2$ -Catalyzed Reactions of 2-Halo-2',4',6'-trimethylbiphenyls with 2-Mesitylmagnesium Bromide

Scheme 6.
Pd(OAc)₂-Catalyzed Reactions of 1-Bromo-2-chlorobenzene with *p*-Tolylmagnesium Bromide

Scheme 7.

Possible Pathways for the Formation of Palladium(II)ClX Associated Aryne Intermediates

Scheme 8.

Pd-Catalyzed Reactions of 2-Bromo-2',4',6'-trimethylbiphenyls with 2-Mesitylmagnesium Bromide

Scheme 9.

Pd(OAc)₂-Catalyzed Reaction of phenyl tosylate with 2-Mesitylmagnesium Bromide

Scheme 10.

Pd(OAc)_2 -Catalyzed Reaction of 2-(2', 4', 6'-Trimethyl)-4-methylphenyl Tosylate with 2-Mesitylmagnesium Bromide

Scheme 11.

$\text{Pd}(0)/t\text{-Bu}_3\text{P}$ -Catalyzed Reaction of 2-Bromo-4-methylphenyl Tosylate with 2-Mesitylmagnesium Bromide

Table 1Ligand Effect on Palladium-Catalyzed Reactions of 1-Bromo-2-chlorobenzene with Grignard Reagents ^a

Entry	Catalyst	Conversion (%)	Ratio (%) ^b	
			I	III
1	1.5% Pd ₂ (dba) ₃ + 6% Ph ₃ P	99	69	31
2	1.5% Pd ₂ (dba) ₃ + 6% Cy ₃ P	99	91	9
3	1.5% Pd ₂ (dba) ₃ + 6% <i>t</i> -Bu ₃ P	99	90	10
4	3% Pd(PPh ₃) ₄	99	88	12
5	1.5% Pd ₂ (dba) ₃ + 6% 	22	99	<1
6	1.5% Pd ₂ (dba) ₃ + 3% DPPE	5	99	<1
7	1.5% Pd ₂ (dba) ₃ + 3% DPPB	27	99	<1
8	1.5% Pd ₂ (dba) ₃ + 3% BINAP	8	99	<1
9	1.5% Pd ₂ (dba) ₃	99	<3	>97
10	3% Pd(PhCN) ₂ Cl ₂	99	<3	>97
11	3% Pd(OAc) ₂	99	<3	>97
12	3% PdCl ₂	93	<3	>97
13	3% Pd/C (10% weight)	11	<3	>97
14	None	0	-	-

^aReaction conditions: 1-bromo-2-chlorobenzene (1.0 equiv.), Grignard reagent (2.5 equiv.), THF (2 ml).^bRatio based on ¹H NMR.

Table 2
Pd(OAc)₂-Catalyzed Reactions of 1, 2-dibromobenzene with 2-MesitylImagnesium Bromide ^a

Entry	X	X'	Temperature	Conversion (%)	I:III ^b
1	F	Br	60 °C	84	17:83
2	Cl	Br	60 °C	99	<3:>97
3	Cl	Br	rt	73	<3:>97
4	Br	Br	rt	99	0:100
5	F	Cl	60 °C	72	63:37
6	Cl	Cl	60 °C	24	3:97

^a Reaction conditions: 1,2-dihalobenzene (1.0 equiv.), Grignard reagent (2.5 equiv.), THF (2 ml).

^b Ratio based on ¹H NMR.

Table 3

Pd(OAc)₂-Catalyzed Cross-couplings of 1-Chloro-2-halobenzenes and 1-Bromo-2-fluorobenzene with Hindered Grignard Reagents ^a

$\text{R}_1\text{-C}_6\text{H}_3\text{Cl-X} + \text{BrMg-C}_6\text{H}_3\text{-R} \xrightarrow[\text{THF, 60 } ^\circ\text{C., 20 h}]{3\% \text{ Pd(OAc)}_2} \text{R}_1\text{-C}_6\text{H}_3\text{-C}_6\text{H}_3\text{-R}$				
entry	dihalide	BrMg-C ₆ H ₃ -R	product	yield(%), ^{bc}
1		BrMg-2,4,6-trimethylphenyl		95
2		BrMg-2,4,6-trimethylphenyl		86
3		BrMg-2,4,6-trimethylphenyl		89
4		BrMg-2,4,6-trimethylphenyl		93
5		BrMg-2,4,6-trimethylphenyl		92
6		BrMg-2,4,6-trimethylphenyl	/	84 ^d
7		BrMg-2,4,6-trimethylphenyl	(77:23) ^e / (80:20) ^e	89 ^d
8		BrMg-2,4,6-trimethylphenyl	(78:22) ^e /	86
9		BrMg-2,4,6-trimethylphenyl	(85:15) ^e /	82
10		BrMg-2,4,6-trimethylphenyl		61 ^f
11		BrMg-2,4,6-trimethylphenyl		59 ^g

^a Reaction conditions (not optimized): 1,2-dihalobenzene (1.0 equiv), Grignard reagent (2.5 equiv), Pd(OAc)₂ (3%), THF (2 mL).

^b Isolated yields.

^c 2-chlorobiaryls were observed in less than 5% yields.

^d 1.5 %Pd₂(dba)₃ as catalyst.

^e Ratio based on ¹H NMR.

^f Refluxing in THF, 40 h, 16% 2-chloro-2',4',6'-trimethylbiphenyl was observed.

^g 17% 2-fluoro-2',4',6'-trimethylbiphenyl was observed.

Table 4

Pd-Catalyzed Cross-Couplings of 1,2-Dibromobenzenes and 1-Bromo-2-iodobenzene with Hindered Grignard Reagents ^a

Entry	Dihalide		Substituted Fluorene	Yield(%) ^d
1				99
2				99
3				92
4				99 (42/58) ^c
5				97 ^d
6				94 ^d
7				99
8				99
9				0

^a Reaction conditions (not optimized): 1,2-dihalobenzene (1.0 equiv.), Grignard reagent (2.5 equiv.), 3% Pd catalyst, THF (2 ml), room temperature.

^b Isolated Yields.

^c ratio based on ¹H NMR.

^d Reaction temperature: 60 °C.

Table 5
Pd-Catalyzed Reactions of 1,2-Dihalobenzenes with Limited Amount of 2-Mesitylmagnesium Bromide ^a

1 equiv. : 0.5 equiv.

Entry	X	R	Pd Catalyst	Conversion(%)	Ratio	
					I	III
1	Br	H	3 % Pd(OAc) ₂	46	0	100%
2	Br	H	1.5% Pd ₂ (dba) ₃ /t-Bu ₃ P	48	7%	93%
3	Br	CH ₃	3 % Pd(OAc) ₂	49	0	100%
4	Br	CH ₃	1.5% Pd ₂ (dba) ₃ /t-Bu ₃ P	45	12%	88%

^aReaction conditions: 1,2-dibromobenzene (1.0 equiv.), Grignard reagent (0.5 equiv.), 3% Pd catalyst, THF (2 ml), r.t., 20 h. Ratios of **I** to **III** were based on GC-MS data.

Table 6Pd-Catalyzed Cross-couplings of 1,2-Dibromobenzene with Hindered Grignard Reagents ^a

Entry			Yield(%) ^b
	Substituted Fluorene		
1			99
2			68
3			29
4			72 ^c
5			<1

^a Reaction conditions (not optimized): dihalide (1.0 equiv.), Grignard reagent (2.5 equiv.), 3% Pd(OAc)₂, THF (2 ml), room temperature.^b Isolated Yields.^c Reaction temperature: 60 °C.

Table 7

Pd(OAc)₂-Catalyzed Reactions of 2-Haloaryl Arenesulfonates with Hindered Grignard Reagents ^a

$\begin{array}{c} \text{R}_1 \\ \\ \text{R}_2-\text{C}_6\text{H}_3-\text{OAs} \\ \\ \text{X} \end{array} + \text{BrMg}-\text{C}_6\text{H}_3-\text{R} \xrightarrow[\text{THF, reflux, 20 h}]{3\% \text{ Pd(OAc)}_2} \begin{array}{c} \text{R}_1 \\ \\ \text{R}_2-\text{C}_{10}\text{H}_6-\text{C}_6\text{H}_3-\text{R} \end{array}$				
Entry	Tosylate	ArMgBr	Product	Yield(%) ^b
1				92
2				87
3				83
4				95
5				86
6				92
7				82
8				74
9				69
10				49
11				79
12				71
13				47
14				< 2
15				< 2

^a Reaction conditions (not optimized): 2-haloaryl arenesulfonate (1.0 equiv), Grignard reagent (2.5 equiv), Pd(OAc)₂ (3%), THF (2 mL), reflux.^b Isolated yields.^c Ratio based on ¹H NMR.