of my Bangladeshi patients I find that their haemoglobin A_{1c} concentration consistently reverts to near normal during long holidays in Bangladesh and doubles inexorably in London; they attribute this to taking less exercise and eating fewer fresh vegetables in London. Group work in the community may be a more effective way of helping understanding and modifying risk for those at high risk.³ It must be recognised that the relation between genetics and past and present environment remains complex and unresolved in relation to an increased risk in south Asians.⁴

ANNA ELERI LIVINGSTONE
General practitioner

Limehouse Practice,
London E14 8HQ

- 1 Gupta S, de Belder A, Hughes LO. Avoiding premature coronary deaths in Asians in Britain. *BMJ* 1995;311:1035-6. (21 October.)
- 2 McKeigue PM, Shah B, Marmot MG. Relation of central obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in south Asians. *Lancet* 1991;337:382-6.
- 3 Kelleher D, Islam S. The problem of integration: Asian people and diabetes. *J R Soc Med* 1994;87:414-7.
- 4 Cruickshank JK. Challenging the orthodoxy of insulin resistance. *Lancet* 1995;346:772-3.

Benefit from controlling blood pressure was understated

EDITOR,—Sandeep Gupta and colleagues press the case for coronary prevention among British Indo-Asians.¹ While we continue research into causes we also have to emphasise what we already know and may not be applying. The authors mention adequate detection and control of high blood pressure only in passing.

Results of controlled trials in groups at high risk, including people in the Medical Research Council's trial in older adults and its American equivalent in systolic hypertension, showed a dramatic benefit from low dose diuretics with or without a potassium retainer, which produced respectively 44% and 27% reductions in coronary events when compared with placebo. Even updated overviews hide this benefit because of the negligible impact of monotherapy with β blockers on coronary events. The issue is not only to detect hypertension and start treatment but then to achieve target blood pressures of <150/90 mm Hg.² Current evidence from surveys suggests that British Indo-Asians have a high rate of hypertension (roughly 30% in middle age), which combines with and adds to the excess risk of diabetes.³ Does that matter?

No direct evidence is available for Indo-Asians. The case is clearest in diabetes in a study of early middle aged Americans followed up for 10-12 years.⁴ There were some 170 coronary deaths and probably double that number of non-fatal events, totalling 512 coronary events per 1000 diabetic men with systolic blood pressure >160 mm Hg. As effective control of blood pressure reduces this figure by a minimum of 16%, some 80 such events could be prevented. If recent evidence from Manchester applies across the nation the wasted opportunity for coronary prevention by proper control of hypertension is considerable: only 14% (55/387) of known hypertensive people with diabetes had blood pressures below 150/90 mm Hg, indicating that some 70 of these 80 potential events would not be prevented. These are conservative estimates.

Once hypertension is established, reaching and maintaining the target of <150/90 mm Hg should be the goal for all patients (particularly Indo-Asians and those at excess risk) as well as nurses, doctors, and purchasers. We have the evidence that it is worth doing.

J K CRUICKSHANK
Senior lecturer

Clinical Epidemiology Unit,
University of Manchester Medical School,
Manchester M13 9PT

- 1 Gupta S, de Belder A, Hughes L. Avoiding premature coronary deaths in Asians in Britain. *BMJ* 1995;311:1035-6. (21 October.)
- 2 Bulpitt CJ, Palmer AJ, Fletcher AK, Beevers DG, Coles EC, Ledingham JG, et al. Optimal blood pressure control in treated hypertensive patients. Report from the Department of Health hypertension care computing project (DHCCP). *Circulation* 1994;90:225-33.
- 3 Cruickshank JK, Cooper J, Burnett M, MacDuff J, Drubra U. Ethnic differences in fasting plasma C-peptide and insulin in relation to glucose tolerance and blood pressure. *Lancet* 1991;338:842-7.
- 4 Stamler J, Vaccaro O, Neaton JD, Wentworth D. Diabetes, other risk factors and 12 year cardiovascular mortality for men screened in the multiple risk factor intervention trial. *Diabetes Care* 1993;16:434-44.
- 5 Cruickshank JK, Du X, Redmond P. Poorly managed hypertension in diabetes: frequency, postural drops and potential for preventing premature stroke and coronary events. *Diabetic Med* 1995;12(suppl 2):541.

Guidelines for pharmacological intervention are needed

EDITOR,—Rates of coronary artery disease in young Asian Indians aged under 40 are three to 10 times higher than those in other populations.¹ I applaud Sandeep Gupta and colleagues for advocating an "effective campaign of primary prevention" of coronary artery disease in Asian Indians² but doubt its efficacy unless there is a diligent search for and aggressive treatment of dyslipidaemias.¹

Among the one million Asian Indians settled in the United States 27 000 are physicians, including 2000 cardiologists. These physicians have made major changes in their lifestyle. In a study of 1478 Asian Indian physicians and their family members in the United States only 68 smoked any cigarettes and 18 smoked over 10 cigarettes a day. Obesity defined as >120% of ideal body weight was noted in 105 (E A Enas *et al*, proceedings of the international symposium on atherosclerosis, Rosemont, IL, US, 6-11 Oct 1991). These physicians have a low saturated fat intake of 14 g (8% of energy) and high levels of exercise (averaging 136 minutes a week).³ Nearly half of them are vegetarians.¹ Yet 110 of the 898 men in the study had coronary artery disease, which is a fivefold higher prevalence than that in American white subjects (12.2% *v* 2.5%) (E A Enas *et al*, proceedings of the international symposium on atherosclerosis, 1991). The rate of admission to hospital for coronary artery disease in Asian Indians in the United States is also four times higher.¹ Contrary to common belief, insulin resistance cannot account for this "Asian Indian paradox," since native Pima American Indians, who have the world's highest rate of glucose intolerance (79%) and a high prevalence of other conventional risk factors, have the lowest prevalence of coronary artery disease of all Americans (six in 546 men (1%))—"the American Indian paradox."

Higher serum lipoprotein(a) concentrations, which are a powerful, independent risk factor for premature coronary artery disease, have been reported in Asian Indians. This concentration is largely genetically determined and minimally influenced by diet and exercise. The pathological effects of high concentrations are augmented by high serum low density lipoprotein, high triglyceride, and low high density lipoprotein concentrations. This "lipid tetrad" best explains the malignant atherosclerosis in young Asian Indians and the resulting high morbidity and mortality from clinical events.¹ Recently a "comprehensive lipid tetrad index" has been proposed to quantify the total burden of dyslipidaemias.⁴ The index is much higher in Asian Indians and is derived by the product of total cholesterol, triglyceride, and lipoprotein(a) values divided by the high density lipoprotein value.

Since serum cholesterol concentrations greatly underestimate the total burden of dyslipidaemia in Asian Indians, specific guidelines that allow

pharmacological treatment to be started at a lower low density lipoprotein concentration and at a lower age are urgently needed. Pharmacological intervention similar to that in secondary prevention of coronary artery disease seems justified as primary prevention in Asian Indians at high risk, because of the persistent excess of coronary artery disease even after intense modification of lifestyle. This strategy may be far more cost effective than waiting and eventually paying for "tertiary prevention," such as coronary angioplasty and bypass surgery.

ENAS A ENAS
Medical director

Coronary Artery Disease in Asian Indians Research,
Woodridge,
Illinois,
60517 USA

- 1 Enas EA, Mehta J. Malignant coronary artery disease in young Asian Indians: thoughts on pathogenesis, prevention and therapy. *Clin Cardiol* 1995;18:131-5.
- 2 Gupta S, de Belder A, Hughes LO. Avoiding premature coronary deaths in Asian Indians in Britain. *BMJ* 1995;311:1035-6. (21 October.)
- 3 Yagalla MV, Song WO, Garg A, Enas EA. Dietary fat, carbohydrate, anthropometrics and leisure-time physical activity as risk factors for atherogenic serum lipids in Asian Indian males in the US. *J Am Diet Assoc* (in press).
- 4 Enas EA. Rapid angiographic progression of coronary artery disease in patients with elevated lipoprotein(a). *Circulation* 1995;92:2353-4.

β Blockers have important role in pulmonary oedema due to mitral stenosis

EDITOR,—I read with interest C A Morley and B A Lim's Lesson of the Week about previously undiagnosed mitral valve disease causing abrupt deterioration in pregnancy.¹ Colleagues and I have had similar experience of young Asian women with pulmonary oedema due to mitral stenosis.

Most of these patients are in sinus rhythm. As the authors say, the patients can develop pulmonary oedema rapidly as tachycardia reduces the time for left atrial emptying during diastole, but the authors fail to mention the important role of β blockers in management. The fall in stroke volume leads to a vicious spiral of a reflex increase in the heart rate, a further fall in stroke volume, and an ever increasing left atrial pressure, all of which can be reversed by β adrenergic blockade.

Giving a β blocker in sufficient dose to counteract the powerful adrenergic drive and slow the heart rate to 80 or 90 beats/min has an almost miraculous effect in reducing left atrial pressure and is followed by immediate improvement. The left ventricle is not in trouble so no inotropes are required (they are best avoided because they raise the heart rate), and only a small initial dose of diuretic is needed. This gives time to plan the best strategy and, if necessary, to transfer the patient safely. Most of these patients' valves are suitable for balloon valvuloplasty, but before this procedure was available closed mitral valvotomy was much safer for the fetus than open valvotomy during cardiac bypass.

Familiarity plus tachycardia was probably responsible for the admitting team's failure to recognise the auscultatory features of mitral stenosis. However, this coupled with the delay in performing chest radiography or echocardiography, let alone in getting a cardiologist, make this a lesson in missed diagnosis and consequent mismanagement.

CELIA M OAKLEY
Professor of clinical cardiology

Hammersmith Hospital,
London W12 0NN

- 1 Morley CM, Lim BA. The risks of delay in diagnosis of breathlessness in pregnancy. *BMJ* 1995;311:1083-4. (21 October.)