sion of HIV can be reduced to less than 5% is almost unbelievable—but he does not stop there. He then wags a righteous finger at women who do not know they have HIV infection for not finding out, by not having themselves tested antenatally, in spite of all the known reasons why women do not get tested antenatally. The Department of Health had a consultation with a large group of women from a range of voluntary organisations involved in HIV/AIDS work in January 1994, in which the problematic issues of antenatal testing were thoroughly aired. It seems that the message—that the supposed benefits to both mother and baby are not at all clear cut—has been lost.

One hospital, Steer says, emphasises to women that “knowing their HIV status may save their baby’s life.” In so many words, he is saying that 95 out of 100 babies of HIV positive women need no longer be at risk of HIV infection. This is patently false. It is ridiculous and dangerous to reduce the situation in this manner, especially since someone might believe him.

MARGE BERER
Editor

Reproductive Health Matters,
London EC1M 3JB

- 1 Steer P. Recent advances in obstetrics. *BMJ* 1995;311:1209-12. (4 November.)
- 2 Vernal DAA. Practising obstetrics and gynaecology in areas with a high prevalence of HIV infection. *Lancet* 1995;346:293-5.
- 3 Dabis F, Mannelbrot L, Msellati P, Van de Perre P. Zidovudine to decrease mother-to-child transmission of HIV-1: is it good for developing countries? *AIDS* 1995;9:204-6.

Author's reply

EDITOR,—My review (PS) recommended that pregnant women should be advised to have a test for HIV antibody so that if the results were positive they could opt for interventions to minimise the risk of transmitting the virus to their baby.¹ The discussion of HIV infection was but a small component of the paper, and it was not possible to cite all the documentary evidence in support of this proposal.

We stand by the statement that the evidence for the role of breast feeding in the transmission of HIV is incontrovertible. In our maternity service 85% of mothers breast feed. We know from anonymous testing during 1994 that nine of the HIV positive women who delivered in this unit were not tested and, as far as is known, were unaware of their HIV infection. If eight of these women breast fed it is possible that one or two of their babies became infected with HIV when this might have been prevented by their avoiding breast feeding. In addition, they were denied the opportunity of discussing delivery by caesarean section or treatment with zidovudine.

People have a right to accurate information, and this is important in informing their choice regarding HIV antibody testing. This is an essential aspect of pretest counselling, whether the patient is pregnant or not. When presented with the evidence most mothers we look after who know that they are HIV positive choose both caesarean section and zidovudine. They clearly disagree with Marge Berer's conclusions.

Berer refers to a meeting she attended, together with a variety of representatives from women's groups, hosted by the Department of Health in January 1994. One of us (JRS) attended such a meeting, but after two days of discussions no agreement could be reached between the women's groups' representatives and the medical staff present, with the former being reluctant to accept the value of the scientific evidence being reviewed.

It is not clear to us why Berer takes such issue with our attempts to offer women the choice of methods which we believe reduce vertical transmission. We strongly support further trials of these and other strategies but feel that the current evidence justifies us recommending to women that

they consider interventions to benefit the health of their children.

PHILIP STEER
Professor
J RICHARD SMITH
Senior lecturer

Department of Obstetrics and Gynaecology,
Charing Cross and Westminster Medical School,
London SW10 9NH

SIMON BARTON
Consultant

John Hunter Clinic,
Chelsea and Westminster Hospital,
London SW10 9NH

- 1 Steer P. Recent advances in obstetrics. *BMJ* 1995;311:1209-12. (4 November.)

Evidence-Based Medicine

Commentaries should be evidence based too

EDITOR,—As enthusiastic supporters of the recent drive towards evidence based health care we eagerly looked forward to the first issue of the journal *Evidence-Based Medicine*, published by the BMJ Publishing Group.¹ When we read it, however, we were disappointed both by the way in which it reviews individual articles rather than focusing on areas of clinical practice and by the lack of objectivity of the commentaries on the articles reviewed.

We were struck by the inclusion of unsubstantiated statements in the commentaries, four of 23 commentaries being unreferenced. The value of such unsubstantiated commentaries has recently been questioned.² Furthermore, no detail is given about the process used to select the authors of commentaries, which leads to concerns about the representativeness of the views expressed and about potential conflict of interest.

There has been a welcome trend towards basing conclusions on evidence from a variety of sources, including randomised controlled trials, meta-analyses, and, particularly, systematic reviews of all published and unpublished studies.³ Selecting individual articles from the world literature moves away from this approach. This could potentially undermine the value of systematic reviews as perceived by the clinical community.

We advocate that the commentaries in *Evidence-Based Medicine* should be substantiated by references, that the process for choosing commentators should be published, and that the findings of systematic reviews should be incorporated in the journal. Finally, we would request that future journals published by the BMJ Publishing Group should be encouraged to include a letters section (*Evidence-Based Medicine* does not do so).

N J PEARSON
Senior registrar in public health medicine

J SARANGI
Senior registrar in public health medicine

R FEY
Senior registrar in public health medicine

Avon Health,
Bristol BS2 8EE

- 1 Davidoff F, Hayne B, Sackett D, Smith R. Evidence based medicine. *BMJ* 1995;310:1085-6.
- 2 Tonks A. Reviewers chosen by authors. *BMJ* 1995;311:210. (22 July.)
- 3 Sackett D, Oxman A. *The Cochrane Collaboration handbook*. Oxford: Oxford Collaboration, 1994.

Editors' reply

EDITOR,—Evidence based medicine requires the integration of individual clinical expertise with best available external evidence from systematic research.¹ Good doctors use both. Without the former, practice quickly becomes “evidence tyrannised”; without the latter, practice quickly

becomes out of date. Journals such as *Evidence-Based Medicine* strive to bring both to a readership that can find less than 30 minutes a week to read about subjects relevant to patients. We serve our readers by monitoring both the current literature and other sources (including both the Cochrane and the York reviews) and extracting just the 2% of reports that are both pertinent to front line clinical practice and scientifically rigorous. They are summarised in structured abstracts (as external clinical evidence), and one of our panel of seasoned clinicians (who are required to disqualify themselves if they have any conflicts of interest) places them in clinical context through accompanying commentaries, synthesising the clinical evidence with clinical experience.

The page headed “Purpose and procedure” appears in every issue of the journal and answers the points raised by N J Pearson and colleagues. Because commentators provide the context of personal clinical experience they add references only when they introduce evidence not already cited in the abstracted original. Thus two of the four unreferenced commentaries in our first issue accompanied exhaustive systematic reviews, rendering any further references repetitious. The third commented on placement of a central line under ultrasonographic guidance and provided exactly what was called for: personal clinical experience with the procedure.

Pearson and colleagues' claim that we have ignored systematic reviews is wrong. Our inaugural issue contained a Cochrane review on stroke units and a routinely published systematic review on β agonists, and our next issue contains several more. But good systematic reviews are rare, and they necessarily lag behind important and often definitive trials by several years. Moreover, they are just beginning to address clinically important topics in diagnosis and aetiology.

Although we provide an opportunity for authors to append brief letters to the pages on which their work appears, we have decided not to have a letters section for two reasons. Firstly, authors of the most helpful letters we have received (which are leading to improvements in the journal) do not request publication. Secondly, having a letters section, although it might provide interesting “theatre,” would mean fewer abstracts and commentaries, and we do not think that the trade off would serve our readers.

DAVID L SACKETT
Editor, *Evidence-Based Medicine*

Centre for Evidence Based Medicine,
John Radcliffe Hospital,
Oxford OX3 9DU

R BRIAN HAYNES
Editor, *Evidence-Based Medicine*

Department of Clinical Epidemiology and Biostatistics,
McMaster University,
Hamilton,
Canada

- 1 Sackett DL, Rosenberg WMC, Muir Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ* 1996;312:71-2. (13 January.)

Management of blood loss in children of Jehovah's Witnesses

EDITOR,—David Busuttill and Adrian Coplestone's editorial on the management of blood loss in Jehovah's Witnesses touches on, but does not resolve, the clinical and ethical issues of treating children of Jehovah's Witnesses.¹ The authors concentrate on the use of recombinant erythropoietin in the postoperative management of anaemia at the expense of considering other pharmacological agents, such as aprotinin or desmopressin, which can reduce intraoperative blood loss. As paediatric surgeons we are aware of the need to preserve circulating blood volume and have been early to adopt techniques that provide