Treatment for children with brain tumours

Paediatric oncologists collaborate internationally

EDITOR,—Karol Sikora's editorial¹ and M C G Stevens and colleagues' article² highlight problems in dealing with cancer, especially in childhood. Although over the past two to three decades major advances in overall survival have been made for most children with cancer, some tumours remain a huge challenge. One change has been the expectation of society in general and individual parents in particular that children should not die. This wish obviously motivates families and oncology units to maximise survival and minimise toxicity and to use innovative treatments when primary treatment has failed. However, the increasing trend for families to seek further, often multiple opinions, though understandable, can be damaging for a child patient's physical and emotional wellbeing. The quality of the terminal illness may be considerably worsened by frequent long distance travel and the family's anguish when hyped treatment does not live up to expectations. Sadly, not all clinical colleagues around the world are equally honest and realistic; some, in particular, seem to treat Britain as if it were a large, lucrative source of extracontractual referrals.

Friends, relatives, and the general public living near patients with cancer are often sucked into raising funds for a project that may actually be harmful to the patient and the family. The media must be held accountable for the extensive publicity given to dramatic mercy dashes across the world, which are not accompanied by follow up of the patients. Rarely are all the subtleties of the pros and cons discussed. The press so easily forgets today's human interest story, and yet the patient and family have to go on coping with the consequences of the drama blown up out of proportion by the publicity.

Advice to authors

We receive more letters than we can publish: we can currently accept only about one third. We prefer short letters that relate to articles published within the past four weeks. Letters received after this deadline stand less chance of acceptance. We also publish some "out of the blue" letters, which usually relate to matters of public policy.

When deciding which letters to publish we favour originality, assertions supported by data or by citation, and a clear prose style. Wit, passion, and personal experience also have their place.

Letters should have fewer than 400 words and no more than five references (including one to the BMJ article to which they relate); references should be in the Vancouver style. We welcome pictures.

Letters should be typed and signed by each author, and each author's current appointment and address should be stated. We encourage you to declare any conflict of interest.

Please enclose a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.

As Stevens and colleagues emphasise, it is rare for patients (particularly those with brain tumours) not to be able to receive all aspects of even the most innovative treatment in Britain. Most specialists in paediatric oncology fully recognise and are happy to satisfy the public's increasing desire to seek second opinions. What the public must recognise, however, is the close collaboration among specialists in childhood cancer in Britain, Europe, and the United States; we all speak to each other, communicate our results, and respond if results look superior in a particular centre or country. We share our successes and failures in order to optimise patients' care. Finally, the media too often give the impression that optimal treatment might be withheld because of a particular physician's ignorance or on financial grounds. There is no evidence that this has occurred, and the public should be aware that we will fight to ensure that it will not occur in future.

O B EDEN
Professor of paediatric oncology

Manchester Children's Hospital,
Pendlebury,
Manchester M27 4HA

1 Sikora K. Second opinions for patients with cancer. *BMJ* 1995;311:1179-80. (4 November.)

2 Stevens MCG, Hockley AD, Spooner D, Walsh AR. Treatment for children with brain tumours. *BMJ* 1995;311:1213-4. (4 November.)

Referral pattern should change so that neurosurgeons can develop their experience

EDITOR,—M C G Stevens and colleagues bring welcome, reasoned analysis to an emotive subject.¹ In our attempts to optimise treatment for children with brain tumours in Britain it may be useful to bear in mind two factors that parents of children with tumours with a poor prognosis find particularly appealing.

Firstly, in the United States, partly because of the greater number of patients attracted by centres, a pattern of neurosurgical referral has developed that has enabled individual surgeons to build up unrivalled experience in the management of certain types of tumour. This compares with the more traditional British approach, whereby surgeons tackle whatever cases are referred to their centre. The British approach results in some children being operated on by surgeons who may not specialise in children's brain tumours, let alone have a specific interest in a particular type of tumour.

Secondly, Americans' positive attitude to problem solving is reflected in an optimistic willingness to go on trying, which has understandable appeal for desperate parents.

I believe that we need to improve our referral pattern to help our neurosurgeons to develop their skills and experience. We should be prepared, without prejudicing the best interests of the children, to articulate our determination to fight for their survival. A 50% chance of death can be better expressed as a 50% chance of survival.

ELISABETH A MACDONALD
Consultant paediatric neuro-oncologist

Guy's Hospital,
London SE1 9RT

1 Stevens MCG, Hockley AD, Spooner D, Walsh AR. Treatment for children with brain tumours. *BMJ* 1995;311:1213-4. (4 November.)

Some treatment is only available abroad

EDITOR,—I am the parent of a child who had a brain tumour with a poor prognosis and who was treated at New York University by Professor Patrick Kelly. Consequently, I wish to comment on M C G Stevens and colleagues' concerns that apparently innovative treatments available outside Britain that offer better prospects for cure than treatments available in Britain have attracted media interest.¹

Their article suggests that Britain is at the forefront in all areas of neurological research; it fails to acknowledge that techniques that are available only outside Britain have not only been trialled but also become established.² The authors argue that benefits have been claimed for innovative treatments on the basis of unscrutinised data. If they refer to Kelly's published research data they will discover that his authority and methodology pass all scrutiny.

The authors question the validity of innovative treatments and then acknowledge that responsible and informed innovation is a necessary part of the development of new treatment strategies. The book *Computers in Stereotactic Neurosurgery* shows that much responsible and informed innovation is taking place outside Britain.³

The authors mention the need for specialist centres to which referrals can be made. My daughter was initially treated at the Royal Manchester Children's Hospital. It was because her consultant paediatric neurosurgeon was aware of treatments that are available elsewhere that we were able to avail ourselves of Kelly's treatment. The authors refer to specific treatment centres acquiring a higher profile than they deserve. One such centre in Britain received media coverage, which claimed that it could have carried out my daughter's operation. The subsequent inquiry by my area health authority to check the validity of this claim elicited an unpublished admission that the centre could not have carried out the procedure.

Stevens and colleagues state that the public needs to be educated about the process of medical decision making. So do doctors. In my daughter's case I know that better treatment was available elsewhere.

Finally, the authors are concerned about public perceptions of underfunding and resource limitations in the NHS and their consequent effect on patients' confidence. These perceptions may be due to several factors. Political decisions are beyond doctors' control, but perhaps the profession could address factors such as medical politics, ego, and empire building, which may also be undermining the patient-doctor relationship with regard to critical decision making.

R W McCLUCKIE

84 Micawber Road,
Poynton,
Cheshire SK12 1UP

1 Stevens MCG, Hockley AD, Spooner D, Walsh AR. Treatment for children with brain tumours. *BMJ* 1995;311:1213-4. (4 November.)

2 Kelly PJ. Stereotactic imaging, surgical planning and computer assisted volumetric resection of intracranial lesions: methods and results. In: Symon L, ed. *Advances and technical standards in neurosurgery*. Vol 17. New York: Springer-Verlag, 1990: 77-118.

3 Kelly PJ, Kall BA, eds. *Contemporary issues in neurological surgery: computers in stereotactic neurosurgery*. Cambridge, MA: Blackwell Scientific, 1992.