Indications for open surgery

- For a stone that cannot be removed by other means
- In a morbidly obese patient (other procedures are technically impossible)
- In a patient whose poor health precludes other (lengthier) procedures
- For large, complex staghorn calculi

Open surgery is now rare

Choice of surgery

Of the surgical options for urinary stone disease, extracorporeal shock wave lithotripsy is the treatment of choice for most renal stones less than 2 cm in diameter when no evidence of obstruction exists. For stones larger than 2 cm or if the caliceal neck is obstructed then percutaneous nephrolithotomy is preferred. Treating renal stones larger than 2 cm with extracorporeal shock wave lithotripsy is not worth while unless the patient is unfit to undergo another procedure as the results are in general poor, and the risk of ureteric obstruction from stone fragments increases.

Key references

Drach GW. Urinary lithiasis: etiology, diagnosis, and medical management. In: Walsh PC, Retik AB, Stamey TA, Vaughan ED, eds. *Campbell's urology*. Philadelphia: W B Saunders, 1992

Preminger GM. Renal calculi: pathogenesis, diagnosis and medical therapy. *Semin Nephrol* 1992; 12: 200-16

Stones in the upper ureter may be treated with in situ lithotripsy, provided that no evidence of sepsis exists in an obstructed upper tract. If sepsis is present then a percutaneous nephrostomy should be performed urgently to relieve the obstruction and drain the kidney. Retrograde manipulation of the stone into the renal pelvis with a ureteric stent has been largely abandoned as the results of in situ lithotripsy are equivalent, and the morbidity associated with a ureteric stent is thus avoided. Lithotripsy of the mid-ureter is difficult because stones may not be easy to see in this area. Open surgery may be needed because stones located here may be too low for antegrade percutaneous removal but too high for easy ureteroscopic removal. Stones in the lower ureter or at the vesicoureteric junction may be treated with lithotripsy or with ureteroscopy with similar results. Stones in the bladder may be treated initially with lithotripsy. Most bladder stones result from bladder outflow obstruction (for example, due to benign prostatic hyperplasia or a urethral stricture), and this condition should also be treated to prevent a recurrence.

The ABC of Urology is edited by Chris Dawson, a senior registrar in urology at the Edith Cavell Hospital, Peterborough, and Hugh Whitfield, a consultant urologist at the Central Middlesex Hospital and the Institute of Urology and Nephrology, London.

Lesson of the Week

Recurrent pelvic endometriosis and bilateral ureteric obstruction associated with hormone replacement therapy

R J Brough, K O'Flynn

Endometriosis after the menopause may be reactivated by hormone replacement therapy

Between 2.5% and 15% of women of reproductive age have endometriosis,¹ yet the ureter is only rarely affected.² We report the case of a patient presenting with bilateral ureteric obstruction secondary to recurrent endometriosis. She had previously undergone a total hysterectomy and bilateral salpingo-oophorectomy for endometriosis and had been taking unopposed oestrogen for postmenopausal symptoms in the five years after surgery.

Case report

A 47 year old health care worker was suddenly admitted to hospital; she complained of three months of increasing bilateral loin pain, change in bowel habit, and general malaise. Six years earlier she had had a hysterectomy and bilateral salpingo-oophorectomy for endometriosis. Surgery showed that she had extensive disease in the pelvis with adhesions obliterating the pouch of Douglas. She was given a six month course of danazol because of the likelihood of residual disease. Histological examination of the ovaries and uterus showed extensive endometriosis affecting the peri-

toneum and both ovaries, with haemorrhagic adhesions throughout the specimen. Hormone replacement therapy in the form of a combined oestrogen and progestogen pill was offered after the course of danazol, but the patient declined.

Two years after surgery she complained of hot flushes, irritability, and vaginal soreness. Her general practitioner diagnosed postmenopausal symptoms, and she was started on a course of regular oral oestrogen (ethinyloestradiol, 10 µg a day). Shortly afterwards she began to have intermittent right loin pain and was referred to the general surgeons, who diagnosed early diverticular disease on colonoscopy and took no further action. She continued taking hormone replacement therapy up to the time of her acute admission.

At that time pelvic examination showed a lower abdominal mass, anterior to the rectum. Serum urea and electrolyte concentrations were normal. An intravenous urogram and renogram showed the right kidney to be completely obstructed and contributing only 11% to overall renal function. The left kidney was partially obstructed and contributing 89% of all renal

Department of Urology,
Manchester Royal
Infirmary, Manchester
M13 9WL
R J Brough, research registrar
K O'Flynn, senior registrar

Correspondence to:
Mr Brough.

BMJ 1996;312:1221-2

function (fig 1). A pelvic ultrasound scan and computed tomogram both showed a soft tissue mass 9 cm in diameter, anterior to the rectum (fig 2).

Cystoscopy and examination under anaesthesia were performed, and a biopsy specimen was taken from the mass. Histologically this was confirmed to be a recurrence of endometriosis with no sign of malignancy. An attempt was made to insert stents into both ureters at the time of cystoscopy, but this was unsuccessful. Consequently, a nephrostomy tube was inserted to decompress the left kidney, and later an antegrade double J stent was manipulated between the left kidney and the bladder. The right kidney was left untreated owing to its low level of function.

Oral oestrogen was discontinued and the patient given a monthly depot dose of goserelin, a synthetic gonadotrophin releasing hormone agonist, administered as a 3.6 mg implant. The implant suppresses oestrogen production from the ovaries and was used in this case because of the small risk of residual ovarian tissue being left after the original operation and to accelerate the involution of the endometriosis.

Subsequent computed tomography (fig 3) showed a progressive decrease in the size of the endometriotic mass, and the double J stent was removed six weeks

after treatment had been started. Follow up renography showed good function and satisfactory elimination from the left kidney but little change in the relative function of the right side. The depot injections of goserelin were stopped after six months, and the patient has been well since, with no change in function or drainage from the left kidney or increase in the size of the mass.

Discussion

Conservative surgery for endometriosis has a five year recurrence rate of about 40%.³ More radical surgery in patients with severe disease who no longer want a family is aimed at removing all the ovarian and uterine tissue together with any endometrial deposits in the pelvis. Langmade suggested that hormone replacement therapy would not cause recurrence of the disease if all the ovarian tissue had been removed.⁴ More recent reports, however, have shown that in postmenopausal patients endometriosis can be reactivated by oestrogen treatment^{5,6}; indeed, the recurrence may be more severe than the original disease, with a greater chance that the ureters will be affected.⁷

The incidence of ureteric disease in endometriosis is estimated to be around 1.2% of cases.⁸ In those patients with ureteric obstruction as many as 25% will sustain irreversible renal damage because of the delay in diagnosis.² Surgery to remove the recurrence and reimplant the ureters used to be the mainstay of treatment in these cases, but with improved interventional radiological techniques and the new gonadotrophin releasing hormone analogues a trial of temporary stent drainage and medication is worth while, with surgery reserved for those failing to respond. Although useful in the treatment of endometriosis, danazol, a synthetic derivative of testosterone, does not seem to alter the course of ureteric obstruction once fibrosis has occurred.⁹ The use of unopposed oestrogen treatment in postoperative patients has been questioned, and Manyonda *et al* have suggested that the addition of a progestogen might have a depressive effect on any residual endometriosis and have recommended its use in hormone replacement therapy after radical surgery for endometriosis.⁵

It is suggested that routine imaging, either intravenous urography or preferably isotope renography, should be used to assess any patient with endometriosis^{4,5,10} to prevent the fibrotic complications and permanent renal damage that can occur with this disease. The possibility of ureteric obstruction should be considered in all patients with a history of endometriosis who are receiving hormone replacement therapy, even if they have already undergone radical surgery for the disease.

Fig 1—Renogram at time of presentation

Fig 2—Computed tomogram showing soft tissue mass before treatment

Fig 3—Computed tomogram after treatment

- 1 Barbieri RL. Etiology and epidemiology of endometriosis. *Am J Obstet Gynecol* 1990;162:565-7.
- 2 Moore JG, Hibbard LT, Growdon WA, Schiffrin BS. Urinary tract endometriosis: enigmas in diagnosis and management. *Am J Obstet Gynecol* 1979;134:162-72.
- 3 Wheeler JM, Malinak LR. Recurrent endometriosis: incidence, management, and prognosis. *Am J Obstet Gynecol* 1983;146:247-50.
- 4 Langmade CF. Pelvic endometriosis and ureteral obstruction. *Am J Obstet Gynecol* 1975;122:463-9.
- 5 Manyonda JT, Neale EJ, Flynn JT, Osborn DE. Obstructive uropathy from endometriosis after hysterectomy and oophorectomy; two case reports. *Eur J Obstet Gynecol Reprod Biol* 1989;31:195-8.
- 6 Dick I, Lang DW, Bergman RT, Bhatnagar BNS, Selvaggi F. Postmenopausal endometriosis with ureteral obstruction. *Br J Urol* 1973;45:153-5.
- 7 Lam AM, French M, Charnock FM. Bilateral ureteric obstruction due to recurrent endometriosis associated with hormone replacement therapy. *Aust NZ J Obstet Gynaecol* 1992;32:1:83-4.
- 8 Ball TL, Platt MA. Urological complications of endometriosis. *Am J Obstet Gynecol* 1963;84:1516-21.
- 9 Rivlin ME, Krueger RP, Wiser WL. Danazol in the management of ureteral obstruction secondary to endometriosis. *Fertil Steril* 1985;44:274-6.
- 10 Kane C, Drouin P. Obstructive uropathy associated with endometriosis. *Am J Obstet Gynecol* 1985;151:207-11.

(Accepted 9 January 1996)