GP's often fail to give early parenteral benzylpenicillin in suspected meningococcal infection

EDITOR,—Recent national guidelines and communications from the chief medical officer have emphasised the importance of early benzylpenicillin in the management of suspected invasive meningococcal disease.¹ Such action may prove life saving.² The only contraindication for the use of benzylpenicillin in such circumstances is anaphylaxis after previous treatment with penicillin.¹

In a recent study Colbridge *et al* found that 85% of general practitioners carried benzylpenicillin in their emergency bags.³ In Rotherham the local medical audit advisory group recently conducted an audit which found that 107 of 114 general practitioners carried benzylpenicillin in their emergency bags. Unfortunately, carriage of benzylpenicillin does not equate with its use when required.⁴

Between 8 December 1995 and 11 January 1996 an outbreak of meningococcal septicaemia occurred in Rotherham.⁵ General practitioners saw five of the patients before their admission to hospital. Parenteral benzylpenicillin was given to one of these patients. In one case benzylpenicillin was not given because a different diagnosis was considered, and in another case the patient referred himself to hospital. In seven cases of suspected invasive meningococcal infection that occurred in Rotherham during the following six weeks parenteral benzylpenicillin was given by general practitioners to only one patient despite increased awareness of the condition. In one of these cases the admitting doctor, from a deputising agency, reported that the antibiotic had not been in the emergency bag.

This experience emphasises, yet again, the need for general practitioners to be regularly reminded to give parenteral benzylpenicillin in cases of suspected meningococcal infection in accordance with the chief medical officer's advice and national guidelines.¹ It also highlights the fact that deputising agencies should ensure that benzylpenicillin is routinely carried in emergency bags given to their deputising doctors.

DAVID IRWIN
Consultant in communicable disease control
JAMES M MILLER
Senior registrar in public health medicine
S JOHN CORNELL
Registrar in public health medicine

Rotherham Health Authority,
Rotherham S5 2QU

- 1 Chief Medical Officer. Meningococcal infection. *CMO Update* 1995;7:2.
- 2 Cartwright K, Reilly S, White D, Stuart J. Early treatment with parenteral penicillin in meningococcal disease. *BMJ* 1992; 305:143-7.
- 3 Colbridge MJ, Bailey GG, Dunbar EM, Ong ELC. Antibiotics carried in general practitioners' emergency bags. *BMJ* 1995; 310:29-30.
- 4 Wood A, O'Brien S. Antibiotics carried in general practitioners' emergency bags. *BMJ* 1995;310:737.
- 5 Communicable Disease Surveillance Centre. Meningococcal disease in England and Wales: outbreak due to serogroup C 2b. *Commun Dis Rep CDR Wkly* 1996;6:17.

Admissions due to overdoses of aromatic analgesics have increased in Scotland

EDITOR,—Philip McLoone reports considerable inequalities in suicide rates in Scotland.¹ The largest increases among men in deprived areas were in the rates of self poisoning, and McLoone suggests that restricting the availability of drugs used for self poisoning might reduce deaths by suicide. We have reported previously on the increase in the absolute number of admissions due to parasuicide in Scotland² and have subsequently examined the

nature of this increase. Our findings are relevant to McLoone's suggestion.

In Scottish hospital records the diagnosis of self poisoning can be entered in any one of six diagnosis fields. For completeness, records were extracted if self poisoning was mentioned in any of the fields. Figure 1 shows that the number of admissions related to parasuicide by ingestion of aromatic analgesics (mainly paracetamol) has increased in Scotland, in line with earlier findings in England.³ The admissions comprise all those for which each drug was entered in any position on the SMR1 discharge form, to exclude potential bias from preferential recording of paracetamol in first position because of clinical concern.

Fig 1—Emergency admissions for deliberate self poisoning by type of substance, Scotland, 1981-94

The rate of overdose of aromatic analgesics in males aged 15-24 increased from 31/100 000 in 1981 to 149/100 000 in 1994, and the rate in those aged 25-34 increased from 26/100 000 to 126/100 000. The figures for females were 58/100 000 increasing to 260/100 000 in those aged 15-24 and 32/100 000 increasing to 153/100 000 in those aged 25-34.

Trends in parasuicide and suicide among females continue to diverge.⁴ The link between changes in the epidemiology of suicide and of parasuicide is unclear. We are examining a cohort of 8304 people discharged from Scottish general hospitals in 1981 after parasuicide (*International Classification of Diseases* codes E950-959), who were followed up for 13 years by record linkage. In each age group a higher proportion of males than females went on to complete suicide. In those aged 25-34, for example, 1.4% of the women but 4.4% of the men died by suicide during the 13 years. Further review of these data and the addition of further cohorts will help to identify changes in the relation between suicide and parasuicide over time.

The morbidity and mortality resulting from overdoses of aromatic analgesics require further examination. Public health measures as suggested by McLoone and by Gunnell *et al*,⁵ such as a limit on the number of paracetamol tablets sold at one time, deserve further consideration.

S HAGEN
Epidemiologist
Department of Public Health,
University of Aberdeen,
Aberdeen AB9 1FX

Crichton Royal Hospital,
Dumfries DG1 2SD

Ayrshire and Arran Health Board,
Ayr KA7 4DW

Argyll and Clyde Health Board,
Paisley PA1 1DU

D HALL
Consultant psychiatrist

C STARK
Consultant in public health medicine

H SMITH
Consultant in public health medicine

- 1 McLoone P. Suicide and deprivation in Scotland. *BMJ* 1996; 312:543-4. (2 March.)
- 2 Stark C, Smith H, Hall D. Increase in parasuicide in Scotland. *BMJ* 1994;308:1569-70.
- 3 Hawton K, Fagg J. Trends in deliberate self-poisoning and self-injury in Oxford, 1976-90. *BMJ* 1992;304:1409-11.
- 4 Lester D. Suicide and parasuicide in Scotland. *Perceptual and Motor Skills* 1995;80:862.
- 5 Gunnell D, Frankel S. Prevention of suicide: aspirations and evidence. *BMJ* 1994;308:1227-33.

Children with an avulsed tooth may need antibiotic prophylaxis against bacterial endocarditis

EDITOR,—In their editorial on the emergency management of children with an avulsed tooth Anthony S Blinkhorn and Iain C Mackie mention prophylactic antibiotics.¹ Children with a congenital cardiac abnormality require special consideration—for example, a 10 year old with an avulsed tooth might be given amoxycillin 250 mg three times daily as routine prophylaxis but require a dose of 1.5 g if at risk of bacterial endocarditis.²

One study showed a 61% incidence of bacteraemia in children after the extraction of two to four permanent teeth that were not diseased.³ A cardiac defect is the commonest congenital abnormality. The development of bacterial endocarditis in a child with a small ventricular septal defect may represent a life threatening complication of a previously asymptomatic abnormality. There is evidence of incomplete awareness among parents of the need for prophylaxis against bacterial endocarditis.⁴ When presenting to an accident and emergency department with their child, parents may not appreciate that traumatic avulsion of a tooth and its replantation by a doctor necessitate the same prophylaxis as treatment in a dentist's surgery, particularly if they are preoccupied with a cosmetic defect. The doctor in the minor injuries area of the accident and emergency department may concentrate on the presenting injury without being aware of the child's other medical problems.

The assessment of any child with an avulsed tooth should include a check to see whether prophylaxis against bacterial endocarditis is required. If it is, treatment will be guided by the relevant recommendations on antibiotic prophylaxis.

DANIEL WALLIS
Senior registrar

Accident and Emergency Department,
Newham General Hospital,
London E13 8RU

SUSAN MATTHAI
Senior registrar

Accident and Emergency Department,
Alder Hey Children's Hospital,
Liverpool L12 2AP

- 1 Blinkhorn AS, Mackie IC. My child's just knocked out a front tooth. *BMJ* 1996;312:526. (2 March.)
- 2 Alder Hey book of children's doses. 6th ed. Liverpool: Royal Liverpool Children's Hospital (Alder Hey), 1994.
- 3 Peterson LJ, Peacock R. The incidence of bacteraemia in pediatric patients following tooth extraction. *Circulation* 1976;53:676-9.
- 4 Cetta F, Bell TJ, Podlecki DD, Ros SP. Parental knowledge of bacterial endocarditis prophylaxis. *Pediatr Cardiol* 1993;14: 220-2.

Advance directives are compatible with good medical practice

EDITOR,—Bromley division of the BMA believes that advance directives are incompatible with good medical practice and ethics.¹ I disagree strongly with this view.

Good medical practice requires doctors to advise patients of treatment options (including no treatment at all) and the likely outcomes. Patients may then choose the treatment most acceptable to them. This may be not the treatment that produces the best medical

outcome but the one that produces the outcome that the patient favours. When patients are mentally incompetent an advance directive may express their wishes. Having ascertained that the directive was made in the light of knowledge of the medical facts and a clear understanding of the implications, doctors should be morally and legally obliged to respect the patient's wishes.

The concern that developments in treatment will not be accommodated is unrealistic. Sudden cures for incurable diseases are regrettably rare. If a previously untreatable disease became treatable, however, I believe that the courts would support a doctor who disregarded an advance directive drawn up before the treatment was available.

A more regular occurrence in my practice is that I wish to know an unconscious patient's views on what is an acceptable quality of life, given that he or she is now critically ill in an intensive care unit. Often the best we can achieve is survival with severe neurological damage or other impairment. Once we have ensured survival with aggressive, unpleasant, and costly treatment and have delivered a quality of life that is unacceptable to the patient neither we nor the patient can undo the harm.

An increasing number of patients express their wishes about limitation of treatment before their admission to our unit. Often they are former patients who found the quality of life on discharge barely acceptable and cannot contemplate survival with additional physical or mental impairment. I am happy to respect those wishes, as I would like my wishes to be respected were I in that position. I see no difference between this scenario and that of a Jehovah's Witness who refuses a blood transfusion—a situation in which the law and medical ethics are quite clear.² When a patient's wishes require a doctor to act contrary to his or her conscience the doctor must refer the patient to another practitioner. The alternative, of doctors carrying out treatments contrary to the patient's stated wishes and getting away with it because the patient is not mentally or physically competent to resist, is something that no doctor or the BMA could sanction.

SIMON FINFER
Specialist in intensive care

Royal North Shore Hospital,
Intensive Therapy Unit,
St Leonards,
NSW 2065,
Australia

1 Jessiman IMCD. BMA division disagrees with association's stance over advance directives. *BMJ* 1996;312:850-1. (30 March.)

2 Finfer S, Howell S, Miller J, Willett K, Wilson-MacDonald J, Wilson DH, et al. Managing patients who refuse blood transfusions: an ethical dilemma. *BMJ* 1994;308:1423-6.

Coffee intake and death from coronary heart disease

Coffee may have both short and long term effects

EDITOR,—I Stensvold and colleagues found that a previously reported association between coffee consumption and mortality from coronary heart disease had been weakened by six more years of follow up, and they suggest that this could be due to a decreased intake of boiled coffee.¹ When, however, they excluded subjects who were referred to a doctor because of their coronary risk, and when they analysed more than 23 000 subjects included some years later in an identical study (thus using a more recent evaluation of the distribution of risk factors), the coefficient for coffee drinking was slightly higher.

Between 1990 and 1992 we performed two case-control studies in patients with first confirmed acute myocardial infarction and ischaemic stroke

Table 1—Odds ratio of myocardial infarction and ischaemic stroke according to daily coffee intake

Daily coffee intake (cups)	Myocardial infarction		Ischaemic stroke	
	Cases/controls	Odds ratio* (95% confidence interval)	Cases/controls	Odds ratio* (95% confidence interval)
0	56/97	1.0	57/57	1.0
1	91/134	1.0	61/80	0.6 (0.3 to 1.2)
2	121/156	1.1 (0.6 to 1.8)	63/59	1.4 (0.7 to 2.8)
3	90/59	1.9 (1.0 to 3.4)	30/25	0.7 (0.3 to 1.7)
4	66/32	2.9 (1.3 to 6.5)	10/14	0.6 (0.2 to 1.7)
≥5	89/35	3.8 (1.8 to 7.9)	16/2	15.3 (2.4 to 97.5)

*Conditional multiple logistic model adjusted for social class, education, alcoholic beverages, smoking habit, diabetes mellitus, hypertension, blood cholesterol concentration, body mass index, physical activity, and family history of myocardial infarction and stroke.

confirmed by computed tomography; these were conducted in parallel and with the same operational characteristics (same hospitals, interviewers, etc) in Abruzzo (southern Italy). Altogether 513 patients with acute myocardial infarction (median age 58) and 237 with ischaemic stroke (median age 63) were recruited along with hospital controls (matched by five year age group and sex) with acute diseases not related to cardiovascular risk factors. Our results (table 1) accord with those of a recent meta-analysis² and with the first six years of follow up in Stensvold and colleagues' cohort study and support the existence of a short term effect of coffee intake. No clear relation could be found between coffee intake and ischaemic stroke, possibly because of the low number of subjects with high levels of coffee consumption who were older. In Greenland's meta-analysis recent cohort studies had more positive results than older ones, while a fairly homogeneous increased risk was found in case-control studies.²

Cohort studies cannot easily allow for the changes in lifestyle habits that occur during follow up: coffee intake peaks between ages 40 to 49 and then declines,³ and assessment of coffee consumption 10 or more years before death from coronary heart disease may be a poor indicator of more recent consumption.⁴ In addition, non-fatal cardiovascular events (for example, acute myocardial infarction, angina pectoris, revascularisation procedures) could result in patients stopping smoking as well as reducing their coffee intake. The effect of coffee intake in Stensvold and colleagues' study (this information was collected at baseline) could therefore have been weakened as follow up was prolonged.

In summary, the existence of a short term effect of coffee intake in addition to a long term one (on atherosclerosis) cannot be easily dismissed. Negative results of cohort studies could be associated with a modification of lifestyle habits (including stopping drinking coffee and lowering intake in addition to changing the type of coffee consumed), due either to aging of the cohort or a worsening health status during follow up.

This work was supported by the Italian National Research Council (Convenzione CNR Consorzio Mario Negri Sud).

ROBERTO MARCHIOLI
Senior investigator
ROCCO DI MASCO
Consultant
ROSA MARIA MARFISI
Research assistant
FELICE VITULLO
Research assistant
GIANNI TOGNONI
Head of department

On behalf of the Progetto 3A Investigators

Department of Clinical Pharmacology and Epidemiology,
Istituto di Ricerche Farmacologiche Mario Negri—
Consorzio Mario Negri Sud,
Santa Maria Imbaro (Chieti),
Italy

1 Stensvold I, Tverdal A, Jacobsen BK. Cohort study of coffee intake and death from coronary heart disease over 12 years. *BMJ* 1996;312:544-5. (2 March.)

2 Greenland S. A meta-analysis of coffee, myocardial infarction, and coronary death. *Epidemiology* 1993;4:366-74.

3 Christensen L, Murray T. A review of the relationship between coffee consumption and coronary heart disease. *J Community Health* 1990;15:391-408.

4 Franceschi S. Coffee and myocardial infarction: review of epidemiological evidence. In: Garattini S, ed. *Caffeine, coffee, and health*. New York: Raven Press, 1993:195-211.

Coffee drinking was compared with tea drinking in monozygotic twins in 18th century

EDITOR,—One of the more peculiar attempts to throw light on the question of whether drinking coffee is bad for one's health¹ was carried out in the 18th century by King Gustaf III of Sweden. He is better known for instituting the Swedish Academy, the august body of 18 (18 because the king liked the sound of the Swedish word for that number, *ardeton*) whom Alfred Nobel later selected to award his prize in literature.

A pair of monozygotic twins had been sentenced to death for murder. Gustaf III commuted their death sentences to life imprisonment on the condition that one twin drank a large bowl of tea three times a day and that the other twin drank coffee. The twin who drank tea died first, aged 83—a remarkable age for the time. Thus the case was settled: coffee was the less dangerous of the two beverages. The king, on the other hand, was murdered at a masked ball in 1792 at the age of 45 and became the subject of an opera by Verdi.

LARS BREIMER
Clinical lecturer

Department of Chemical Pathology and Human Metabolism,
Royal Free Hospital School of Medicine
London NW3 2QG

1 Stensvold I, Tverdal A, Jacobsen B. Cohort study of coffee intake and death from coronary heart disease over 12 years. *BMJ* 1996;312:544-5. (2 March.)

Care management

Care programme approach constitutes good management

EDITOR,—Max Marshall is correct in distinguishing between "standard" case management and "assertive community treatment."¹ Clearly, the benefits suggested by research on assertive community teams cannot be expected for routine community teams, which have lower ratios of staff to patients. This does not mean, however, that the care programme approach offers no benefits to patients and their relatives. It is hard to see how the components of the care programme approach can be regarded as anything but good practice for people with a mental illness. Surely patients should have their needs assessed and care plans recorded. And, surely, any care that is instigated should be reviewed and there should be one professional