Radcliffe Hospital and Oxfordshire Health Authorities decided to collaborate in producing their own card. This is now in wide use throughout the county, having replaced the national card not only at the Oxford Radcliffe Hospital but also, more importantly in numerical terms, in all general practice surgeries and community pharmacies. It has also been offered to the other trusts in the county to use if they wish.

The card was approved by the Royal Pharmaceutical Society before the substitution was made. The Department of Health was also given notice of our intentions and raised no objection.

A further concern arose about the need for guidance as to when the card should and should not be issued. A double sided, A4 sized advice sheet was therefore produced, using the best available evidence, and sent to all general practitioners and community pharmacists with the first batch of the new steroid cards. Part of its intention is to rationalise the issuing of the steroid cards. They are not, for instance, recommended for patients receiving low doses of inhaled corticosteroids, in whom there is no increased risk from chickenpox.

The same guidance sheet also includes locally tailored advice as to what should be done if a patient at risk who is receiving corticosteroids does come into contact with chickenpox or shingles.

JOHN REYNOLDS
Clinical lecturer in clinical pharmacology

University of Oxford,
Oxford OX3 9DU

TOM JONES
Primary care medical adviser
JULIE MADDEN
Pharmaceutical adviser

Oxfordshire Health,
Oxford OX3 7LG

SARAH SCHOFIELD
Formulary pharmacist

Oxford Radcliffe Hospital,
Oxford OX3 9DU

1 Feher MD, Simms JP, Lant AF. History of chickenpox and steroid cards: a new warning? *BMJ* 1996;312:542-3. (2 March.)

Doctors Reform Society in Australia defends its reputation

EDITOR,—Events since the publication of Simon Chapman's Focus article from Sydney have shown the shallow analysis of his report.¹ His attempt to tarnish the Doctors Reform Society by saying that it is obedient to a political party requires a response.

The society has always been concerned with a better health system and is jealous of its independent reputation and professionalism. In 23 years of medical politics it has stood firm against moves by both the main political parties to whittle away the universality and accessibility of Medicare. All political views are represented among current members of its executive, who are united in their commitment to the best ideals of medicine. The society has a fine reputation in Australia, and I hope that readers will not have been misled.

CON COSTA
National president

Doctors Reform Society,
Box 14, 4 Goulburn Street,
Sydney, NSW 2000,
Australia

1 Chapman S. Health versus personality in Australian politics. *BMJ* 1996;312:334. (10 February.)

Evaluation of a primary care anticoagulant clinic

Authors did not present enough data

EDITOR,—Sheena H Macgregor and colleagues' report on the evaluation of a primary care anticoagulant clinic managed by a pharmacist is

so lacking in methodological detail and supporting data that it is difficult to see how they can justify many of their conclusions.¹ No description of the study setting or study population is given. The potential for selection bias cannot therefore be determined, and it would be inappropriate to generalise the results beyond this particular general practice.

The authors chose to report only the proportion of the total number of measurements of the international normalised ratio that fell outside the target range, but it would be useful to know how many patients had abnormal ratios at any time during the study: did a small number of "difficult" patients contribute all the abnormal values or did most patients have an abnormal ratio at some time during the study? In addition, the subsequent course of those subjects with abnormal ratios is not reported.

No details are given as to how patients' knowledge was measured other than "by questionnaire," and no data are provided to substantiate the claim that counselling in hospital was unsatisfactory but subsequent counselling in the clinic improved patients' knowledge.

Despite the stated aims of the study, no costings are provided that might allow readers to assess the validity of the financial conclusion reached. The study cannot provide evidence that a pharmacist led clinic reduces the risks of toxicity and failure of treatment: it is not of an appropriate design to do so. Finally, only in the statement on funding does it become apparent that the study may have taken place in a fundholding practice. Given the nature of the study and the current climate of encouraging competition between health care providers, it is arguable that this amounts to a conflict of interest, although none is disclosed.

If the provision of health care in Britain is to improve, policymakers must surely be able to base decisions on evidence of a higher quality than this.

PETER W FOWLER
MRC training fellow in health services research

University of Dundee,
Ninewells Hospital and Medical School,
Dundee DD1 9SY

1 Macgregor SH, Hamley JG, Dunbar JA, Dodd TRP, Cromarty JA. Evaluation of a primary care anticoagulant clinic managed by a pharmacist. *BMJ* 1996;312:560. (2 March.)

Issue of quality control was not addressed

EDITOR,—Sheena H Macgregor and colleagues conclude that "good therapeutic control" and a "cost effective" anticoagulant service can be provided in the general practice surgery.¹ Near patient testing of the international normalised ratio is certainly an attractive means of providing an anticoagulant monitoring service. The authors' paper fails, however, to address the central issue of quality control and does not quantify the surgery costs.

Optimal patient care requires that near patient testing services meet the same standards of quality control that are required of accredited laboratories. Calibration of a coagulometer does not ensure that the measurements of the international normalised ratio produced on a day to day basis are either accurate or reproducible. The Joint Working Group on Quality Assurance² and the British Committee for Standards in Haematology³ recommend that near patient testing schemes should be subject to a regular, properly documented, quality control process administered in cooperation with an accredited laboratory. This is particularly important for those portable coagulometers (such as the Biotrack 512 used in the study) that analyse uncitrated blood and consequently cannot be entered into national external quality assurance

schemes. Without quality control data, comparison of international normalised ratios produced by different analytical systems, as in this study, cannot be meaningful.

Macgregor and colleagues refer only to the marginal costs of their clinic and do not quantify these. They draw a comparison with the hospital clinic's charge of £35. This is inappropriate, particularly as the hospital clinic has been available to fundholding general practitioners in Dundee free of charge.

Purchasers require detailed information about the quality, acceptability, and cost of different systems for monitoring anticoagulant treatment. The studies necessary to generate useful comparative data require cooperation rather than competition between primary and secondary care. In Dundee one such study involving four general practices and the haematology department is currently in progress.

SURJEET ADLAKHA
Staff grade haematologist
PHILIP G CACHIA
Consultant haematologist
MARTIN J PIPPARD
Professor of haematology

Department of Haematology,
Ninewells Hospital and Medical School,
Dundee DD1 9SY

PETER G DAVEY
Clinical director

Pharmacoeconomic Research Centre,
Ninewells Hospital and Medical School,
Dundee DD1 9SY

BARCLAY M GOUDIE
General practitioner

Westgate Health Centre,
Dundee DD2 4AD

- 1 Macgregor SH, Hamley JG, Dunbar J A, Dodd TRP, Cromarty J A. Evaluation of a primary care anticoagulant clinic managed by a pharmacist. *BMJ* 1996;312:560. (2 March.)
- 2 Joint Working Group on Quality Assurance. Appendix: guidelines on the control of near-patient tests and procedures performed on patients by non-pathology staff. In: Wood K, ed. *Standard haematology practice* 2. Oxford: Blackwell Science, 1994:278-9.
- 3 England JM, Hyde K, Lewis SM, Mackie IJ, Rowan RM. Guidelines for near patient testing: haematology. *Clin Lab Haematol* 1995;17:301-10.

Reporting of results should be standardised

EDITOR,—Sheena H Macgregor and colleagues report that primary care anticoagulant clinics run by a pharmacist achieve substantially better results than previously reported.¹ However, they give no indications of the patients' clinical conditions, no data on adverse events or quality of life, no information on how patients' preference was identified, and unhelpful data on costs.

A further major problem is their decision to report the proportion of measurements of the international normalised ratio that were within the target range as $\pm 10\%$ of the British Society for Haematology's guidelines. (We also have to assume that control was assessed against the desired range for the clinical indications for warfarin, although this is not stated.) This means that patients for whom the recommended reference range is 3.0-4.5 could have had actual values of between 2.7 and 4.95. This is a very wide therapeutic window, and it is therefore not surprising that 84-90% of patients achieved it. This study highlights the need for standardisation in reporting results from anti-coagulant clinics.

While such point prevalence data are of value, the percentage of time spent within the target range is a more discriminating assessment of therapeutic quality control.² One method of assessing the degree of therapeutic control, which we have reported, is to give the mean (± 1 SD) international normalised ratio for the clinic.³ Such analysis allows direct comparison between different environments—for example, primary care and hospital based clinics—and could be applied to the different therapeutic ref-

erence ranges. We therefore suggest that the problems of standardising the reporting of results for anticoagulation clinics would be reduced by use of the proportion of patients achieving therapeutic international normalised ratios, the percentage of time for which the ratios are within the therapeutic reference range, and the mean (± 1 SD) ratio.

The model of care reported by Macgregor and colleagues may have a place in anticoagulant treatment, but their small observational study cannot draw the broad conclusion that if we "are willing to devolve management to pharmacists then good therapeutic control is achievable in the surgery."

D A FITZMAURICE
Lecturer
F D R HOBBS
Professor

Department of General Practice,
University of Birmingham,
Medical School,
Birmingham B15 2TT

- 1 Macgregor SH, Hamley JG, Dunbar JA, Dodd TRP, Cromarty JA. Evaluation of a primary care anticoagulant clinic managed by a pharmacist. *BMJ* 1996;312:560. (2 March.)
- 2 Azar AJ, Dekers JW, Rosendaal FR, van Bergen PF, van de Meer FJ, Jonker JJ, et al. Assessment of therapeutic quality control in a long term anticoagulant trial in post myocardial infarction patients. *Thromb Haemostas* 1994;72:347-51.
- 3 Fitzmaurice DA, Hobbs FDR, Rose PE, Murray ET. A randomised controlled trial of oral anticoagulation management in general practice utilising computerised decision support (DSS) and near patient testing (NPT) with traditional management. *Fam Pract* 1995;12:253-4.

Authors' reply

EDITOR,—The lack of methodological detail, information on the patients studied, and data to support our conclusions are a consequence of publication as a short report. The full peer reviewed report of our study includes all these data.¹

The study included all patients prescribed anticoagulants, and hence all indications for warfarin. Of the 23 measurements of the international normalised ratio (in 12 patients) that were outside the target range, 15 were in patients recently discharged from hospital and not yet stabilised. Four occurred after interacting medicines were started (two during hospital admission and two by a general practitioner). Referral to the next clinic by the prescribing general practitioner avoided adverse effects, which is one advantage of monitoring the ratio at the point of prescribing. The remaining four cases were probably attributable to changes in cardiac status.

Patients' knowledge was assessed by structured interview when they first attended the surgery clinic. Altogether 24% did not know what action to take if they missed a dose of anticoagulant, 69% could not state signs of excessive ratios, 34% were unaware of the effects of alcohol, and 14% did not know that they should avoid medicines containing aspirin. After reinforcing information was given over three months the results were 0%, 14%, 7%, and 4% respectively.

Surjeet Adlakha and colleagues rightly point out the need for near patient testing to meet good quality control standards. Evaluation of the Biotrack has found good correlation with laboratory methods of evaluating the prothrombin time.² Contrary to the authors' comments that portable coagulometers cannot be entered into national external quality assurance schemes, it is possible to register with the schemes for quality assessment of tests that use a Boehringer-Mannheim coagulometer. The most important issue in assessing suitability is whether differences between results obtained by the instrument and by laboratory methods would alter the dose of warfarin prescribed. In this study random venous samples were submitted to the hospital laboratory. On no occasion would the

dose prescribed have been different had the laboratory result been used instead.

We agree with D A Fitzmaurice and F D R Hobbs that results from anticoagulant clinics should be standardised, provided the system can accommodate inappropriate changes of dose in stabilised patients with international normalised ratios marginally outside the target range. Adjusting the dose in these patients may be more detrimental than beneficial.

The management of chronic diseases should consider patients' convenience if possible. Several patients discharged from hospital have required home monitoring or initiation of treatment. Near patient testing facilitates this. Having up to date information on medical problems and treatment available in medical notes is also advantageous. Our patients asked to continue with management in the surgery when the practice became fundholding.

SHEENA H MACGREGOR
Clinical pharmacist
JAMES A DUNBAR
Senior partner

Downfield Surgery,
Dundee DD3 8NE

- 1 Macgregor SH. The development and implementation of a general practice based clinic to manage patients prescribed oral anticoagulants. Aberdeen: Robert Gordon University, 1995. (MSc thesis.)
- 2 Jennings I, Luddingstone RJ, Baglin T. Evaluation of the Ciba-Corning Biotrack 512 coagulation monitor for the control of anticoagulation. *J Clin Pathol* 1991;44:950-3.

GPs in Glasgow are in favour of primary care emergency centres

EDITOR,—Val Lattimer and colleagues ascertained views about the future provision of out of hours primary medical care from general practitioners who were members of two research networks; their response rate was 74% (n=199).¹ A similar survey was carried out among the 610 general practitioners registered with Greater Glasgow Health Board; the response rate of the doctors was also 74% (n=454), and that of the practices was 88%. The objective was to find out the extent to which proposals for developing primary care emergency centres were in tune with professional opinion and, therefore, the likelihood of their acceptance.

Of the general practitioners in Greater Glasgow who responded to the survey, 327 used a commercial deputising service out of hours. More than half of those who responded (238) thought that primary care emergency centres should be established but that home visits out of hours should continue for those requiring them. Most general practitioners (249) thought that primary care emergency centres would be an effective and efficient method of delivering out of hours care, although they were uncertain whether patients would find the service acceptable and thought that trivial calls might increase.

General practitioners over the age of 50 and female general practitioners were more likely to endorse these centres (83/130 v 159/322 (P<0.01) and 109/179 v 129/270 (P<0.01) respectively). This may be related to the finding that these two groups contained a high proportion of doctors who used a commercial deputising service almost all of the time or who had no out of hours commitment at all. Could this be a consequence of the pressures of home visiting (often in the dark) in a city like Glasgow? Is the job perceived as too dangerous and too stressful? These questions should be researched and debated as the results may affect the way in which out of hours care is delivered in the larger cities.

The general practitioners who responded to this study believed that primary care emergency centres should be situated locally (not necessarily in casualty departments) and that they should be staffed

by at least two doctors, one nurse, and one receptionist and should have a security guard. Altogether 198 general practitioners stated they would be prepared to work in these centres.

The results of this survey informed the decision to set up six primary care emergency centres in Glasgow covering 900 000 patients and over 90% of the city's general practitioners. These centres opened last March.

JAMES O'NEIL
General practitioner

Lightburn Medical Centre,
Glasgow G32 6NB

- 1 Lattimer V, Smith H, Hungin P, Gasper A, George S. Future provision of out of hours medical care: a survey with two general practitioner research networks. *BMJ* 1996;312:352-6. (10 February.)

Evidence-Based Medicine

Reviews may not be sufficiently critical of evidence

EDITOR,—The deficiencies of the approach taken by *Evidence-Based Medicine* in reviewing research papers, pointed out by NJ Pearson and colleagues,¹ are highlighted by the almost diametrically opposite conclusions reached in its review of a recent trial of interferon beta-1b² and in a more thorough review of the same evidence in the *Drug and Therapeutics Bulletin*.³

The review in the *Drug and Therapeutics Bulletin* critically examined the design of the trial⁴ and pointed out that the results were not analysed on an intention to treat basis and that this might have favoured interferon beta-1b. Also, the double blind nature of the design did not hold since the side effects of the drug identified those in the arm given active treatment. The review further made the point that this is not an idle criticism of the design since a fifth of the relapses (a primary outcome measure) were self reported. If these self reported relapses were excluded and the analysis confined to objectively verified cases of relapse then the difference between the groups became non-significant. None of these points are brought out in the commentary in *Evidence-Based Medicine*.

These criticisms may not vitiate the broad conclusions of the trial, but the difference in approach of the two publications led to very different bottom line statements. *Evidence-Based Medicine* concluded, "Physicians are advised to consider the use of [interferon beta-1b] in their patients with relapsing-remitting [multiple sclerosis]." The *Drug and Therapeutics Bulletin*, on the other hand, concluded, "In selected patients with [multiple sclerosis] interferon beta-1b possibly reduces the number and severity of clinical relapses. The drug is expensive and the evidence for its benefit is weak."

The moral of the story seems to be that even hard evidence may not be all that it is made out to be. Those who depend on reviews of published research rather than the original reports should be warned that such reviews may not always be sufficiently critical of the evidence.

JAMMI N RAO
Consultant in public health
J D MIDDLETON
Director of public health

Sandwell Health Authority,
West Bromwich,
West Midlands B70 9LD

- 1 Pearson NJ, Sarangi J, Fey R. Evidence-Based Medicine. *BMJ* 1996;312:380. (10 February.)
- 2 Absher JR. Interferon beta-1b reduced exacerbations in multiple sclerosis. *Evidence-Based Medicine* 1996;1:53.
- 3 Interferon beta-1b—hope or hype? *Drug Ther Bull* 1996;34:9-11.
- 4 IFNB Multiple Sclerosis Study Group, the University of British Columbia MS/MRI Analysis Group. Interferon beta-1b in the treatment of multiple sclerosis: final outcome of the randomized controlled trial. *Neurology* 1995;45:1277-85.