The only difference will be in the particular set of financial incentives that affect the people concerned. In a largely private, insurance based system such as that in the United States this may encourage inefficiency—that is, oversupply for some and no supply for others. Social insurance systems such as those in France and Germany may also be overresourced. Proposing other forms of finance is no escape from the fundamental issue. It merely alters the way in which the people concerned respond to inevitable scarcity.

In conclusion it is worth making clear that we do not propose any fundamental changes in the methods by which the NHS is financed. We support the continuation of a publicly financed NHS. However, we wish to promote an ongoing, open, and informed debate on how to make the hard choices about who should benefit from its limited resources.

Members of the Rationing Agenda Group are: Mr John Cairns, University of Aberdeen; Dr Nan Carle, King's Fund Management College, London; Dr Angela Coulter, King's Fund Development Centre, London; Professor Len Doyal, St Bartholomew's and the London Hospital School of Medicine and Dentistry, London; Professor John Gabbay, University of Southampton; Professor Raanan Gillon, Imperial College, London; Ms Heather Goodare, Horsham; Dr Iona Heath, Kentish Town Health Centre, London; Mr Chris Hegin-

botham, East and North Hertfordshire Health Authority, Welwyn Garden City; Mr John James, Kensington and Chelsea and Westminster Health Commissioning Agency, London; Professor Alison Kitson, Royal College of Nursing, London; Dr Robert Maxwell, King's Fund, London; Professor Alan Maynard, Nuffield Provincial Hospitals Trust, London; Professor Sheila McLean, University of Glasgow; Rabbi Julia Neuberger, London; Mr Bill New, King's Fund Policy Institute, London; Ms Marianne Rigge, College of Health, London; Mr Nick Ross, London; Professor Peter Rubin, Queen's Medical Centre, Nottingham; Dr Richard Smith, *BMJ*, London; Professor Albert Weale, University of Essex; Professor Alan Williams, University of York; Ms Karen Brown, London.

Funding: The King's Fund provided travel and accommodation funding for some members.

Conflict of interest: None.

- 1 Peters BG. *The politics of bureaucracy*. 3rd ed. London: Longman, 1989.
- 2 Harrison S, Hunter D. *Rationing health care*. London: Institute for Public Policy Research, 1994.
- 3 Doyal L. Needs, rights and equity: moral quality in healthcare rationing. *Quality in Health Care* 1995;4:273-83.
- 4 Maxwell R, ed. *Rationing health care*. London: Churchill Livingstone, 1995.
- 5 Walley T, Barton S. A purchaser perspective of managing new drugs: interferon beta as a case study. *BMJ* 1995;311:796-9.
- 6 Redmayne S, Klein R. Rationing in practice: the case of in vitro fertilisation. *BMJ* 1993;306:1521-4.

(Accepted 17 May 1996)

Lesson of the Week

Addison's disease presenting as reduced insulin requirement in insulin dependent diabetes

Lynne Armstrong, Patrick M Bell

Recurrent severe hypoglycaemia in patients with previously well controlled diabetes may be due to Addison's disease

Diagnosing Addison's disease requires a high degree of clinical suspicion since it is rare and its presenting symptoms are often non-specific. An association between Addison's disease and insulin dependent diabetes is well recognised. Whereas about 10-18% of patients with Addison's disease have insulin dependent diabetes, Addison's disease remains relatively rare among patients with insulin dependent diabetes.¹⁻³ We report two cases of patients with previously well controlled diabetes presenting with recurrent severe hypoglycaemia despite reduction in insulin doses.

Case 1

A 43 year old man with a 15 year history of insulin dependent diabetes well controlled with a stable regimen of twice daily injections of human insulin was referred to the diabetic clinic, Royal Victoria Hospital, complaining of increased frequency of hypoglycaemic episodes on waking. On one occasion he required intramuscular glucagon administered by his wife. On a second occasion he collapsed at work and required medical assistance. He also complained of tiredness and nausea. He was advised to reduce his evening dose of long acting insulin and was given an appointment for further review.

Before this date he was referred back by his general practitioner with further episodes of severe hypoglycaemia despite decreasing his insulin dose from a total of 30 units to 15 units. Tiredness, nausea, and vomiting had become more prominent, and he had noted a weight loss of about 6 kg in the preceding two weeks. Admission was arranged for stabilisation of diabetes and assessment of autonomic function in view of his apparent lack of hypoglycaemic awareness.

On admission he was noted to be pigmented, thin and dehydrated. Pigmentation was present particularly in the palmar creases and buccal mucosa. Postural hypotension was demonstrable. Investigations revealed serum concentrations of sodium 128 mmol/l, potassium 6.6 mmol/l, and urea 10.2 mmol/l. Primary adrenal failure was confirmed with a short synacthen test (0.25 µg intramuscularly)—plasma cortisol concentration was 91 nmol/l at baseline and 89 nmol/l at 30 minutes. Adrenocorticotrophic hormone concentration was 590 ng/l, and adrenal and gastric parietal cell antibodies were present. Antibodies to thyroglobulin and thyroid microsomes were not detected.

The patient rapidly recovered after treatment with intravenous fluids and hydrocortisone. His insulin requirements increased to 36 units daily. He remains well with replacement therapy of hydrocortisone 30 mg and fludrocortisone 0.1 mg daily.

Case 2

A 31 year old woman with insulin dependent diabetes diagnosed at two years of age presented at her routine diabetic clinic review complaining of increased hypoglycaemia despite repeatedly decreasing her insulin dose. She generally took a careful interest in her diabetes, with frequent monitoring of blood glucose and a multiple pen injection regimen resulting in levels of haemoglobin A_{1c} of 5.6-6.7% (normal <6%) over the previous two years.

She was noted to be pigmented, and her blood pressure was 130/90 mm Hg supine and 80/60 mm Hg standing. A short synacthen test (0.25 µg intramuscularly) carried out at the clinic confirmed the diagnosis of Addison's disease (plasma cortisol concentrations at baseline and 30 minutes both <40 mmol/l). Adrenal

Sir George E Clark
Metabolic Unit, Royal
Victoria Hospital, Belfast
BT12 6BA
Lynne Armstrong, research
fellow
Patrick M Bell, consultant
physician

Correspondence and reprint
requests to: Dr Bell.

BMJ 1996;312:1601-2

antibodies were present, but antibodies to thyroglobulin, thyroid microsomes, and gastric parietal cells were not detected. She was treated with oral hydrocortisone and fludrocortisone, and after adequate replacement her insulin requirements rose from 25 units to 35 units.

Discussion

The problems of hypoglycaemia in insulin dependent diabetes have received much attention.⁴ Loss of awareness of the classic symptoms of hypoglycaemia, sometimes associated with impaired autonomic nervous system and counter regulatory hormone responses to hypoglycaemia, may be caused in part by strict glycaemic control. Restoration of awareness of symptoms may follow careful avoidance of hypoglycaemia.⁵ Given the potential seriousness of severe hypoglycaemia, a natural clinical response is to urge careful monitoring of blood glucose and a reduction in insulin dose. The two cases described remind us to consider alternative diagnoses. Both patients presented to the same diabetologist eight years apart. The first was deteriorating with acute adrenal crisis before the diagnosis was made, whereas the second was diagnosed relatively quickly.

An association between autoimmune Addison's disease and insulin dependent diabetes is well recognised. Other features of the polyendocrine autoimmune syndrome may be present.¹ The prevalence of insulin dependent diabetes in series of patients with Addison's disease ranges from 10% to 18%.¹⁻³ By contrast, the prevalence of Addison's disease in patients with insulin dependent diabetes is lower, perhaps less than 1.9%, and is therefore more likely to be overlooked.⁶ Occasionally, both may present acutely with diabetic ketoacidosis provoking adrenal crisis.

In one series of patients who developed Addison's disease some time after insulin dependent diabetes, hypoglycaemia was a prominent presenting feature, although it usually merits only a brief mention in standard textbooks of diabetes and endocrinology.⁷ Addison's disease may present with spontaneous hypoglycaemia in the absence of diabetes mellitus.⁸ The cause of the hypoglycaemia in both situations is presumably increased sensitivity to insulin when the cortisol concentration is low or absent.

Reduced insulin requirements leading to hypoglycaemia in patients with previously well controlled diabetes can result from several causes, including the so called honeymoon phase of partial pancreatic recovery in the first year after diagnosis, changes in diet and exercise, and, usually in disease of long duration, reduced insulin clearance due to declining renal function. Increased insulin sensitivity due to adrenal insufficiency is a further cause which must not be overlooked.

- 1 Irvine WJ, Barnes EW. Adrenal insufficiency. *Clin Endocrinol Metab* 1972;1:549-94.
- 2 Maisey MN, Lessof MH. Addison's disease. A clinical study. *Guy's Hospital Reports* 1969;118:363-72.
- 3 Nerup J. Addison's disease. Clinical studies: a report of 108 cases. *Acta Endocrinologica* 1974;46:127-41.
- 4 Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long term complications in insulin dependent diabetes mellitus. *N Engl J Med* 1993;329:977-86.
- 5 Dagogo-Jack S, Rattarasarn C, Cryer PE. Reversal of hypoglycaemic awareness but not defective glucose counter-regulation in IDDM. *Diabetes* 1994;43:1426-34.
- 6 Riley WJ, Maclaren NK, Neufeld M. Adrenal autoantibodies and Addison's disease in insulin-dependent diabetes mellitus. *J Pediatr* 1980;97:191-5.
- 7 Beaven DW, Nelson DH, Renold AE, Thorn GW. Diabetes mellitus and Addison's disease. *N Engl J Med* 1959;261:443-54.
- 8 Dunlop D. Eighty-six cases of Addison's disease. *BMJ* 1963;iii:887-91.

(Accepted 22 March 1996)

A MIRACLE REMEMBERED

A very close shave

It was 29 August 1969. As a commander in the Biafran army, I had received intelligence that reinforcements for the Nigerian troops would be passing through the little town of Afugiri that night. I was ordered to use every means available to destroy the reinforcements. I selected a spot no more than 150 m from the main road leading through the town. Directly south of the spot the road traversed a hill. Traffic negotiating it would have to be very slow, greatly improving the chances of a successful strike. In the total darkness my men and I waited to spring our deadly ambush with mortars.

Close to midnight we heard the Nigerian military transport vehicles as they struggled up the incline. To our great surprise, they had their headlights on. It was therefore extremely easy to identify the individual vehicles. All I had to do was bring heavy fire down on to the target. With the mortars raining bombs on the convoy, the explosions came in mighty clusters, and the resulting fire illuminated the entire area for miles around. The illumination also revealed to the Nigerian sniper, no more than 70 m from my troops, who was the officer commanding the attack. He opened fire and hit me over my heart.

So great was the impact that I was knocked over. I just noted the small fire burning on my shirt as I realised that I had been killed in battle. I felt a profound sense of peace but nothing else. Then I heard the staff sergeant shouting to the men, "Let's remove the body before the vandals get here and chop it to pieces." The Biafrans always referred to the Nigerian troops as "vandals."

As my men began to drag me through the undergrowth so great was the pain in the left half of my

chest that I asked them to stop. There was utter consternation as the men, who had just seen me killed by the sniper, heard me speak—and even more so when we all saw a pile of bullets, still sparking flames. I had been hit over the heart by a burst of some 20 incendiary bullets and though they had set fire to my shirt and caused a huge haematoma not one of them had pierced my skin. Though I was glad to be alive, I was totally confused and it was only the pain that helped me to focus my mind on what was going on. By this time, the Nigerian survivors of my attack had begun to counterattack. My mortar command joined in to contain the counterattack.

The war ended five months later in January 1970. But for the entire period between that shooting and the end of the war, whenever a battle was joined, I noticed a definite aggregation of soldiers around me. It was only towards the final battles that my batman told me what had been happening. Word had got about that I had been hit by several bullets at close range, and that the bullets had not pierced my skin. The troops reasoned, therefore, that anyone near me during a battle would be unlikely to be injured. The death of a couple of my men who had been deployed at some distance during an attack only served to consolidate this belief.

Twenty five years on I still have no idea why not one of the 20 or so incendiary bullets, fired from only 70 m, had cut my skin. All I can think is that I was somehow protected by forces beyond anything I understand, for which I am grateful.—I KECHUKWU OBIALO AZUONYE is a consultant psychiatrist in Middlesex