Authors' reply

EDITOR,—Tom O'Connell reiterates our point that the differences in psychosocial outcomes in our study, although significant, were small. Consequently their importance is uncertain. We will soon begin a two year follow up of women in the antenatal care project to evaluate possible long term effects of reduced schedules of visits on maternal psychological health, the relationship between mother and child, and use of the health service postnatally. We hope that this study will help to clarify the clinical importance of the differences in psychosocial outcome that we found.

O'Connell also questions the generalisability of our findings, given our non-participation rate of 26%. Similar trials have reported non-participation rates of 33%¹ and 52%.² Non-participation may indicate women's dissatisfaction with reduced schedules; indeed, this is what we found from our questionnaire for non-participants. We advocate that the views of non-participants are sought in all trials in which there is considerable non-participation.

Sarah Vause and colleagues state that we failed to address the influence of parity on psychosocial outcomes and maternal satisfaction. While we were aware of the potential influence of parity, we would have required a much larger sample for subgroup analysis to study this. Unplanned post hoc comparison of subgroups is, of course, inappropriate. We have, however, investigated the influence of parity (using logistic regression); we found that it did not predict satisfaction.³ The fact that appreciably fewer nulliparous women took part in the trial suggests, however, that reduced schedules may be less appropriate for first time mothers.

We disagree with Vause and colleagues' view on measures of continuity. The absolute number of times that a woman sees her main carer is arguably a more appropriate index of whether she will have an opportunity to develop a relationship with that carer than is the proportion of visits in which she sees her main carer.

Like Douglas Tincello, we appreciate the importance of postnatal depression as a key psychosocial outcome of pregnancy. We reported data from the Edinburgh postnatal depression scale as means and SD because of the basic principle that statistical power is lost by dichotomising continuous variables. When we compared the proportions of women scoring above the threshold score of 12 we found no significant difference between the two groups, with 112 (13%) of 842 women in the intervention group scoring above 12 compared with 114 (14%) of 829 women in the control group ($\chi^2 = 0.039$, $df = 1$, $P = 0.844$).

We agree with Ruth Bell that further evaluation is needed of changes in the pattern of postnatal care that have the potential to affect psychosocial outcomes.

Kim Hinshaw and colleagues highlight a possible discrepancy between the indicators of success in *Changing Childbirth* and results of research on women's preferences. Their caveats about familiarity with a style of care are important. Nevertheless, we agree with them that the concept of woman centred care should be based on research evidence about women's preferences as well as their psychosocial and clinical needs.

JIM SIKORSKI SARAH CLEMENT
General practitioner Lecturer
JENNIFER WILSON
Honorary research midwife

Department of General Practice,
United Medical and Dental School,
London SE11 6SP

SARAH DAS
Project coordinator

National Maternity Record Project,
London NW1 5SD

NIGEL SMEETON
Lecturer in medical statistics

Department of Public Health Medicine,
United Medical and Dental School,
Guy's Hospital, London SE1 9RT

- 1 McDuffie RS, Beck A, Bischoff K, Cross J, Orleans M. Effect of frequency of prenatal care on perinatal outcome among low-risk women: a randomized controlled trial. *JAMA* 1996;275:847-51.
- 2 Binstock MA, Wolde-Tsadik G. Alternative prenatal care: impact of reduced visit frequency, focused visits and continuity of care. *J Reprod Med* 1995;40:507-12.
- 3 Clement S, Sikorski J, Wilson J, Das S, Smeeton N. Women's satisfaction with traditional and reduced antenatal visit schedules. *Midwifery* (in press).

Recent advances in ophthalmology are not solely technological

EDITOR,—Because they concentrate on technological advances in ophthalmology Hamish M A Towler and Susan Lightman fail to mention a recent advance in knowledge that could prevent or reduce blindness and that has been subjected to a rigorous randomised controlled clinical trial.¹ We refer to the results of the diabetes control and complications trial.² In this trial of intensive treatment for patients with insulin dependent diabetes tight glycaemic control reduced the risk of retinopathy developing by 76% in patients with no baseline retinopathy and slowed the progression of retinopathy by 54% in patients with mild baseline retinopathy. Results from a longitudinal observational study,³ a randomised controlled clinical trial,⁴ and expert consensus⁵ also suggest that improved glycaemic control will reduce microvascular complications (such as retinopathy) in people with non-insulin dependent diabetes.

Translating these results into daily medical practice poses several challenges, and many primary care practitioners do not yet understand the need to improve glycaemic control to a level that would achieve reductions in microvascular disease.⁶ Including these results as a recent advance in ophthalmology would be helpful.

ALAN D PENMAN
Medical epidemiologist
MICHAEL M ENGELGAU
Medical epidemiologist

Epidemiology and Statistics Branch,
Division of Diabetes Translation,
National Center for Chronic Disease Prevention and Health Promotion,
Centers for Disease Control and Prevention,
4770 Buford Highway, NE (Mailstop K-10),
Atlanta, GA 30341-3724,
USA

- 1 Towler HMA, Lightman S. Recent advances: ophthalmology. *BMJ* 1996;312:889-92. (6 April.)
- 2 Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977-86.
- 3 Klein R, Klein BE, Moss SE, Cruikshanks KJ. The Wisconsin epidemiologic study of diabetic retinopathy. XV. The long-term incidence of macular edema. *Ophthalmology* 1995;102:7-16.
- 4 Okhubo Y, Kishikawa H, Araki E, Miyata T, Isami S, Motoyoshi S, et al. Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized prospective 6-year study. *Diabetes Res Clin Pract* 1995;28:103-17.
- 5 Fifth Regenerstrief conference. Risks and benefits of intensive management in non-insulin-dependent diabetes mellitus. *Ann Intern Med* 1996;124:81-5.
- 6 Feder G, Griffiths C, Highton C, Eldridge S, Spence M, Southgate L. Do clinical guidelines introduced with practice based education improve care of asthmatic and diabetic patients? A randomised controlled trial in general practices in east London. *BMJ* 1995;311:1473-8. (2 December.)

Evidence based medicine

Rich sources of evidence are ignored

EDITOR,—It is reassuring that David L Sackett and colleagues have made two important concessions from their previous extreme position relating to evidence based medicine^{1,2}: firstly,

they have recognised the importance of clinical skill, as distinct from the consideration of scientific evidence; and, secondly, they have allowed other forms of evidence to take a step towards the hallowed position held by the randomised controlled trial.

A contradiction then ensues. Clinical skill is essentially derived from experience and is expressed as judgment in decision making.² But the variation in the ways in which different individuals interpret experience and formulate judgments renders this aspect difficult to expose to "big statistical ways of thinking."¹ It is therefore difficult to apportion scientifically the appropriate use of "best available clinical evidence"² in any particular decision process. We are thus led back or on to a wider view of medicine as a humane art that is supported by science.³ In this view the discipline of evidence based medicine is important for certain aspects of clinical decision making but is a relatively small part of the practice of medicine.

Forms of evidence allowable in evidence based medicine, while now extending beyond the randomised controlled trial, remain heavily numerate. This encourages emphasis on the quantifiable and physical aspects of any clinical dilemma, which may be inappropriate. Denial of social and psychological aspects may be detrimental, and ignoring the less readily measured dimensions may be dangerous. Rich sources of evidence also include the anecdotal, which are so often slated,⁴ and the qualitative, which is not mentioned by this lobby yet has developed considerably and provides illuminative results.⁵ The search for justification continues.

BLAIR H SMITH
Lecturer in general practice

Department of General Practice,
University of Aberdeen,
Foresterhill Health Centre,
Aberdeen AB9 2AY

- 1 Davidoff F, Haynes B, Sackett D, Smith R. Evidence based medicine. *BMJ* 1995;310:1085-6.
- 2 Sackett DL, Rosenberg WMC, Muir Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ* 1996;312:71-2. (13 January.)
- 3 Charlton BG. Holistic medicine or the humane doctor? *Br J Gen Pract* 1993;43:475-7.
- 4 Macnaughton J. Anecdotes and empiricism. *Br J Gen Pract* 1995;45:571-2.
- 5 Pope C, Mays N. Reaching the parts other methods cannot reach: an introduction to qualitative methods to health and health services research. *BMJ* 1995;311:42-5.

Scientific method and raw data should be considered

EDITOR,—Proponents of evidence based medicine claim that it is the conscientious, explicit, and judicious use of current best available external evidence about medical care.¹ What about scientific method and raw data? Scientific method—the fundamental intellectual activity of obtaining data, assessing them by a series of valid arguments, and obtaining a series of conclusions in descending order of likelihood in all subjects—does not seem to merit consideration.

Paradoxically, evidence based medicine seems to avoid all contact with the first hand evidence by replacing original findings with subjectively selected, arbitrarily summarised, laundered, and biased conclusions of indeterminate validity or completeness. It has been carried out by people of unknown ability, experience, and skills using methods whose opacity prevents assessment of the original data.

The new journal *Evidence-based Medicine* makes no reference in its first issue to methodology, experimental design, or statistics. No reason is given for the selection of the 24 papers that are published. No explanation of the sources and techniques of the added manipulations is included. No evidence of authorship