sample size would have given us 99.9% power to detect a difference in the rate of change in the scores obtained in the paired associate learning test of ≥ 0.1 point a year between treatments.

Although it is desirable to test multiple cognitive domains, there are practical limits to what can be achieved in an investigation that has been piggybacked on to a trial with its own demanding protocol. Expediency is necessary to retain the cooperation of the investigators and subjects in this important collaboration. Nevertheless, not all of C D Irvine and S H Irvine's criticisms of our outcome measures are justified. Their contentions that the paired associate learning test may not be a valid measure of intellectual function in late life and that it lacks internal consistency are not supported by the reference that they cite. In research in older populations the validity of the test is established as an indicator of decline on the abbreviated mental test score,¹ as a correlate of cognitive and functional decline as measured by the psychogeriatric assessment schedule,¹ and by psychiatrists' clinical ratings of the organic brain syndrome.² The narrow range of scores obtained in the paired associate learning test that we reported was not a function of the limited time available for learning. As Carroll observed, the speeded component of this test increases its variance.³

We acknowledge the presence of learning effects with the Reitan trail making test. We are not convinced, however, that they invalidate the test as a marker of cognitive change. The practice effect reflects a useful cognitive ability, impairment of which might suggest appreciable cognitive decline.

Finally, Irvine and Irvine observe that for those subjects who have stable test scores with time there will be no significant within subject correlation between the test score and time. This is true but irrelevant. The natural course of cognitive function in this age group is for it to decline in some subjects, to improve in others, and to remain relatively stable in many. There is therefore a continuum of change with time. This constitutes an intuitively reasonable, robust, and statistically conservative outcome measure for an investigation of group differences in cognitive functioning over time. A more detailed justification and description of this approach are given in a recent publication.⁴

MARTIN PRINCE
Lecturer in psychiatry
ANTHONY MANN

Professor of epidemiological psychiatry

Institute of Psychiatry,
London SE5 8AF

ROBERT BLIZARD
Lecturer in statistics
ANNE BIRD
Consultant psychiatrist

Royal Free Hospital,
London NW3 2QG

- 1 Little A, Hemsley D, Bergmann K, Volans J, Levy R. Comparison of the sensitivity of three instruments for the detection of cognitive declines in elderly living at home. *Br J Psychiatry* 1987;150:808-14.
- 2 Kuriansky J, Gurland B, Cowan D. The usefulness of a psychological test battery. *Int J Aging Hum Dev* 1976;7:331-42.
- 3 Carroll JB. *Human cognitive abilities*. New York: Cambridge University Press, 1993.
- 4 Prince M, Lewis G, Bird A, Blizard R, Mann A. A longitudinal study of factors predicting change in cognitive test scores over time, in an older hypertensive population. *Psychol Med* 1996;26:555-68.

Reduced schedule of antenatal visits

Differences in psychosocial outcomes between groups were not large

EDITOR,—Jim Sikorski and colleagues' paper comparing two different schedules of antenatal visits

highlights the wide variation in practices within the European Union.¹ The average number of antenatal visits is five in Luxembourg; seven in Belgium, France, and Italy; and 10 in Germany and Denmark.² The authors' main conclusions are that a reduced schedule of antenatal visits may lead to reduced psychosocial effectiveness and dissatisfaction with the frequency of visits.

The study was conducted in three inner city locations in London, and roughly 30% of the participants in each group were drawn from eight or more ethnic minorities. The non-participation rate was 26.3%, and among those who agreed to participate the non-response rate to the questionnaire asking about patients' satisfaction ranged from roughly 30% to 37%. Thus care must be exercised in assessing the applicability of these results to the wider population of Britain.

The authors' conclusion that the psychosocial outcome was poorer in the group for whom the schedule of visits was reduced is partially based on the use of the Cambridge worry scale and a scale scoring positive and negative attitudes towards the fetus and baby. While the women in the group for whom the schedule of visits was reduced had a significantly higher worry score for two of 16 variables examined ($P < 0.01$), the actual mean differences were small: with the possible score ranging from 0 to 5, the mean values for whether the baby was thriving differed by only 0.2 between the two groups and those for whether the mother was coping differed by only 0.1. Similarly, while the score for a negative attitude towards the fetus was significantly higher in the women in the group for whom the schedule of visits was reduced ($P < 0.05$), the actual difference was small: with a range of possible scores of 0-21, the mean difference between the groups was only 0.6.

Thus readers must bear in mind the special nature of the study population used and the fact that the differences in psychosocial outcomes between the two groups were not particularly large.

TOM O'CONNELL
Postgraduate student

Department of Public Health Medicine,
University College Dublin,
Dublin 2,
Republic of Ireland

- 1 Sikorski J, Wilson J, Clement S, Das S, Smeeton N. A randomised controlled trial comparing two schedules of antenatal visits: the antenatal care project. *BMJ* 1996;312:546-53. (2 March.)
- 2 Blondel B, Pusch D, Schmidt E. Some characteristics of antenatal care in European countries. *Br J Obstet Gynaecol* 1985;92:565-8.

Study ignored influence of parity on women's needs

EDITOR,—We wish to make two comments about Jim Sikorski and colleagues' trial comparing two schedules of antenatal visits.¹ Firstly, with regard to continuity of carer we think that in any study in which the mean number of visits differs significantly between the two groups, variables such as those suggested by Binstock and Wolde-Tsadik should be used to take account of this difference.² If the number of visits is reduced a woman will probably see fewer carers. Therefore the proportion of care provided by the main carer (continuity index) and the number of carers divided by the number of visits (discontinuity index) would be more meaningful measures. Secondly, the authors fail to address the question of whether the reduced schedule of visits was less acceptable to primiparous than multiparous women even though they note that fewer primiparous women agreed to be recruited and that one of the most common reasons for non-participation was that the women did not want fewer antenatal visits.

Although the authors report that the professionals thought that traditional care met the

non-clinical needs of nulliparous women better than new style care, they do not report whether satisfaction and dissatisfaction differed between multiparous and primiparous women. Even if there was no difference in satisfaction this would be an important finding as it would be contrary to the view of the professionals surveyed. It would also be interesting to know whether non-adherence to the proposed schedule of visits was related to parity—for example, did a larger proportion of primiparous women in the group for whom the schedule of visits was reduced book extra visits and a larger proportion of multiparous women in the group receiving traditional care miss visits?

Women in the new style care group were significantly more likely to choose the same type of care in a future pregnancy. Primiparous women in this study would be multigravidas in a future pregnancy, and this might affect their choice of care in a future pregnancy. Again, it would be important to know whether women viewed the reduced schedule of visits as appropriate for a second or subsequent pregnancy but not for the first.

The authors seem to have disregarded the influence of parity on a woman's needs for education and reassurance in pregnancy. This is unfortunate as the issue of a schedule of antenatal visits that meets both the clinical and the psychosocial needs of women merits urgent consideration.

SARAH VAUSE
Research fellow
MICHAEL MARESH
Director
KHALED KHALED
Research fellow

Royal College of Obstetricians and Gynaecologists Audit Unit,
St Mary's Hospital,
Manchester M13 0JH

- 1 Sikorski J, Wilson J, Clement S, Das S, Smeeton N. A randomised controlled trial comparing two schedules of antenatal visits: the antenatal care project. *BMJ* 1996;312:546-53. (2 March.)
- 2 Binstock MA, Wolde-Tsadik G. Alternative prenatal care: impact of reduced visit frequency, focused visits and continuity of care. *J Reprod Med* 1995;40:507-12.

Effect of fewer visits on postnatal depression is important

EDITOR,—Jim Sikorski and colleagues have conducted a large and detailed trial of the benefits and problems associated with different patterns of antenatal care.¹ Such studies, examining outcomes such as satisfaction with care, anxieties about the health of the fetus, and the ability of new parents to cope with their baby, are essential to allow proper evaluation of the recommendations in *Changing Childbirth*, the report of an expert maternity group.² The authors found that women in the group for whom the schedule of visits was reduced had significantly lower levels of satisfaction, thought that they did not have enough time to share concerns about the pregnancy, and were more worried about their ability to cope postnatally than women in the group who received traditional care. This must make us rethink our attitude to the new style of antenatal care, particularly as many of us have been swayed by the evidence that the pattern of care does not affect important physical outcomes such as severe pre-eclampsia.

Postnatal depression is the commonest complication of pregnancy, affecting up to a tenth of women, and in some cases carries an appreciable risk of suicide and infanticide.³ It may be precipitated by feelings of isolation during pregnancy and anxieties in the puerperium. Sikorski and colleagues' data are potentially of great interest in terms of the effect on the incidence of depression but are not presented in a way that

shows this adequately. The Edinburgh postnatal depression scale is a validated tool for screening for psychiatric morbidity, which is indicated by a score of ≥ 12 or 13.⁴ Presenting these data in the form of a mean and SD does little other than describe the range of emotional and mental states of the sample. To show the incidence of postnatal depression it would have been better to include the number or percentage of patients in each group who exceeded the threshold score and to analyse the significance of any difference between these. As the paper stands, we are left with the information that women randomised to fewer antenatal visits were more anxious, worried, and likely to have negative feeling about their babies than women randomised to traditional care, but we do not have sufficient information to determine whether more of these women are at risk of postnatal depression.

DOUGLAS TINCELLO

Specialist registrar in obstetrics and gynaecology

Liverpool Women's Hospital,
Liverpool L8 7SS

- 1 Sikorski J, Wilson J, Clement S, Das S, Smeeton N. A randomised controlled trial comparing two schedules of antenatal visits: the antenatal care project. *BMJ* 1996;312:546-53. (2 March.)
- 2 Department of Health. *Changing childbirth: report of the expert maternity group*. London: HMSO, 1993.
- 3 Henshaw CA, Cox JL. Postnatal depression. *Current Obstetrics and Gynecology* 1995;5:70-4.
- 4 Cox JL, Holden JM, Sagavosky R. Detection of postnatal depression: development of the 10-item Edinburgh postnatal depression scale. *Br J Psychiatry* 1987;150:782-6.

Postnatal care also requires evaluation

EDITOR,—Jim Sikorski and colleagues report that reducing the number of scheduled antenatal visits may worsen psychosocial outcomes and increase dissatisfaction with care.¹ Postnatal care, while not being a specific focus of the policy initiative outlined in *Changing Childbirth*,² has also changed recently, with increasingly short lengths of stay and a move away from daily visits by midwives.

I investigated the use of maternity services in rural Northumberland, comparing women resident in two local government districts in which isolated general practitioner maternity units were situated with those resident in a district much closer to the consultant obstetric unit that served as the main referral unit for these isolated units. Information on a range of demographic and clinical variables was obtained from routine information systems and medical records for all singleton deliveries to women resident in these areas in 1988. The most striking difference between the two groups was the difference in the length of postnatal stay; the women resident in the districts with the general practitioner units spent significantly longer in hospital after delivery, and this did not seem to be accounted for by differences in clinical variables (table 1). The general practitioner units were used extensively for postnatal convalescence by local women who had been discharged soon after delivery from the consultant unit.

I did not investigate postnatal complications in detail or ask for women's views of their care, and there are several potential explanations for the increased length of stay in the areas with general practitioner units. Possibly, however, women who are in a small, comfortable local unit with relatively little pressure on beds are less enthusiastic to return home soon after delivery. Certainly, support from a midwife may be less readily available for these women after discharge.

Increasingly, plans for new maternity units suggest a move towards American-style "labour, delivery, and recovery" rooms, where postnatal stay is expected to be a matter of hours. In the community there is a trend towards fewer scheduled visits, as with antenatal care. If reduced care before delivery may worsen psychosocial out-

Table 1—Details of singleton deliveries in hospital to mothers resident in three local government districts in rural Northumberland in 1988. Figures are numbers (percentages)

	Districts A and B	District C
Singleton deliveries	582	479
Deliveries in general practitioner maternity unit	232 (39.9)	3 (0.6)
Deliveries in consultant unit	350 (60.1)	476 (99.4)
Primiparous women	262 (45.0)	190 (39.7)
Caesarean sections	67 (11.5)	62 (12.9)
Assisted vertex deliveries	69 (11.9)	36 (7.5)*
Babies admitted to SCBU	57 (9.8)	53 (11.1)
Women staying in hospital <48 h after delivery	95 (16.3)	144 (30.1)**
Women staying >5 days after delivery	178 (30.6)	88 (18.4)**

SCBU = Special care baby unit.
 χ^2 test: * $P < 0.05$, ** $P < 0.01$.

comes then the same may hold for care after delivery, when women may need support with breast feeding and with adjusting to the arrival of the baby. Changes to postnatal care should be rigorously evaluated by randomised controlled trials to assess their impact on clinical and non-clinical variables.

RUTH BELL

Registrar in public health medicine

County Durham Health Commission,
Durham DH1 5XZ

- 1 Sikorski J, Wilson J, Clement S, Das S, Smeeton N. A randomised controlled trial comparing two schedules of antenatal visits: the antenatal care project. *BMJ* 1996;312:546-53. (2 March.)
- 2 Department of Health. *Changing childbirth: report of the expert maternity group*. London: HMSO, 1993.

Attention should be paid to what women want

EDITOR,—Jim Neilson highlights the significant increase in maternal anxiety that may be provoked by reducing the number of routine antenatal visits.^{1,2} This reduction is one of 10 "key indicators of success" listed in the report *Changing Childbirth*,³ which must be implemented by 1998.⁴ Maternity services should become more "woman centred," with balanced input from midwives, general practitioners, and obstetricians, according to each woman's needs. Some of the key indicators should increase women's feeling of involvement in their pregnancy care, but others seem directive and inflexible and have little supportive evidence at either national or local level.

We surveyed 332 unselected women (beyond 34 weeks of pregnancy) attending our hospital antenatal clinic; 40% were nulliparous and 60% multiparous. The hospital serves an urban popu-

lation in north east England and has 4000 deliveries a year. A consultant led, shared care system is used, with women at "low risk" attending the hospital at booking and 34, 38 (nulliparous women only), and 41 weeks.

Jim Sikorski and colleagues report that women in a group for whom the schedule of antenatal visits was reduced were significantly more anxious, had more negative attitudes, and were more dissatisfied with the number of visits than women who had traditional care, although outcomes of pregnancy were no different.² We asked our subjects specifically about visits to hospital clinics. Altogether 287 thought that the number was "just right" (table 1). The attitudes of nulliparous and multiparous women were not significantly different ($P = 0.19$, χ^2 test). Two hundred and ninety two women thought that visits to a hospital antenatal clinic were necessary in an uncomplicated pregnancy (table 1); 61 thought that they gave "reassurance/peace of mind" and 113 that they confirmed that "everything was all right." Such data are open to criticism, as most women will have experienced only one style of care and are satisfied with whatever care they receive.⁵ We think, however, that our results support Sikorski and colleagues' findings. Before the number of hospital visits is reduced further, perhaps we should acknowledge that many women value hospital care, even in uncomplicated pregnancies.

Another key indicator states that "at least 30% of women should have the midwife as the lead professional."³ We asked the women which of five styles of antenatal care they would prefer in an uncomplicated pregnancy; table 1 shows their preferences. Again, we accept that most women may have chosen the third option because of familiarity. We support the principle of midwifery led care for women who prefer it, but why should an arbitrary figure of 30% be imposed? Is this really moving towards woman centred care?

KIM HINSHAW

Consultant obstetrician

AMR EL-HORISHY

Senior house officer in obstetrics and gynaecology

SUZANNE BATES

Audit officer

City Hospitals Sunderland NHS Trust,
Sunderland SR4 7TP

- 1 Neilson J. Antenatal care on trial. *BMJ* 1996;312:524-5. (2 March.)
- 2 Sikorski J, Wilson J, Clement S, Das S, Smeeton N. A randomised controlled trial comparing two schedules of antenatal visits: the antenatal care project. *BMJ* 1996;312:546-53. (2 March.)
- 3 Expert Maternity Group. *Changing childbirth: report of the expert maternity group*. London: HMSO, 1993.
- 4 Moores Y, Winyard G. *Woman-centred maternity services*. Leeds: NHS Management Executive, 1994. (EL(94)9.)
- 5 Tucker JS, Hall MH, Howie PW, Reid ME, Barbour RS, Florey CduV, et al. Should obstetricians see women with normal pregnancies? A multicentre randomised controlled trial of routine antenatal care by general practitioners and midwives compared with shared care led by obstetricians. *BMJ* 1996;312:554-9. (2 March.)

Table 1—Women's responses to questions seeking their opinion on antenatal care

	No
Opinion on No of visits to hospital clinic (n = 325):	
Just right	287
Too few	37
Too many	1
Opinion on whether it is necessary to go to hospital antenatal clinic for uncomplicated pregnancy (n = 328):	
Yes	292
No	10
Not sure	26
Style of antenatal care preferred in uncomplicated pregnancy (n = 317):	
All care with community midwife (who would refer you to your GP or hospital if there were any problems)	33
All care with community midwife and GP (who would refer you to hospital if there were any problems)	51
Most care with GP and community midwife, with some visits to hospital clinic	209
All care with GP (who would refer you to hospital if there were any problems)	2
All care at hospital clinic	22

GP = General practitioner.