has a protective function. It protects the vulnerable from misinformation due to mistakes by or the ignorance of the informer, from pressure by those with malintent, from economically driven judgments on their future, and from much more. It also protects us, as doctors, from ourselves: our ignorance or arrogance, any temptation to cover up medical mistakes, our difficulty in asking for help from a colleague, overinvolvement with a patient that colours our judgment, our fatigue, or personal prejudice or bias about clinical or social conditions. It protects us from undue pressure by relatives weary of caring or who stand to gain financially. Managers cannot put pressure on us to clear those who are dying from our beds rapidly, and purchasers cannot question why we strive to provide quality care to patients with a poor prognosis.

As a pathologist Kevorkian may be desensitized to corpses. We provide long term care and bereavement support and are increasingly aware of the absolute import of death. Currently, prognosis cannot be predicted accurately, there are errors of diagnosis, depression is difficult to diagnose in medically ill people, patients' priorities alter often during the course of a life threatening illness, hope can re-emerge from hopelessness, we find some patients' problems overwhelming at times, and sometimes our judgment is clouded by ignorance or fatigue. Why no cries to enshrine in law the right of all patients to a second opinion if their suffering remains intractable for a week? Why call for legalising carelessness?

ILORA FINLAY (senior lecturer in palliative medicine),
PHILIP ROUTLEDGE (professor of therapeutics),
ANDREW FREEDMA (consultant in HIV and infectious diseases), KEN WOODHOUSE (professor of geriatric medicine), D P DAVIES (professor of child health),
A B HAWTHORNE (consultant gastroenterologist),
MIKE PRITCHARD (consultant rheumatologist),
MOLLY HALL (consultant general physician and rheumatologist), PETER BECK (consultant physician),
CLARE WILKINSON (senior lecturer in general practice)

University of Wales College of Medicine,
Cardiff CF4 4XN

1 Roberts J, Kjellstrand C. Jack Kevorkian: a medical hero. *BMJ* 1996;312:1434. (8 June.)

His actions are the antithesis of heroism

EDITOR,—John Roberts and Carl Kjellstrand are entitled to their view that Jack Kevorkian is a medical hero.¹ They should, however, be aware that those who are fortunate enough to have avoided coming into contact with him do not share their view.

Kevorkian is famous for taking at their word those who are sick or disabled who say that they want to die, and of "helping" by killing them. His defence has always been that his aim is to kill their pain and suffering and that the death of the organism is an unfortunate side effect of this laudable intention. Where this value system falls down is in its assumption that death is the best—indeed the only—remedy for intractable suffering and that sick and disabled people are right to want to die while able bodied people are inherently wrong to want to die, even though people in both groups may request death equally fervently and for much the same reason.

Roberts and Kjellstrand are right to say that the medical profession must say "enough" to pain and suffering. The point they miss is that there are ways of saying this that do not entail killing the patient.

I might well once have sought out Kevorkian's "services." I am severely disabled, and some years ago it was thought that my life expectancy was severely reduced. Additionally, I was (and still am) suffering great pain, and several unconnected factors combined to make me decide that

I wanted to die—a wish that lasted many years. I would have satisfied all the "strict criteria" proposed by the voluntary euthanasia lobby, let alone the much more lax standards set by Kevorkian himself. Had he been practising in my vicinity, I would quite possibly have availed myself of Kevorkian's services and thus have been denied the chance to see again the beauty of life, albeit a life still restricted both by my disability and by severe pain.

I suggest that the real heroes of sick and disabled people are those who give of themselves; who stay with us, hold our hands, and, when the best efforts of modern medicine fail, say that they will not desert us. People who are sick and feel hopeless need the very best that medicine, in its widest sense, can offer. What they do not need is to be told, "Yes, you are right; death is the only answer to your problems." That, I suggest, is the ultimate desertion and the antithesis of heroism.

ALISON DAVIS

35 Stileham Bank,
Milborne St Andrew,
Blandford Forum,
Dorset DT11 0LE

1 Roberts J, Kjellstrand C. Jack Kevorkian: a medical hero. *BMJ* 1996;312:1434. (8 June.)

Editorial's objectivity is in doubt

EDITOR,—It is interesting that nowhere do John Roberts and Carl Kjellstrand explicitly support what Jack Kevorkian is doing; indeed, they record "actions that most of us find dubious" and say that "to be a hero does not mean being right."¹ They do, however, give explicit support for why he acts thus, and many readers will interpret this as implicit approval of his actions. This subjective relativism is a betrayal of the academic objectivity we expect and deserve from the *BMJ*. There has been recent, rational discussion of all the issues in Britain, and the profession and parliament have overwhelmingly rejected euthanasia.^{2,3}

Another commentary adds to the catalogue of Kevorkian's dubious actions: "For his next trick, Dr Kevorkian will assist at a suicide and then, with the prior consent of the deceased and the appropriate medical tests, his or her organs will be removed soon after death for use in transplant surgery" and "He courted controversy early in his career with his suggestion that death-row prisoners could be used for medical experimentation just prior to death and that organs be harvested from executed criminals." The omission of these relevant facts from their editorial casts further doubt on the authors' objectivity.

ANDREW FERGUSSON
Chairman

Healthcare Opposed to Euthanasia,
PO Box 10550,
London N1 1NN

1 Roberts J, Kjellstrand C. Jack Kevorkian: a medical hero. *BMJ* 1996;312:1434. (8 June.)

2 Beecham L. Legalised euthanasia is categorically rejected. *BMJ* 1993;307:132.

3 Select Committee on Medical Ethics. *Report*. London: HMSO, 1994. (Para 237.)

4 Hunt L. Can Dr Death be a true hero? *Independent* 1996 Jun 8:19.

Doctors should indeed cry, "Enough"

EDITOR,—Thomas Carlyle wrote, "The Hero can be Poet, Prophet, King, Priest or what you will."¹ John Roberts and Carl Kjellstrand choose Jack Kevorkian as their hero.² How do they square this choice—which is akin to choosing Barabbas—with the Hippocratic Oath or the Geneva Declaration, in which doctors promise to maintain the utmost respect for human life? Dr Everett Koop, a former surgeon general in the

United States, predicted that such choices would be made before the century was out when he wrote in 1980 that practices once labelled unthinkable would be considered acceptable. He went on to plead: "Let it never be said by historians in the latter days of this century that there was no outcry from the medical profession. Let it never be said that a euthanasia programme for various categories of citizens could never have come about if physicians had stood for the moral integrity that recognises the worth of every human life."³

Is it not time for us as a profession to decry this form of hero worship and indeed cry "enough" of this perverse destruction of the principles of our professional founders?

ROBERT P BALFOUR

Consultant obstetrician and gynaecologist

Princess of Wales Hospital,
Bridgend CP31 1RQ

1 Goldberg MK, Bratin GJ, Engel M. *On heroes, hero worship, and the heroic in history*. Oxford: Oxford University Press, 1993.

2 Roberts J, Kjellstrand C. Jack Kevorkian: a medical hero. *BMJ* 1996;312:1434. (8 June.)

3 Schaeffer FA, Everett Koop C. *Whatever happened to the human race?* London: Marshall Morgan and Scott, 1980.

*We received 17 other letters about the editorial, all of which expressed views similar to those published here.—EDITOR

Haem iron intake in young children

Other health promotion activities would have higher priority in Africa

EDITOR,—As a result of their study of haem iron intake and serum ferritin concentration in children aged 12–36 months in Australia, Michael Mira and colleagues urge that lean meat be introduced at 6–9 months and state the amount of meat that will give a daily intake of haem iron of 0.71 mg/day.¹

We have three questions. Firstly, what is the experience of vegetarians, who have better health than omnivorous eaters? Are their young children likely to be detectably disadvantaged? Vegetarian women, described as being in good health, have low ferritin concentrations (mean 13.6 µg/l).² Could a low ferritin concentration have a different connotation with regard to health in different contexts?

Secondly, what are the implications for infants in the Third World, especially those in impoverished Africa? The regimen suggested by the authors is almost impossible there, especially in high parity families, because in most populations meat is eaten at most once or twice a week. The alternative of general prophylaxis with iron is far beyond the means of the masses. Since in such populations there are so many other adverse factors, dietary and non-dietary, would the particular drawback of low ferritin concentrations be likely to be clinically discernible? In an African village would the group in the lowest quartile of ferritin concentration be at a demonstrable disadvantage if compared with the group in the highest quartile, apart from in areas where malaria and hookworm are endemic?

Thirdly, what is the magnitude in young children of disabilities linked with low ferritin concentrations? Much in this field remains unclear.³ Mira and colleagues refer to studies of very young children. In that undertaken in Chile the scores on the mental development index in the contrasting groups differed by 6%.⁴ In the oft quoted Costa Rican study the Woodcock-Johnson scores in the anaemic and non-anaemic groups differed by 1%.⁵ While other reported

differences are slightly larger, the disadvantages of those with low iron status are not major.

In Africa the state has limited financial and other resources for maintaining health and combating disease. In the absence of prospective studies, a widespread improvement of iron status would seem to have lower priority than many other health promotion endeavours, such as water supply, immunisation of children, and 24 hour availability of staff at clinics.

ALEXANDER R P WALKER
Head

Human Biochemistry Research Unit,
South African Institute for Medical Research,
PO Box 1038,
Johannesburg,
South Africa

DEMETRE LABADARIOS
Head

Department of Human Nutrition,
Faculty of Medicine,
University of Stellenbosch,
PO Box 19063,
Tygerberg,
South Africa

- 1 Mira M, Alperstein G, Karr M, Ranmuthugala G, Causier J, Niec A, *et al*. Haem iron intake in 12-36 month old children depleted in iron: case-control study. *BMJ* 1996;312:881-3. (6 April.)
- 2 Alexander D, Ball MJ, Mann J. Nutrient intake and haematological status of vegetarians and age-sex matched omnivores. *Eur J Clin Nutr* 1994;48:534-46.
- 3 Beard J. One person's view of iron deficiency, development, and cognitive function. *Am J Clin Nutr* 1995;62:709-10.
- 4 Walter T, de Andrada I, Chadud P, Perales CG. Iron deficiency anemia: adverse effects on infant psychomotor development. *Pediatrics* 1989;84:7-17.
- 5 Lozoff B, Jimenez EE, Wolf AW. Long-term developmental outcome of infants with iron deficiency. *N Engl J Med* 1991;325:687-94.

Authors' reply

EDITOR,—Alexander R P Walker and Demetre Labadarios's first question is whether children of vegetarians are likely to be disadvantaged in terms of haem iron intake. In a large community prevalence study that we have recently completed we found no significant association between vegetarianism and iron depletion.¹ This is probably because, in Australia, vegetarianism is a matter of choice rather than a cultural dictate and vegetarians are usually well informed about dietary issues such as iron intake. The authors' comparison of vegetarian women and toddlers, in whom the central nervous system is still developing, may not be appropriate. A wealth of literature suggests that, among other effects, iron deficiency in young children impairs cognitive development² and immunity³ and enhances absorption of lead from the environment.⁴

Australia has a high standard of living, a comprehensive social security system, and a well developed system of primary care. A safe water supply is guaranteed, levels of sanitation are high, and a wide range of foods is available at moderate cost. Conclusions drawn from data collected in a country such as Australia may not be applicable to the developing world; for this reason we described in our article the context in which our study took place.

In answer to the question whether the particular drawback of low ferritin concentrations would be clinically discernible, the answer is no. Firstly, it is iron deficiency anaemia rather than iron depletion that is associated with permanently impaired cognitive development. Secondly, the effects of iron deficiency anaemia on cognitive ability are too small to be detected in any individual child but in Australia are important on a population basis. The same is likely to be true for the other effects of iron deficiency. Whether these effects are demonstrable on a population basis in Africa requires research.

Ideally, decisions on the allocation of resources and priorities in any country should be based on an active and informed dialogue

between public health officials and the community. In Africa the water supply, sanitation, availability of food, immunisation, breast feeding, education (especially of women), and political stability are likely to be of higher priority in promoting health and cognitive development than is the isolated problem of iron deficiency.

MICHAEL MIRA
Director

MARGARET KARR
Research officer

Division of General Practice,
Central Sydney Area Health Service,
Balmain NSW 2041,
Australia

GARTH ALPERSTEIN
Area community paediatrician

Community Health Services
Central Sydney Area Health Service,
Camperdown, NSW 2050,
Australia

- 1 Karr M, Alperstein G, Causier J, Mira M, Lammi, Fett MJ. Iron status and anaemia in preschool children in Sydney. *Aust J Public Health* (in press).
- 2 Lozoff B, Jimenez E, Wolf AW. Long-term developmental outcome of infants with iron deficiency. *N Engl J Med* 1991;323:687-94.
- 3 Thibault H, Galan P, Selz F, Preziosi P, Olivier C, Badoual J, *et al*. The immune response in iron-deficient young children: effect of iron supplementation on cell-mediated immunity. *Eur J Pediatr* 1993;152:120-4.
- 4 Mahassey KR. Nutritional factors in lead. *Nutr Rev* 1981;39:353-62.

Tea flavonoids have little short term impact on serum antioxidant activity

EDITOR,—There is currently much interest in the possible benefits of dietary flavonoids in preventing cardiovascular diseases.^{1,2} The apparent protective effect has been attributed to the potent antioxidant activity of flavonoids. In vitro flavonoids are powerful inhibitors of the oxidation of low density lipoproteins, a process that has been strongly implicated in the development of atherosclerosis.³ For some years we have been using enhanced chemiluminescence as a simple method of measuring antioxidant activity in biological and other fluids.⁴ Since black tea may account for over half of the flavonoid intake in Western diets we recently investigated the antioxidant activity of a variety of black teas as well as their potential impact on antioxidant status in vivo.

We prepared eight samples of black tea by adding 0.5 g of tea leaf (from eight popular commercially available brands) to 25 ml of boiling water, which was regularly agitated for three minutes. The mean antioxidant activity in the resulting solutions was 8477 $\mu\text{mol/l}$ (range 4275-12 110 $\mu\text{mol/l}$), which confirms that tea at typically consumed concentrations has powerful antioxidant properties in vitro. These values compare with typical serum antioxidant activities of 350-550 $\mu\text{mol/l}$.⁴ It therefore seemed likely that ingestion of tea would have a considerable impact on serum antioxidant activity in vivo. We investigated the impact of drinking 500 ml of English breakfast tea (1 g/100 ml) on serum antioxidant status in 10 healthy volunteers (five male, five female; mean age 21.1 years; mean body mass index 24.0). After a four hour fast an intravenous cannula was inserted and the volunteers drank tea without milk over 20 minutes at lunchtime. Mean serum antioxidant activity before and at 60, 120, and 180 minutes after ingestion of the tea was 430, 434, 447, and 439 $\mu\text{mol/l}$, respectively (no significant change over time).

These results indicate that the rapid ingestion of large amounts of tea flavonoids has little short term impact on serum antioxidant activity and are in contrast to the results of similar studies by our group examining the impact of red wine flavonoids.⁵ This does not exclude the possibility of a long term cumulative impact of drinking tea

on antioxidant status. We suggest that further studies are required to determine the extent to which dietary flavonoids are absorbed and contribute to antioxidant activity in vivo.

SIMON MAXWELL
Senior lecturer

Division of Clinical Pharmacology,
Leicester Royal Infirmary,
Leicester LE2 7LX

GARY THORPE
Biochemical director

Wolfson Applied Technology Laboratory,
Queen Elizabeth Hospital,
Birmingham B15 2TH

- 1 Muldoon MF, Kritchewsky SB. Flavonoids and heart disease. *BMJ* 1996;312:458-9. (24 February.)
- 2 Knekt P, Jarvinen R, Reunanen A, Maatela J. Flavonoid intake and coronary mortality in Finland: a cohort study. *BMJ* 1996;312:478-81. (24 February.)
- 3 Steinberg D, Parthasarathy S, Carew T, Khoo J, Witztum J. Beyond cholesterol. Modifications of low density lipoprotein that increase its atherogenicity. *N Engl J Med* 1989;320:915-24.
- 4 Whitehead TP, Thorpe GHG, Maxwell SRJ. Enhanced chemiluminescence assay for antioxidant capacity in biological fluids. *Analytica Chimica Acta* 1992;266:265-77.
- 5 Maxwell S, Cruickshank A, Thorpe G. Red wine and antioxidant activity in serum. *Lancet* 1994;344:193-4.

Immediate enteral feeding after gastrointestinal resection

Tests of intestinal permeability were inadequate

EDITOR,—Cornelia S Carr and colleagues report a trial with the important message that early enteral feeding after major surgery is both possible and safe.¹ Their conclusion about the changes seen in intestinal permeability is, however, unjustified.

Tests of intestinal permeability were performed preoperatively and on day 5 postoperatively. No significant change was found in the permeability index (lactulose:mannitol absorption ratio) on these two days in the group given early enteral feeding. This does not mean, however, that no rise occurred in this index: the permeability may have increased but returned to normal by day 5. Wicks *et al* showed that, after liver transplantation, permeability was increased only on day 1 and had returned to normal on day 3.² This is corroborated by our preliminary data showing that permeability in a range of patients in an intensive therapy unit may be greatly increased on admission but has returned to normal by day 6 (fig 1).

We also question the authors' experimental design and methodology with respect to the permeability tests. They state that after the test

Fig 1—Mean (SE) changes in permeability in six patients admitted to general intensive therapy unit