Key messages

- Intrapancreatic activation of trypsinogen to trypsin is a key feature of acute pancreatitis; in blood trypsin forms a complex with α_1 antitrypsin
- This complex can be accurately measured in a sensitive immunoassay
- In this study the diagnostic and prognostic accuracy of serum concentrations of trypsin 2- α_1 antitrypsin complex was determined in acute pancreatitis
- The complex was more accurate than trypsinogen 2, C reactive protein, and amylase in differentiating between acute pancreatitis and extrapancreatic disease and in predicting a severe course for the disease
- If the immunoassay could be automated determination of concentrations of trypsin 2- α_1 antitrypsin complex could greatly improve the diagnosis of this common and potentially lethal disease

Funding: Grants from the Finska Läkaresällskapet, the Sigrid Juselius Foundation, and the Finnish Academy of Sciences.

Conflict of interest: None.

- 1 Williamson RC. Early assessment of severity in acute pancreatitis. *Gut* 1984;25:1331-9.
- 2 Ranson JHC, Spencer FC. The role of peritoneal lavage in severe acute pancreatitis. *Ann Surg* 1978;187:565-75.
- 3 Levitt RG, Geisse GS, Sagel SS, Stanley RJ, Evens RG, Koehler RE, et al. Complementary use of ultrasound and tomography in studies of the pancreas and kidney. *Radiology* 1978;126:149-52.
- 4 Dammann HG, Grabbe E, Runge M. Computed tomography and acute pancreatitis. *Lancet* 1980;2:860.

- 5 Block S, Maier W, Bittner M, Bühler M, Malfertheiner H, Beger HG. Identification of pancreas necrosis in severe acute pancreatitis: imaging procedures versus clinical staging. *Gut* 1986;27:1035-42.
- 6 Kivisaari L, Somer K, Standertskjöld-Nordenstam C-G, Schröder T, Kivilaakso E, Lempinen M. A new method for the diagnosis of acute hemorrhagic-necrotizing pancreatitis using contrast-enhanced CT. *Gastrointest Radiol* 1984;9:27-30.
- 7 Steinberg WM, Goldstein SS, Davis ND, Shamma'a J, Anderson K. Diagnostic assays in acute pancreatitis. A study of sensitivity and specificity. *Ann Intern Med* 1985;102:576-80.
- 8 Salt WB, Schenker S. Amylase—its clinical significance: a review of the literature. *Medicine* 1976;55:269-89.
- 9 Mayer AD, McMahon MJ, Corfield AP, Cooper MJ, Dickson AP, Shearer MG, et al. Controlled clinical trial of peritoneal lavage for the treatment of severe acute pancreatitis. *N Engl J Med* 1985;312:399-404.
- 10 Imrie CW, Benjamin IS, Ferguson JC, McKay AJ, Mackenzie I, O'Neill J, et al. A single-centre double-blind trial of Trasylol therapy in primary acute pancreatitis. *Br J Surg* 1978;65:337-41.
- 11 Corfield AP, Cooper MJ, Williamson RCN. Acute pancreatitis: a lethal disease of increasing incidence. *Gut* 1985;26:724-9.
- 12 Ohlsson K, Eddeland A. Release of proteolytic enzymes in bile induced pancreatitis in dogs. *Gastroenterology* 1975;69:668-75.
- 13 Lasso A, Ohlsson K. Protease inhibitors in acute human pancreatitis. Correlation between biochemical changes and clinical course. *Scand J Gastroenterol* 1984;19:779-86.
- 14 Ohlsson K. Acute pancreatitis. Biochemical, pathophysiological and therapeutic aspects. *Acta Gastroenterol Belg* 1988;51:3-12.
- 15 Itkonen O, Koljonen E, Hurme M, Alfthan H, Schröder T, Stenman U-H. Time-resolved immunofluorometric assays for trypsinogen-1 and 2 in serum reveal preferential elevation of trypsinogen-2 in pancreatitis. *J Lab Clin Med* 1990;115:712-8.
- 16 Durie PR, Gaskin KJ, Geokas MC, O'Rourke M, Largman C. Plasma immunoreactive anionic pancreatic trypsin in cystic fibrosis. *J Pediatr Gastroenterol Nutr* 1982;1:337-43.
- 17 Borgström A, Ohlsson K. Studies on the turnover of endogenous cathodal trypsinogen in man. *Eur J Clin Invest* 1978;8:379-82.
- 18 Hedstrom J, Leinonen J, Sainio V, Stenman U-H. Time-resolved immunofluorometric assay of trypsin-2 complexed with alpha 1-antitrypsin in serum. *Clin Chem* 1994;40:1761-5.
- 19 Gudgeon AM, Heath DJ, Hurley P, Jehanli A, Patel G, Wilson C, et al. Trypsinogen activation peptides assay in the early prediction of severity of acute pancreatitis. *Lancet* 1990;335:4-8.
- 20 Fischer CL, Gill C, Forrester MG, Nakamura R. Quantitation of acute phase proteins postoperatively. Value in detection and monitoring of complications. *Am J Clin Pathol* 1976;66:840-6.
- 21 Zweig MH, Campbell G. Receiver-operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. *Clin Chem* 1993;39:561-77.
- 22 Koivunen E, Itkonen O, Halila H, Stenman U-H. Cyst fluid of ovarian cancer patients contains high concentrations of trypsinogen-2. *Cancer Res* 1990;50:2375-8.

(Accepted 16 May 1996)

Perinatal characteristics in relation to incidence of and mortality from prostate cancer

Anders Ekblom, Chung-cheng Hsieh, Loren Lipworth, Alicja Wolk, Jan Pontén, Hans-Olov Adami, Dimitrios Trichopoulos

Abstract

Objective—To test the hypothesis that factors causing morbidity and mortality from prostate cancer may operate in utero.

Design—Matched case-control study of singleton men born between 1874 and 1946 at one hospital.

Setting—Uppsala University Hospital.

Subjects—250 patients with prostate cancer and 691 controls, including 80 patients who died from prostate cancer and their 196 matched controls.

Main outcome measures—Mother's age at menarche, parity, pre-eclampsia or eclampsia before delivery, age at delivery and socioeconomic status; case or control's birth length and weight, placental weight, prematurity derived from gestational age, and presence of jaundice.

Results—Both pre-eclampsia (odds ratio 0, 95% confidence interval 0 to 0.71) and prematurity (0.31, 0.09 to 1.04) were inversely associated with incidence of prostate cancer. Among subjects born full term, placental weight, birth weight, and ponderal index (weight/height³) showed non-significant positive associations with prostate cancer incidence, and stronger associations with mortality.

Conclusion—Prenatal exposures that are likely correlates of pregnancy hormones and other

growth factors are important in prostate carcinogenesis and influence the natural course as well as the occurrence of this cancer.

Introduction

Prostate cancer comprises about 8% of all new cancers among men globally,¹ and the incidence seems to be increasing in many countries and ethnic groups.² Although this increase may be partly due to increased detection rates,³ a genuine increase in the incidence of clinically aggressive cancers is likely since mortality from the disease is also increasing.² The highest incidences have been reported in North America and northern Europe and the lowest in south east Asia.¹⁻⁴ In the United States, incidence and mortality are substantially higher among African-Americans at all ages.⁴

Evidence exists that early life events have a role in prostate cancer, although it is not as strong as for breast cancer.⁵ The incidence of prostate cancer increases in the first generation after migration to areas where it is more common⁶ and recent epidemiological studies also suggest a link with early life.⁷⁻⁸ Intrauterine factors have already been associated with testicular cancer⁹ and breast cancer.¹⁰⁻¹¹ Some recent reports have addressed the possible role of the intrauterine environment in prostate cancer. Ross and Henderson suggested that the intrauterine period may be aetiologically relevant to

Departments of Cancer

Epidemiology and Pathology, Uppsala University, S-751 85 Uppsala, Sweden
Anders Ekblom, associate professor
Alicja Wolk, associate professor
Jan Pontén, professor
Hans-Olov Adami, professor

Cancer Center, University of Massachusetts Medical Center, Worcester, MA, USA
Chung-cheng Hsieh, professor

Department of Epidemiology and Center for Cancer Prevention, Harvard School of Public Health, Boston, MA, USA
Loren Lipworth, research fellow
Dimitrios Trichopoulos, professor

Correspondence to: Dr Ekblom.

BMJ 1996;313:337-41

Table 1—Age at diagnosis of all men who developed prostate cancer and those who died plus numbers of matched controls

Year of birth	All cases				Fatal cases			
	Age < 70 years		Age ≥ 70 years		Age < 70 years		Age ≥ 70 years	
	Cases	Controls	Cases	Controls	Cases	Controls	Cases	Controls
1874-99	2	4	21	41	1	1	11	20
1900-19	44	123	113	275	23	54	27	60
1920	65	227	5	21	18	61	0	0
Total	111	354	139	337	42	116	38	80

prostate cancer¹²; a small study of 21 prostate cancer cases indicated a positive relation between high birth weight and the risk of prostate cancer¹³; and molecular researchers have postulated that early life genomic events may be essential for the subsequent occurrence and progression of adult cancer,^{14 15} leading to the inference that early life events could have a more powerful influence on mortality than incidence.¹⁶

As part of a programme to evaluate intrauterine and perinatal influences on the occurrence of hormone related cancers we have used databases in Sweden that allow linkage of pregnancy and birth records since the last century with national cancer and death registry records on prostate cancer.

Subjects and methods

Because there is no private inpatient treatment in Sweden, hospital medical services are population based and linked to the county in which the patient lives. We identified for study all men who were born at the University Hospital in Uppsala, resident in Uppsala county since 1 January 1947, and were alive in 1958 using the unique 10 digit national registration numbers introduced for all Swedish citizens in 1947.¹⁷ The first six digits provide birth date (year, month, and day). The seventh and eighth digits provide information on county of residence.

All newly diagnosed malignant tumours in Sweden must be reported by both the diagnosing physician and the pathologist or cytologist to one of six regional cancer registries, which pass data to the national cancer registry established in 1958.¹⁸ At the time of this study the national cancer registry was complete up to 31 December 1991, and the Uppsala regional cancer registry was complete up to 31 December 1994.

All men in the national and regional cancer registry who had prostate cancer (ICD-7 code 177) and the code for Uppsala county in their national registration number were included in the study. Cases could have been missed if a man born at the University Hospital moved out of the county before 1947 and subsequently developed prostate cancer.

From the parish in which each patient lived when prostate cancer was diagnosed, we were able to establish his place of birth and family name at birth. At this stage, those who were not born at the University Hospital were excluded. Two hundred and fifty cases of prostate cancer were identified among singletons in the study cohort. We used the Swedish death registry to identify patients who were still alive at the time of censoring the follow up information (December 1994), thus enabling us to evaluate both incidence of and mortality from prostate cancer.

For each case, singleton males born live to the first four or more mothers who were admitted after the case's mother were selected as potential controls. This insured that one individually matched control was alive at the time prostate cancer was diagnosed in the case. We used the cancer and death registries to check that the potential control man was alive and had not had prostate cancer diagnosed at the time of diagnosis or

death of the case. A total of 691 controls were matched to the incident cases and 196 to the fatal cases.

For all cases and controls we manually abstracted from the standardised hospital maternity charts information on the mother's age at menarche, parity, pre-eclampsia or eclampsia before delivery, age at delivery, and socioeconomic status. We also recorded the following data on the child: singleton or twin, birth length and weight, placental weight, prematurity derived from gestational age, and presence of jaundice. The completeness of records exceeded 95% for every item used in this study.

We analysed the data using logistic regression conditional on the matching process.¹⁹ Different sets of models were generated for assessing incidence of and death from prostate cancer. For analysis of death rates we used only cases in whom prostate cancer was the cause of death and excluded controls who did not survive until the age of death of the corresponding case. In addition to the indicators of birth size (birth weight, birth length, placental weight) we used the ponderal index of body mass (kg/m³). This indicator is often used in neonatology to assess intrauterine growth,^{19 20} and it is believed to be particularly sensitive to growth during the last few weeks of gestation. For each of the birth size indicators that were alternatively introduced into the model to avoid problems of collinearity, the unit measure was set as the closest round number corresponding to one standard deviation increase. The level of significance was set at $P < 0.05$ (two sided test).

Results

Table 1 shows the distribution of cases of prostate cancer and controls according to birth cohort and age at diagnosis of all cases. The mean age at diagnosis was 70.5 years; 72% of the subjects were born before 1920.

Table 2 gives the adjusted odds ratios for prostate cancer in relation to perinatal or maternal characteristics. Pre-eclampsia or eclampsia predicted a low risk of prostate cancer and was the only significant association. Prematurity and higher parity were non-significantly associated with low risk, and birth size and jaundice showed non-significant positive associations.

Table 3 presents mutually adjusted odds ratios for death from prostate cancer in relation to perinatal or maternal characteristics. The effect of pre-eclampsia or eclampsia is no longer significant because of small numbers, but prematurity is significantly associated with reduced mortality from prostate cancer. All birth size indicators (with the exception of birth length, for which random measurement error is always proportionately high) were more strongly related to death from prostate cancer than to incidence, and the association with ponderal index was significant. The effects of both neonatal jaundice and parity were larger than for incidence of prostate cancer but still non-significant.

Discussion

This case-control study nested within an explicitly defined cohort, and as such its validity is identical to that of the underlying cohort. Thus, information bias could occur only by non-differential misclassification, which weakens the associations found. Moreover, selection cannot introduce bias because it is unrelated to ascertainment of outcome. Some controls may have had subclinical prostate cancer but the study base (population at risk) for clinical prostate cancer also includes people in this category. Whereas implications of chance variation can be and were measured, confounding requires the existence of risk factors and demonstration of associations of these risk factors with the exposures under consideration. The only established risk factors for prostate cancer are age and race, and both of them were accounted for through matching and conditional

Table 2—Results of multiple logistic regression analysis for incidence of prostate cancer in relation to perinatal or maternal characteristics from a nested case-control study of 250 cases and 691 controls.* Values are numbers (percentages) unless stated otherwise

Characteristic	Cases	Controls	Category or incremental unit for odds ratio	Odds ratio (95% confidence interval)	P value
Mean (SD) maternal age (years)	28.4 (6.5)	28.2 (6.3)	5 years	1.09 (0.94 to 1.25)	0.25
Socioeconomic status:					
Low	212 (84.8)	558 (80.6)	1 class	0.79 (0.54 to 1.16)	0.23
Medium	37 (14.8)	119 (17.2)			
High	1 (0.4)	14 (2.2)			
Mean (SD) age at menarche (years)	14.9 (1.7)	14.9 (1.6)	1 year	0.96 (0.87 to 1.06)	0.39
Parity:					
1	95 (38.2)	256 (37.0)	1	1.00 (reference)	0.61
2	154 (61.8)	435 (63.0)	≥2	0.91 (0.63 to 1.32)	
Pre-eclampsia or eclampsia:					
No	250 (100)	678 (98.1)	No	1.00 (reference)	0.02†
Yes	0 (0)	13 (1.9)	Yes	0.00 (0.00 to 0.71)†	
Neonatal jaundice:					
No	242 (96.8)	675 (97.7)	No	1.00 (reference)	0.30
Yes	8 (3.2)	16 (2.3)	Yes	1.80 (0.60 to 5.41)	
Prematurity (35 wks):					
No	241 (98.8)	633 (95.2)	No	1.00 (reference)	0.06
Yes	3 (1.2)	32 (4.8)	Yes	0.31 (0.09 to 1.00)	
Mean (SD) birth weight (g)	3543.4 (475.5)	3504.0 (528.0)	500	1.04 (0.88 to 1.23)	0.63
Mean (SD) birth length (mm)‡	513 (23)	512 (23)	20	1.00 (0.87 to 1.16)	0.97
Mean (SD) placental weight (g)‡	649.9 (135.0)	643.6 (132.8)	150	1.12 (0.93 to 1.35)	0.24
Ponderal index‡	26.2 (2.8)	26.1 (2.9)	3	1.07 (0.90 to 1.29)	0.44

*Some missing values.

†Exact 95% confidence interval and P value from crude data.

‡Introduced one at a time in place of birth weight.

analysis for age and because there are few non-white Swedish citizens. There is no agreement about the quantitative importance of other suggested risk factors for prostate cancer, including fat intake,²²⁻²⁴ vasectomy,²⁴⁻²⁶ and physical activity,²⁷ and no evidence that any of these is associated with perinatal characteristics.

SEX STEROIDS

Sex steroids may affect prostate cancer given their role in normal growth and function of the prostate.²⁸⁻²⁹ However, epidemiological studies of some steroid hormones (oestradiol, testosterone, and androstenedione) and prostate cancer in humans have proved

inconclusive.³⁰⁻³³ The activity of 5α-reductase (involved in the conversion of circulating testosterone to the biologically active dihydrotestosterone) is higher among white Americans than among low risk native Chinese,³⁴ but its role has not been evaluated in an epidemiological study of prostate cancer.

Although we were primarily examining the effect of intrauterine influences on the incidence of prostate cancer, mortality may be more strongly related to perinatal variables than incidence.¹⁶ Our choice of exposure variables was constrained by availability, but several of them are thought to reflect intrauterine growth, which is partly modulated by steroid hormones.³⁵⁻³⁸ Ponderal index is an indicator of late stage intrauterine

Table 3—Results of multiple logistic regression analysis for death from prostate cancer in relation to perinatal or maternal characteristics from a nested case-control study of 80 cases and 196 controls*

Characteristic	Cases	Controls	Category or incremental unit for odds ratio	Odds ratio (95% confidence interval)	P value
Mean (SD) maternal age (years)	28.4 (6.2)	28.1 (6.5)	5 years	1.06 (0.82 to 1.36)	0.66
Socioeconomic status:					
Low	69 (86.2)	164 (83.7)	1 class	0.86 (0.39 to 1.88)	0.21
Medium	11 (13.8)	30 (15.3)			
High	0 (0)	2 (0.9)			
Mean (SD) age at menarche (years)	15.0 (1.9)	14.9 (1.8)	1 year	0.94 (0.79 to 1.12)	0.51
Parity:					
1	31 (38.8)	62 (31.6)	1	1.00 (reference)	0.07
2	49 (61.3)	134 (68.4)	≥2	0.52 (0.26 to 1.04)	
Pre-eclampsia or eclampsia:					
No	80	193	No	1.00 (reference)	0.56†
Yes	0	3	Yes	0.00 (0.00 to 3.11)†	
Neonatal jaundice:					
No	77 (96.3)	194 (99.0)	No	1.00 (reference)	0.09
Yes	3 (3.8)	2 (1.0)	Yes	8.51 (0.72 to 101.06)	
Prematurity (35 wks):					
No	78 (100)	175 (94.1)	No	1.00 (reference)	0.02†
Yes	0 (0)	11 (5.9)	Yes	0.00 (0.00 to 0.72)†	
Mean (SD) birth weight (g)	3582.9 (431.5)	3484.1 (506.9)	500	1.22 (0.87 to 1.70)	0.75
Mean (SD) birth length (mm)‡	51.3 (2.3)	51.4 (2.2)	20	0.91 (0.68 to 1.21)	0.51
Mean (SD) placental weight (g)‡	674.5 (132.4)	642.9 (129.3)	150	1.43 (0.99 to 2.06)	0.06
Ponderal index‡	26.5 (2.8)	25.7 (2.7)	3	1.60 (1.07 to 2.39)	0.02

*Some missing values.

†Exact 95% confidence interval and P value from crude data.

‡Introduced one at a time in place of birth weight.

Key messages

- Prostate cancer is the most common malignancy among men in the West, but little is known about the aetiology
- Recent evidence suggests that events in utero may affect the risk of prostate cancer later in life
- In this study 250 men born since 1874 at one hospital and who developed prostate cancer were identified and compared with 691 controls
- Pre-eclampsia and prematurity were associated with a reduced risk of prostate cancer and ponderal index was associated with an increased risk of mortality from prostate cancer
- Prenatal exposures that are likely correlates of pregnancy hormones and other growth factors seem to be of importance in the prostate carcinogenesis

growth.²⁰ Moreover, concentrations of oestrogens and other hormones are generally lower in women with pre-eclampsia or eclampsia,^{35 8 40} and there is some evidence that primiparous women have higher oestrogen concentrations in pregnancy than multiparous women.^{41 42} We found that prematurity and pre-eclampsia or eclampsia reduced the risk of prostate cancer and that birth size was more strongly associated with mortality than incidence. These findings agree with our hypothesis and support a role for sex hormones in prostate cancer.

MORTALITY

The existence of associations between perinatal factors and prostate cancer indicate that early life is important in the long carcinogenesis process. It seems unlikely that the associations were confounded by post-natal exposures, and there is no evidence to suggest confounding. The substantially stronger associations of most of the perinatal characteristics with death from prostate cancer than with incidence further suggest that not only the future occurrence but also the biological features of the tumour are programmed in utero. This may explain why the strong geographical variations in incidence of prostate cancer are not mirrored by a similar pattern in the prevalence of subclinical disease detected at necropsy.⁴³

Loeb and other investigators have postulated that an early phenomenon, perhaps affecting factors controlling genetic stability, is needed to explain the high mutation rate required for the accumulation of the genetic changes leading to overt cancer.^{14 15 44} According to this hypothesis, if clonal cells remain genetically unstable even after the appearance of cancer, perinatal determinants of the early event should be expected to forecast metastasis of, and death from, a particular cancer even better than its incidence. This theory provides a possible explanation for our epidemiological findings. It is unclear how high concentrations of growth hormones in utero would influence subsequent mutation rates, but it could be through enlargement of a pool of potential target cells for malignant transformation or cell proliferation.

Our working hypothesis and results are compatible with two of the most striking descriptive characteristics of the disease. Clinical prostate cancer and mortality from prostate cancer are substantially higher among Americans and Europeans than among Asians,^{1 4} and within the United States incidence of and mortality from the disease are substantially higher among black than among white Americans.^{4 9} Birth weight is lower among Japanese and Chinese than Americans,⁴⁵ and pregnancy steroids are much higher among black

women.⁹ Moreover, long term increases in the incidence of prostate cancer and mortality have been preceded by long term increases in weight and obesity of the newborn.

It may be stretching biologic plausibility that a cancer which appears clinically in elderly people would have an origin in utero. However, it has been shown recently that prostatic intraepithelial neoplasia is already quite common among men in their 20s.^{46 47}

We thank Ann Almqvist, Ulrika Lund, and Monica Rundgren for abstracting the data.

Conflict of interest: None.

Funding: This study was supported by grants from the Swedish Cancer Society. LL received a training award in cancer epidemiology from the US National Institutes of Health/ National Cancer Institute (T32CA09001-19A2).

- 1 Parkin DM, Pisani P, Ferlay J. Estimates of the worldwide incidence of eighteen major cancers in 1985. *Int J Cancer* 1993;54:594-606.
- 2 Wingo PA, Tong T, Bolden S. Cancer statistics 1995. *CA* 1995;45:8-30.
- 3 Potosky AL, Kessler L, Gridley G, Brown CC, Horn JW. Rise in prostatic cancer incidence associated with increased use of transurethral resection. *J Natl Cancer Inst* 1990;82:1624-8.
- 4 Parkin DM, Muir CS, Whelan SL, Gao Y-T, Ferlay J, Powell J, eds. *Cancer incidence in five continents*. Vol 6. Lyons, France: International Agency for Research on Cancer, 1992. (IARC scientific publication No 120.)
- 5 Lipworth L. Epidemiology of breast cancer. *Eur J Cancer Prev* 1995;4:7-30.
- 6 Tomatis L, ed. *Cancer: Causes, occurrence and control*. Lyon, France: International Agency for Research on Cancer, 1990. (IARC scientific publication No 100.)
- 7 Andersson S-O, Baron J, Wolk A, Lindgren C, Berstrom R, Adami H-O. Early life risk factors for prostate cancer: a population-based case-control study in Sweden. *Cancer Epidemiol Biomark Prev* 1995;4:187-92.
- 8 Rotkin ID. Studies of the epidemiology of prostatic cancer: expanded sampling. *Cancer Treat Rep* 1977;61:173-80.
- 9 Henderson BE, Bernstein L, Ross RK, Depue RH, Judd HL. The early in utero oestrogen and testosterone environment of blacks and whites: potential effects on male offspring. *Br J Cancer* 1988;57:216-8.
- 10 Trichopoulos D. Hypothesis: does breast cancer originate in utero? *Lancet* 1990;335:939-40.
- 11 Ekblom A, Thurfjell E, Hsieh CC, Trichopoulos D, Adami HO. Perinatal characteristics and adult mammographic patterns. *Int J Cancer* 1995;61:177-80.
- 12 Ross RD, Henderson BE. Do diet and androgens alter prostate cancer via a common etiologic pathway? *J Natl Cancer Inst* 1994;86:252-4.
- 13 Tibblin G, Eriksson M, Cnattingius S, Ekblom E. High birth weight as a predictor of prostate cancer risk. *Epidemiology* 1995;6:423-4.
- 14 Loeb LA. Microsatellite instability: marker of a mutator phenotype in cancer. *Cancer Res* 1994;54:5059-63.
- 15 Fishel R, Lescoe MK, Rao MR, Copeland NG, Jenkins NA, Garber J, et al. The human mutator gene homolog MSH2 and its association with hereditary nonpolyposis colon cancer. *Cell* 1993;75:1027-38.
- 16 Trichopoulos D, Lipworth L. Is cancer causation simpler than we thought, but more intractable? *Epidemiology* 1995;6:347-9.
- 17 Lunde AS, Lundeborg S, Lettenstrom GS, Thugesen L, Huebner J. The person-number systems of Sweden, Norway, Denmark and Israel. *Vital Health Stat* [2] 1980;84:5-11.
- 18 Cancer Registry. *Cancer incidence in Sweden 1987*. Stockholm: National Board of Health and Welfare, 1990.
- 19 Breslow NE, Day NE. *Statistical methods in cancer research*. Vol 1. *The analysis of case-control studies*. Lyon, France: International Agency for Research on Cancer, 1980 (IARC scientific publication No 32.)
- 20 Lubchenco LO, Hansman C, Boyd E. Intrauterine growth in length and head circumference as estimated from live births at gestational ages from 26 to 42 weeks. *Pediatrics* 1966;37:403-8.
- 21 Daikoku NH, Johnson JWC, Graf C, Kearney K, Tyson JE, King TM. Patterns of intrauterine growth retardation. *Obstet Gynecol* 1979;54:211-9.
- 22 Giovannucci E, Rimm EB, Colditz GA, Stampfer MJ, Ascherio A, Chute CC, et al. A prospective study of dietary fat and risk of prostate cancer. *J Natl Cancer Inst* 1993;85:1571-9.
- 23 Whittemore AS, Kolonel LN, Wu AH, John EM, Gallagher RP, Howe GR, et al. Prostate cancer in relation to diet, physical activity, and body size in blacks, whites, and Asians in the United States and Canada. *J Natl Cancer Inst* 1995;87:652-61. Hayes RB. Are dietary fat and vasectomy risk factors for prostate cancer? *J Natl Cancer Inst* 1995;87:629-31.
- 24 John EM, Whittemore AS, Wu AH, Kolonel LN, Hislop TG, Howe GR, et al. Vasectomy and prostate cancer: results from a multiethnic case-control study. *J Natl Cancer Inst* 1995;87:662-69.
- 25 Giovannucci E, Ascherio A, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. A prospective cohort study of vasectomy and prostate cancer in US men. *JAMA* 1993;269:873-7.
- 26 Thune I, Lund E. Physical activity and the risk of prostate and testicular cancer: a cohort study of 53 000 Norwegian men. *Cancer Causes Control* 1994;5:549-56.
- 27 Coffey DS, Isaacs JT. Control of prostate growth. *Urology* 1981;17 (Suppl 3):17-24.
- 28 Greenwald P. Prostate. In: Schottenfeld D, Fraumeni JF Jr, eds. *Cancer epidemiology and prevention*. Philadelphia: Saunders, 1982:9138-46.
- 29 Barrett-Connor E, Garland C, McPhillips JB, Khaw KT, Wingard DL. A prospective, population-based study of androstenedione, estrogens and prostatic cancer. *Cancer Res* 1990;50:169-73.
- 30 Hsing A, Comstock G. Serum hormone and risk of subsequent prostate cancer. *Am J Epidemiol* 1989;130:829.
- 31 Drafta D, Proca E, Zamfir V, Schindler AE, Neacsu E, Stroe E. Plasma steroids in benign prostatic hypertrophy and carcinoma of the prostate. *J Steroid Biochem* 1982;17:689-93.

- 32 Andersson SO, Adami HO, Bergstrom R, Wide L. Serum pituitary and sex steroid hormone levels in the aetiology of prostatic cancer—a population-based case-control study. *Br J Cancer* 1993;68:97-102.
- 33 Ross RK, Bernstein L, Lobo RA, Shimizu H, Stanczyk FZ, Pike MC, *et al*. 5- α -reductase activity and risk of prostate cancer among Japanese and US white and black males. *Lancet* 1992;339:887-9.
- 34 Kloppe A, Jandial V, Wilson G. Plasma steroid assay in the assessment of foetoplacental function. *J Steroid Biochem* 1975;6:651-6.
- 35 McPadyen IR, Worth HG, Wright DJ, Notta SS. High oestrogen excretion in pregnancy. *Br J Obstet Gynaecol* 1982;89:994-9.
- 36 Gerhard I, Vollmar B, Runnebaum B, Kliinga K, Haller U, Kubli F. Weight percentile at birth. II. Prediction by endocrinological and sonographic measurements. *Eur J Obstet Gynecol Reprod Biol* 1987;26:313-28.
- 37 Petridou E, Panagiotopoulou K, Katsouyanni K, Spanos E, Trichopoulos D. Tobacco smoking, pregnancy estrogens and birth weight. *Epidemiology* 1990;1:247-50.
- 38 Long PA, Abell DA, Beischer NA. Fetal growth and placental function assessed by urinary estriol excretion before the onset of pre-eclampsia. *Am J Obstet Gynecol* 1979;135:344-7.
- 39 Rosing U, Carlstrom K. Serum levels of unconjugated and total estrogens and dehydroepiandrosterone, progesterone and urinary estriol excretion in pre-eclampsia. *Gynecol Obstet Invest* 1984;18:199-205.
- 40 Bernstein L, Depue RH, Ross RK, Judd HL, Pike MC, Henderson BE. Higher maternal levels of free estradiol in first compared to second pregnancy: early gestational differences. *J Natl Cancer Inst* 1986;76:1035-9.
- 41 Panagiotopoulou K, Katsouyanni K, Petridou E, Garas Y, Tzonou A, Trichopoulos D. Maternal age, parity and pregnancy estrogens. *Cancer Causes Control* 1990;1:119-24.
- 42 Nomura AMY, Kolonel LN. Prostate cancer: a current perspective. *Am J Epidemiol* 1991;133:200-27.
- 43 Loeb LA. Mutator phenotype may be required for multistage carcinogenesis. *Cancer Res* 1991;51:3075-9.
- 44 Wang X, Guyer B, Paige DM. Differences in gestational age-specific birth-weight among Chinese, Japanese and White Americans. *Int J Epidemiol* 1994;23:119-28.
- 45 Sakr WA, Haas GP, Cassin BF, Pontes FE, Crissman JD. The frequency of carcinoma and intraepithelial neoplasia of the prostate in young male patients. *J Urol* 1993;150:379-85.
- 46 Chevillet JC, Bostwick DG. Postatrophic hyperplasia of the prostate. A histologic mimic of prostatic adenocarcinoma. *Am J Surg Pathol* 1995;19:1068-76.

(Accepted 1 June 1996)

Fluctuations of HIV load in semen of HIV positive patients with newly acquired sexually transmitted diseases

M C Atkins, E M Carlin, V C Emery, P D Griffiths, F Boag

Most HIV infections worldwide are acquired sexually, yet most coital episodes involving an infected partner do not result in acquisition of HIV. Acquisition of HIV must depend on both the volume of secretions transferred from the infected partner (donor) and the concentration of HIV present in the secretions. However, exposure alone to such a virus inoculum is clearly insufficient to ensure transmission. The coexistence of other sexually transmitted diseases in either the recipient or donor could potentially increase the risk of transmission by causing genital ulcers or by releasing inflammatory cytokines which increase HIV replication.¹

We measured the HIV load in the semen and blood of HIV infected patients presenting with acute sexually transmitted diseases to determine if viral load in semen decreases when patients receive specific treatment for sexually transmitted diseases.

Subjects, methods, and results

Between October 1994 and June 1995, four asymptomatic homosexual men positive for HIV-1 with CD4 T cell counts between 330 and 600 cells/ μ l participated in the study. The men had been HIV positive for 3-10 years and none was receiving anti-retroviral treatment. The patients presented with a purulent urethral discharge which developed after unprotected orogenital sex. In each case, microscopy of a Gram stained urethral smear revealed 10 or more neutrophils per high power field ($\times 1000$), and in three cases (patients 1, 2, and 4) Gram negative intracellular diplococci were identified; urethral infection with *Neisseria gonorrhoea* was diagnosed at presentation and subsequently confirmed by culture. Non-gonococcal urethritis was diagnosed in patient 3. Patients with gonococcal urethritis were treated with single oral doses of ciprofloxacin 500 mg (patients 1 and 4) or ampicillin 3 g with probenecid 1 g (patient 2). The patient with non-gonococcal urethritis received doxycycline 100 mg bd for seven days. Subsequent urethral samples from patients 1, 3, and 4 were non-purulent and negative on microscopy and culture. Patient 2 had a post-gonococcal urethritis with a persistent discharge which resolved when treated with a course of doxycycline. No patient had any signs or symptoms of prostatitis. Blood

and semen samples were collected on the day of presentation (before treatment) and at subsequent visits.

DNA was extracted from whole blood or semen using a commercially available DNA extraction method (Digene Laboratories, Germany). Quantitative polymerase chain reaction for proviral DNA was performed using nested primers to the *gag* region of HIV-1 (outer primers 5' GAGGAGCCACCCACAATATT and 5' TAGGTGGATTATTTGTTCATCCA; inner primers 5' TGCTAAACACAGTTGGGGGGA and 5' CCTGAAGGGTACTAGTAGTT) and coamplification of an internal control sequence.

Figure 1 shows HIV proviral loads during the period of treatment. HIV-1 proviral load in the semen of these subjects declined ($P < 0.05$ Mann-Whitney test) when their intercurrent sexually transmitted diseases were treated; there were no significant changes in the blood.

Fig 1—Fluctuations in HIV proviral loads in serial semen and blood samples from four patients

Department of Virology,
Royal Free Hospital
School of Medicine,
London NW3 2QG
M C Atkins, research fellow
V C Emery, reader in virology
P D Griffiths, professor of
virology

Department of
Genitourinary Medicine,
Royal Chelsea and
Westminster Hospital,
London SW10 9TH
E M Carlin, senior registrar
F Boag, consultant physician

Correspondence to:
Dr Boag.

BMJ 1996;313:341-2