- 2 Ethnicity, race, and culture: guidelines for research, audit, and publication. *BMJ* 1996;312:1094. (27 April.)
- 3 Senior PA, Bhopal R. Ethnicity as a variable in epidemiological research. *BMJ* 1994;309:327-30.
- 4 Aspinall PJ. Department of Health's requirement for mandatory collection of data on ethnic group of inpatients. *BMJ* 1995;311:1006-9.
- 5 Shaunak S, Lakhani SR, Abraham R, Maxwell JD. Differences among Asian patients. *BMJ* 1986;293:1169.

Links need to be maintained with census definitions

EDITOR,—The definition and meaning of terms relating to race, ethnicity, and culture are issues to which social scientists have devoted hours and pages of thought. Some doubt whether they are indeed real entities (like "society"), but, as many studies have now shown, they are real in their effects. Kwame McKenzie and N S Crowcroft's editorial and "style guide" are helpful pointers out of the maze.^{1,2}

It is important to emphasise that culture and ethnicity are multivariate. Researchers should (for example) make reference to religion as it is likely to be a key factor in behaviour. The style guide seems to omit language, which, at least in relation to people of South Asian descent, may provide as much useful information as does religion—and certainly more than the catch-all term "Indian."

The guidance also emphasises the need to let people identify their own ethnicity.² It is therefore good news for those who, like me, have advocated this and promoted ethnic monitoring that general practitioners in Leicester and Lincolnshire found the procedure so easy to operate.³ Hopefully, such open debate will reassure those who, by virtue of their uncertainty about the value of ethnic monitoring, make the collection of data problematic. The results will be of little value, however, unless linked to sound estimates of the population denominator. For this reason it is imperative that, however freely people are allowed to choose or describe their identity, a link is maintained to the complex and carefully designed structures used in the census.⁴ Without some degree of agreement we shall see no benefit in epidemiology or the delivery of services and a backlash against the collection of data may result.

The provision of well thought out guidelines is a useful start and may help to avoid the confusion that seems to have developed in the United States.⁵ We may now look forward to an end of the muddled thinking embodied in such old and ugly terms as "Europid" and the confusion between nationality as stated in a passport, which has no clinical importance, and ethnic group, which manifestly does.

MARK R D JOHNSON
Senior research fellow

University of Warwick,
Coventry CV4 7AL

- 1 McKenzie K, Crowcroft NS. Describing race, ethnicity, and culture in medical research. *BMJ* 1996;312:1054. (27 April.)
- 2 Ethnicity, race, and culture: guidelines for research, audit, and publication. *BMJ* 1996;312:1094. (27 April.)
- 3 Pringle M, Rothera I. Practicality of recording patient ethnicity in general practice: descriptive intervention study and attitude survey. *BMJ* 1996;312:1080-2. (27 April.)
- 4 Aspinall PJ. Department of Health's requirement for mandatory collection of data on ethnic group of inpatients. *BMJ* 1995;311:1006-9.
- 5 Abdieh L, Hahn RA. Use of the terms "race," "ethnicity" and "national origins"; a review of articles in the American Journal of Public Health, 1980-1989. *Ethnicity and Health* 1996; 1: 95-8.

Guidelines will encourage the thinking that underpins racism in medicine

EDITOR,—I am concerned that Kwame McKenzie and N S Crowcroft's editorial on the need to describe race, ethnicity, and culture in medical

research will encourage prejudice.¹ I am sure that that is not the authors' intention, but it is what they are most likely to accomplish if they continue in their present line of thought. They offer the spurious argument that "if researchers want to be able to compare results of studies now and in future, a framework is needed for the classification of ethnic and cultural groups." Not so. There are more meaningful criteria on which people can be compared, such as sex, age, place of residence, environmental conditions, occupation, income, and lifestyle.

McKenzie and Crowcroft go on to suggest that we need to "agree on how culturally and ethnically to demarcate the people of the world." Why? Can the authors not see that most of our problems are the direct result of just such demarcation? What do they understand to be the underlying cause of such phenomena as racism, "ethnic cleansing," the caste system in the Indian subcontinent, the pogroms in Rwanda and Burundi, and the "troubles" in Northern Ireland?

I am sorry that the *BMJ* lends credibility to the views of these two authors by producing guidelines on describing people by ethnicity, race, or culture in research, audit, and publications.² This only encourages the thinking underpinning the racism that is endemic in the selection of medical students and in the training, employment, and promotion of doctors in Britain.

We need to take a conscious decision not to "demarcate" people but to encourage people to think non-racially and non-ethnically. As one of the most widely read international journals, the *BMJ* is uniquely placed to take a leading role in this process. I would not support censorship, but I would support the *BMJ*'s rejection of all letters, articles, and papers that seek to divide people into distinct racial, ethnic, or cultural groups. This is not just because of a wish to stop supporting prejudice but also because, to quote the *BMJ* itself, "race has limited biological validity" and "culture is difficult to measure and describe."² Lifestyle does, but race, culture, and ethnicity do not, merit inclusion in our list of valid epidemiological variables.

IKECHUKWU OBIALO AZUONYE
Consultant psychiatrist

South Western Hospital,
London SW9 8EA

- 1 McKenzie K, Crowcroft NS. Describing race, ethnicity and culture in medical research. *BMJ* 1996;312:1054. (27 April.)
- 2 Ethnicity, race, and culture: guidelines for research, audit, and publication. *BMJ* 1996;312:1094. (27 April.)

Authors' reply

EDITOR,—We agree with David de Bono's comments and the findings of James McAuley and colleagues that there is as much variability in "white" groups as in any other. In fact, we had suggested that a "white" example should be included in the guidelines to make this very point. Unfortunately, this was not done. Part of the motivation for our editorial and the guidelines was to dissipate the myth that groups defined by skin colour are homogeneous in any meaningful sense other than that which corresponds to racialised thinking.

Mark R D Johnson points out the importance of language and of maintaining links between self coding and census categories so that population denominators are available. McAuley and colleagues' letter, however, underlines the limitations inherent in the use of any particular classification on its own.¹ Furthermore, individuals may change their self coded ethnic group as society changes, making interpretation of time trends problematic.

Ikechukwu Obialo Azuonye speaks from the heart when he calls for a world where racial, ethnic, and cultural differences are not important. We would all like to live in such a world, but we do not at present, and we have to deal with the problems of this one. There are ethnic differences in the rates of illness and in the uptake of health services.² There is racism in society. Research that recognises these facts can lead to better policies and services to combat inequity, and to a better understanding of the aetiology of diseases. How could appropriate interventions be developed if we were unable to measure and compare the effects of racism, genetics, poverty, and culture?

Thousands of papers are published every year that refer to race, ethnicity, or culture. If we are to use such information effectively to improve health and health care then we need a coherent framework within which to report and interpret findings. We wrote our editorial and the guidelines in the hope that such a framework could be agreed internationally and to start a correspondence on the subject. We hope that letters such as these published here will lead to the guidelines being refined so that they meet the needs of both researchers and the subjects of research.

KWAME MCKENZIE

Honorary research fellow
King's College Hospital Medical School and Institute of

Psychiatry,
London SE5 8AX

NATASHA CROWCROFT

Fellow

European Programme for Intervention Epidemiology Training,
Institute of Hygiene and Epidemiology,
Brussels 1050,
Belgium

- 1 McKenzie KJ, Crowcroft NS. Race, ethnicity, culture, and science. *BMJ* 1994;309:286-7.

- 2 Cruikshank JK, Beevers DG. *Ethnic factors in health and disease*. Oxford: Butterworth-Heinemann, 1989.

Addison's disease

Secondary adrenal failure in critically ill patients is underrecognised

EDITOR,—C M Brosnan and N F C Gowing's lesson of the week is a succinct reminder of the diverse presentation of primary adrenal failure and emphasises the risk of shock that is unresponsive to inotropes in these patients.¹ There is a growing realisation that secondary adrenal failure can also occur in critically ill patients and is probably more common than previously thought.² Establishing the diagnosis may be difficult in these patients, since they do not exhibit the normal electrolyte abnormalities of adrenal failure and usually have a serum cortisol concentration in the normal range for the healthy population. Typically, however, they present with septic shock that is unresponsive to inotropes and has no apparent focus of infection.³

Treatment with physiological doses of steroids leads to a rapid resolution of the shock, and inotropes can usually be withdrawn within 48 hours of the start of such treatment.⁴ The adrenal failure is normally transient, so the steroids can be tapered off once the patient's general condition improves.

Adrenal failure in critically ill patients is underrecognised and may be life threatening. The diagnosis should be considered in any patient in whom septic shock occurs without an obvious source of infection.

C GRANGER
Senior registrar

Department of Anaesthesia,
Blackpool Victoria Hospital,
Blackpool FY3 8NR

- 1 Brosnan CM, Gowing NFC. Addison's disease. *BMJ* 1996;312:1085-7. (27 April.)

- 2 Merry WH, Caplan RH, Wickus GG, Reynertson RH, Kiskin WA, Cogbill TH, et al. Post-operative acute adrenal failure caused by transient corticotropin deficiency. *Surgery* 1994;116:1095-100.
- 3 Bouachour G, Tiro P, Varache N, Gouello JP, Harry P, Alquier P. Hemodynamic changes in acute adrenal insufficiency. *Intensive Care Med* 1994;20:138-41.
- 4 Baldwin WA, Allo M. Occult hypoadrenalism in critically ill patients. *Arch Surg* 1993;128:673-6.

Adrenal insufficiency should be excluded before thyroxine replacement is started

EDITOR,—We strongly agree with C M Brosnan and N F C Gowing's comment that Addison's disease remains the unforgiving master of non-specificity and disguise.¹ We report two cases of Addison's disease in which the finding of a raised thyroid stimulating hormone concentration during investigations of weight loss prompted inappropriate treatment with thyroxine and precipitated an Addisonian crisis.

Case 1—A 58 year old woman presented to her general practitioner with persistent nausea, vomiting, and weight loss. Thyroxine was started when investigations showed a thyroid stimulating hormone concentration of 27 mU/l (normal 0.3-5.0). Her condition deteriorated considerably, with abdominal pain and vomiting necessitating emergency admission. On examination she was dehydrated and shocked. Abdominal examination was unremarkable. Investigations showed a low thyroid stimulating hormone concentration (<0.05 mU/l) and a high total thyroxine concentration of 251 nmol/l (65-150). An abdominal ultrasound scan and results of upper gastrointestinal endoscopy were normal. She was noted to be pigmented. A short tetracosactrin test was performed, and intravenous hydrocortisone was started. Her improvement was dramatic, with resolution of all the symptoms. A positive result of the short tetracosactrin test, raised adrenocorticotrophic hormone concentration, and presence of adrenal antibodies confirmed autoimmune Addison's disease. Results of thyroid function tests remained normal after thyroxine was withdrawn.

Case 2—A 39 year old woman was admitted direct from the outpatient department, having been referred for investigation of weight loss, increasing weakness, and epigastric discomfort. Four weeks before her admission thyroxine replacement had been started because her thyroid stimulating hormone concentration was raised. On examination she was unwell and shocked, but otherwise the examination was unremarkable. Investigations showed a low sodium and a high potassium concentration. Withdrawal of thyroxine and replacement with hydrocortisone resulted in a dramatic resolution of her symptoms, as in case 1. Investigations confirmed autoimmune Addison's disease. Results of thyroid function tests remained normal.

The association of autoimmune thyroid and adrenal disease is well recognised and described.² Addisonian crisis precipitated by thyroxine replacement in patients with hypothyroidism has been documented.³ To avoid precipitating a potentially fatal Addisonian crisis in patients with hypothyroidism it is important to exclude adrenal insufficiency before starting thyroxine replacement treatment. In many patients with adrenal insufficiency abnormal blood thyroid variables return to normal after treatment with corticosteroids.⁴

I A OSMAN

Staff grade doctor in general medicine

PETER LESLIE

Consultant physician

Medical Department,
Borders General Hospital,
Melrose TD6 9BS

- 1 Brosnan CM, Gowing NFC. Addison's disease. *BMJ* 1996;312:1085-6. (27 April.)

- 2 Papadopoulos KI, Hallengren B. Polyglandular autoimmune syndrome type II in patients with idiopathic Addison's disease. *Acta Endocrinol* 1990;122:472-8.
- 3 Banitt PF, Munson AK. Addisonian crisis after thyroid replacement. *Hosp Pract* 1986;21:132, 134.
- 4 Gharib H, Hodgson SF, Gastineau CF, Scholz DA, Smith LA. Reversible hypothyroidism in Addison's disease. *Lancet* 1972;iii:734-6.

Hydrocortisone should be started immediately adrenal insufficiency is considered

EDITOR,—The most important lesson to be learnt from the cases of Addison's disease reported by C M Brosnan and N F C Gowing is not stated.¹ In any patient in whom Addisonian crisis is suspected, serum should be saved for measurement of cortisol and adrenocorticotrophic hormone concentrations and treatment with intravenous hydrocortisone 100 mg should be started immediately. This should then be continued four hourly. The search for the precipitating cause can be concurrent.

Case 1 in Brosnan and Gowing's paper is a fairly classic case, and the biochemical abnormalities are suggestive of adrenal insufficiency. The authors state that the diagnosis of Addison's disease was considered, yet no treatment was started. I would hope that the lesson of the week is that, having considered the possibility of adrenal insufficiency, despite its non-specific presentation, you should treat it. Failure to do so has rapidly lethal results, as the case shows.

PETER LEMAN

Registrar

Accident And Emergency Department,
Lewisham Hospital,
London SE13 6LH

- 1 Brosnan CM, Gowing NFC. Addison's disease. *BMJ* 1996;312:1085-7. (27 April.)

N of 1 trials

Research is needed into why such trials are not more widely used

EDITOR,—Ten years after n of 1 trials were discovered for medicine by clinical epidemiologists from McMaster University,^{1,2} Jeffrey Mahon and colleagues' study on theophylline for irreversible chronic airflow limitation merits several comments.³ The authors cautiously state that further studies are needed before widespread use of n of 1 trials can be advocated in routine clinical practice. In 1986 we also wondered what the relevance of the contributions of this design to patients' care would be.⁴ In view of the progress achieved since then we now hope that greater attention will be given to reasons why these designs are not more widely used despite their suitability to many problems.³

N of 1 trials have much in common with the classic crossover clinical trial.⁴ In particular, both types of trials are subject to two sources of bias, sometimes referred to as treatment-period interaction. These sources of bias are, firstly, carryover effects (pharmacological effects persist in the second period) and, secondly, period effects (responses in the second period differ substantially from those in the first period, possibly owing to saturation of some biological response, fluctuations in the severity of disease, patients responding differently because of the "practice" in the first period, or other changes in environmental circumstances). In classic crossover studies, tests for carryover and period effects should routinely be conducted in the early stages of the analysis.⁴ We thus wonder why tests for treatment-period interaction are not more common in n of 1 trials.

A given set of n of 1 trials will also share several intriguing features with another design,

the recently developed case crossover study, which seems particularly appropriate for studying transient effects on the risk of acute events.⁵ Might some of the analytical strategies proposed for the case crossover design be applicable to a study of n of 1 trials such as the one by Mahon and colleagues? We believe that most such studies of trials could benefit from methodological cross fertilisation—that is, from developments achieved both before and after the advent of n of 1 trials.

Beyond methodological matters, several issues that are clinically relevant warrant further scrutiny. One concerns the nature of the inferences made possible by the three types of studies (for example, on the therapeutic effectiveness of a given drug in groups of patients versus the clinical value of the drug in an individual patient). Surely other issues lie ahead.

MIQUEL PORTA

Associate professor

Institut Municipal d'Investigació Mèdica,
Universitat Autònoma de Barcelona,
E-08003 Barcelona,
Spain

FRANCISCO BOLUMAR

Professor

ILDEFONSO HERNANDEZ

Associate professor

JESUS VIOQUE

Associate professor

Department of Public Health,
Universidad de Alicante,
03080-Sant Joan,
Alicante,
Spain

- 1 McLeod RS, Taylor DW, Cohen Z, Cullen JB. Single-patient randomised clinical trial. Use in determining optimum treatment for patient with inflammation of Kock continent ileostomy reservoir. *Lancet* 1986;ii:726-8.
- 2 Guyatt G, Sackett D, Taylor DW, Chong J, Roberts R, Pugsley S. Determining optimal therapy—randomized trials in individual patients. *N Engl J Med* 1986;314:889-92.
- 3 Mahon J, Laupacis A, Donner A, Wood T. Randomised study of n of 1 trials versus standard practice. *BMJ* 1996;312:1069-74. (27 April.)
- 4 Porta M. The search for more clinically meaningful research designs: single-patient randomized clinical trials. *J Gen Intern Med* 1986;1:418-9.
- 5 Mittleman MA, Maclure M, Robins JM. Control sampling strategies for case-crossover studies: an assessment of relative efficiency. *Am J Epidemiol* 1995;142:91-8.

Placebos should be abandoned

EDITOR,—Jeffrey Mahon and colleagues report that n of 1 trials led to less use of theophylline without adverse effects on exercise capacity or quality of life in patients with irreversible chronic airflow limitation in a tertiary centre.¹ This finding is valuable but highly specific and does not justify the larger trial that the authors advocate to confirm the result. As the authors go some way to acknowledging, there are many other diseases, settings, groups of doctors and patients, forms of treatment, periods of follow up, and outcome measures that could be studied, and the results are not generalisable. Moreover, although the authors claim that "the decision about theophylline in the n of 1 trial group was usually governed by an objective, statistical result," among 18 paired treatment periods the difference in symptom scores was conclusive in only eight, and among four subjects in whom theophylline was continued the 95% confidence interval for the difference in diary scores excluded zero in only one.

A more useful approach would be to improve the ways in which we influence and assess response to treatment. There are good arguments for discarding the placebo since its effect cannot be clearly distinguished from that of either specific treatments² or a range of therapeutic skills.³ Discarding the placebo would make it possible to increase patients' involvement in medical decision making, which itself can lead to improved outcomes.⁴ At the very