- 46 Working Party on Prenatal Diagnosis and Genetic Screening. *Prenatal diagnosis and genetic screening: community and service implications*. London: Royal College of Physicians of London, 1989.
- 47 Anionwu EN. Sickle cell and thalassaemia: some priorities for nursing research. *J Adv Nurs* 1996;23:853-6.
- 48 Shickle D, May A. Knowledge and perceptions of haemoglobinopathy carrier screening among general practitioners in Cardiff. *J Med Genet* 1989;26:109-12.

- 49 Choiseul M, Allen A, May A. Training and needs of health visitors in the haemoglobinopathies. *Health Visitor* 1988;61:205-6.
- 50 King's Fund forum consensus statement: screening for fetal and genetic abnormality. *BMJ* 1987;295:1551-3. (Conference report.)
- 51 Anionwu EN. Ethnic origin of sickle and thalassaemia counsellors: does it matter? In: Kelleher D, Hillier S, eds. *Researching cultural differences in health*. Routledge, 1996:160-89.

(Accepted 16 May 1996)

How can impact factors be improved?

Eugene Garfield

Impact factors are widely used to rank and evaluate journals. They are also often used inappropriately as surrogates in evaluation exercises. The inventor of the *Science Citation Index* warns against the indiscriminate use of these data. Fourteen year cumulative impact data for 10 leading medical journals provide a quantitative indicator of their long term influence. In the final analysis, impact simply reflects the ability of journals and editors to attract the best papers available.

Counting references to rank the use of scientific journals was reported as early as 1927 by Gross and Gross.¹ In 1955 I suggested that reference counting could measure "impact,"² but the term "impact factor" was not used until the publication of the 1961 *Science Citation Index (SCI)* in 1963. This led to a byproduct, *Journal Citation Reports (JCR)*, and a burgeoning literature using bibliometric measures. From 1975 to 1989, *JCR* appeared as supplementary volumes in the annual *SCI*. From 1990-4, they have appeared in microfiche, and in 1995 a CD ROM edition was launched.

Calculation of current impact factors

The most used data in the *JCR* are impact factors—ratios obtained from dividing citations received in one year by papers published in the two previous years. Thus, the 1995 impact factor counts the citations in 1995 journal issues to "items" published in 1993 and 1994. I say "items" advisedly. There are a dozen major categories of editorial matter. *JCR*'s impact calculations are based on original research and review articles, as well as notes. Letters of the type published in the *BMJ* and the *Lancet* are not included in the publication count. The vast majority of research journals do not have such extensive correspondence sections. The effects of these differences in calculating journal impact can be considerable.^{3,4}

Absolute citation counts

The ubiquitous and sometimes misplaced use of journal impact factors for evaluation has caused considerable controversy. They are probably the most widely used of all citation based measures. They were invented to permit

Large journals that publish many papers may not have as high an impact as smaller review journals

reasonable comparison between large and small journals. Absolute citation counts preferentially give highest rank to the largest or the oldest journals. For example, in 1994, articles published in the *BMJ* regardless of age were cited 37 600 times. Of these, 5800 citations—about 15%—were to items published in 1992 and 1993.

Table 1 shows absolute citation counts for nine English language clinical journals as well as the *Journal of Biological Chemistry*, which is included to emphasise the difference between absolute and relative citation of large journals. Table 2 lists these same journals ranked by 1994 impact. It is important to note that of the thousands of journals published and cited in *SCI* only 337 achieved a current impact higher than 3.0. The *SCI* processes over 3300 source journals and cites thousands more, but it does not include as sources hundreds of low impact applied and clinical journals. This has been a source of frustration to editors from the Third World. They often ask how they can improve impact so as to warrant inclusion.

Impact of review journals

It should be apparent from reviewing the two tables that the ranking relationship between quality and citation is not absolute. The *Journal of Biological Chemistry* is one of the most cited journals in the history of science but it is also among the largest. Publishing over 4000 research manuscripts a year (as do other high volume journals like *Physical Review*) inevitably leads to considerable variation in quality and impact of individual articles. As a consequence, while these journals publish many papers which become "citation classics," their current impact may not be as high as some smaller journals, especially certain review journals. Indeed, one of the highest impact journals is *Annual Review of Biochemistry*—with a current impact of 42.2. But as the data for *Annual Review of Medicine*

Institute for Scientific Information, 3600 Market Street, Suite 450, Philadelphia, PA 19104, USA
Eugene Garfield, chairman emeritus
email: garfield@aurora.cis.upenn.edu

BMJ 1996;313:411-3

Table 1—Absolute citation counts, 1994 and 1989

Title	Rank in 1994	Times cited (thousands) in 1994 (from all years)	Rank in 1989	Times cited (thousands) in 1989 (from all years)
<i>Journal of Biological Chemistry</i>	1	265.3	1	183.4
<i>New England Journal of Medicine</i>	10	98.9	9	72.4
<i>Lancet</i>	11	85.5	12	68.0
<i>JAMA</i>	41	41.0	40	29.4
<i>BMJ</i>	45	37.7	34	32.6
<i>Annals of Internal Medicine</i>	51	31.7	48	26.5
<i>American Journal of Medicine</i>	75	22.1	60	21.7
<i>Archives of Internal Medicine</i>	138	13.7	130	10.4
<i>Medicine</i>	502	4.3	421	3.9
<i>Annual Review of Medicine</i>	1118	1.6	999	1.4

Table 2—1994 current impact factors

Journal	1994 impact factor (1992-3 articles)	Citations to 1992-3 items (thousands)
<i>New England Journal of Medicine</i>	22.7	18.4
<i>Lancet</i>	17.3	16.6
<i>Annals of Internal Medicine</i>	9.9	5.4
<i>Journal of Biological Chemistry</i>	7.7	60.2
<i>JAMA</i>	6.9	8.1
<i>BMJ</i>	4.4	5.8
<i>Archives of Internal Medicine</i>	4.1	2.1
<i>Medicine</i>	3.9	0.2
<i>Annual Review of Medicine</i>	2.8	0.2
<i>American Journal of Medicine</i>	2.7	1.7

Table 3—Cumulative impact 1981-94 for papers published from 1981 to 1986

Title	Impact	Papers	Citations
<i>New England Journal of Medicine</i>	115.7	2 289	264 768
<i>Annals of Internal Medicine</i>	67.9	1 741	100 770
<i>Medicine</i>	56.5	193	10 903
<i>Journal of Biological Chemistry</i>	50.8	14 483	735 026
<i>Lancet</i>	50.5	3 544	179 125
<i>American Journal of Medicine</i>	28.2	2 932	82 649
<i>Annual Review of Medicine</i>	25.3	260	6 315
<i>JAMA</i>	23.9	3 326	79 339
<i>BMJ</i>	17.4	5 748	100 154
<i>Archives of Internal Medicine</i>	16.8	2 027	34 078

show, this is not an absolute rule. In general, a journal is well advised to publish authoritative review articles if it would increase its impact. Since there are over 40 000 review articles published each year, not all will achieve high impact. Again, selection of reviews about active research fronts is important, as is their timing. Controversial topics may increase impact. A non-medical example is "cold fusion" by Fleischman and Pons,⁵ which has been cited over 500 times. A recent medical example could be the "Concorde study," already cited in 150 papers.⁶ But hundreds of halfbaked controversial ideas are essentially ignored.

Nothing will replace the judgment necessary for editors to select putative citation classics and to reject trivial or outlandish papers. Nevertheless, most reputable journals have at one time or another rejected papers which proved to be blockbusters. In retrospect, we should congratulate editors who publish controversial ideas such as that of Barry Marshall concerning *Helicobacter pylori* and peptic ulcers.⁷ That paper was well on its way to citation classic status when I nominated Marshall for the John Scott Award of Philadelphia—several years before he received the Albert Lasker award this year.

Method papers

It is widely believed that method papers are cited more than the average and thus increase journal impact. How lovely it would be if every method proved to be another Lowry method,⁸ cited over 8000 times in 1994 and over 250 000 in its lifetime. But the fact is that method journals do not achieve extraordinary impact since the vast majority of their papers, like clinical tests, are not unusual.

An editor could select authors on the basis of past performance. By checking their citation histories, one could undoubtedly increase the probability of publishing papers with higher potential impact. Some editors do this instinctively, especially when publishing the first few issues of a new specialty journal. In most cases, the most-cited papers for newer journals appear in the first volume published.

Over the years, the increase in multiauthored papers has been apparent. This is matched by an increase in multinational and multi-institutional clinical and epidemiological studies. At the Institute for Scientific Information (ISI), unpublished studies support the notion that these papers produce greater impact. ISI's *Science Watch* has regularly reported on the most-cited current papers in medicine. There is fierce competition among editors to publish these "hot papers." These undoubtedly contribute to increased current impact. But what about long term impact?

Cumulative impact factors

In table 3, the 14 year citation impact for 1981-94 for the articles published from 1981 to 1986 in the *BMJ* or the *Lancet* are, in general, even higher than suggested by current impact. *Annals of Internal Medicine* is even stronger on this measure. Certain fields or topics require more time to mature because of delayed recognition (as with *H pylori*) or because of the time required to produce experimental or clinical results. It generally takes more time to achieve impact in dermatology than it does in molecular biology or astrophysics. Invidious comparisons between journals even in different fields of medicine do not take these subtleties into account.

No substitute for judgment

Successful editors and publishers know that in order to improve the editorial quality of journals, there is no substitute for judgment, quality, and relevance. Impact and other citation measures merely report the facts. Authors will gravitate to journals with widespread influence. Circulation alone will not increase research impact. Otherwise, *JAMA* and others would rank higher in impact. But only a fraction of doctors do research. Dissemination of research results to the press may increase general awareness.⁹ But that role in research is played primarily by current awareness services such as *Current Contents*, by contacts at meetings, reprint exchange, and reading primary journals. Last but not least, the way to improve impact is to insist that authors cite all of the relevant literature. Editors should avoid artificial limits on bibliography as long as it is not obviously self serving.

In spite of dozens of presentations by myself and others, there continues to be a certain mystique about journal selection for *Current Contents*, the *Science Citation Index*, and the *Social Sciences Citation Index*. My 1990 essay in *Current Contents* is still sent to those making such enquiries.¹⁰

Of the 4500 journals covered by *SCI* and *SSCI*, probably 3000 can be described as biomedically related. Of these, 500 account for 50% of what is published and 75% of what is cited. Of the 3300 covered in *Medline*, hundreds of low impact journals are not included by ISI for similar reasons—space and economics.

Other factors in journal evaluation

New journals continue to appear each year and must be evaluated as early as possible. But even *Nature* in its periodic reviews of new journals requires the passage of time before accepting journals to be reviewed. An experienced evaluator takes into account timing, format, subject matter, past performance, and other indicators such as internationality. The first issue of many journals is full of hope, but they soon exhaust the backlog of material needed to insure continued, timely publication. Clearly a society publisher with years of experience does not launch a journal without a long term commitment—and its editorial standards will be well known. Inexperienced publishers often do not live up to their rosy expectations. The inclusion of abstracts and complete author, street, and email addresses are but a few artefacts that are factored in the judgment of

minimum quality. ISI may also ask how well a particular specialty or country is represented in its coverage. And, after a few years of history, all other criteria being equal, one can look at a journal's impact.

This article has focused on journal impact factors and their role in what Stephen Lock described as "journalology."¹¹ Like individual authors, a variety of indicators can be used to judge journals in a current or historical sense. Impact numbers are probably less important than the rankings which are obtained. Often there are only slight quantitative differences involved. The literature is replete with recommendations for corrective factors that should be considered, but in the final analysis subjective peer judgment is essential.

Caution in use of impacts as surrogates

Journal impact data have been grafted on to certain large scale studies of university departments and even individuals. Sometimes a journal's impact is used as a substitute for the evaluation of recently published articles simply because it takes several years for the average article to be cited. However, a small percentage of articles will experience almost immediate and high citation. Using the journal's average citation impact instead of the actual article impact is tantamount to grading by the prestige of the journal involved. While expedient, it is dangerous. Although journal assessments are important, evaluation of faculty is a much

more important exercise that affects individual careers. Impact numbers should not be used as surrogates except in unusual circumstances.¹²

- 1 Gross PLK, Gross EM. College libraries and chemical education. *Science* 1927;66:385-9.
- 2 Garfield E. Citation indexes for science: a new dimension in documentation through association of ideas. *Science* 1955;122:108-111.
- 3 Garfield E. Which medical journals have the greatest impact? *Ann Intern Med* 1986;105:313-20.
- 4 Garfield E. Why are the impacts of the leading medical journals so similar and yet so different? Item-by-item audits reveal a diversity of editorial material. In: *Essays of an information scientist*. Vol 10. Philadelphia: ISI Press, 1987:7-13.
- 5 Fleischmann M, Pons S, Hawkins M. Electrochemically induced nuclear fusion of deuterium. *Journal of Electroanalysis Chemistry Interface* 1989;261:301-8.
- 6 Aboulker JP, Smart AM. Preliminary analysis of the Concorde trial. *Lancet* 1993;341:889-90.
- 7 Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* 1984;i:1311-5.
- 8 Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry* 1951;193:265-75.
- 9 Van Trigt AM, de Jong-van den Berg LT, Voogt LM, Willems J, Tromp TF, Haaijer-Ruskamp FM. Setting the agenda: does the medical literature set the agenda for articles about medicines in the newspapers? *Soc Sci Med* 1995;41:893-9.
- 10 Garfield E. How ISI selects journals for coverage: quantitative and qualitative considerations. In: *Essays of an information scientist*. Vol 13. Philadelphia: ISI Press, 1990:185-93.
- 11 Lock SP. "Journalology": are the quotes needed? In: Garfield E. *Essays of an information scientist*. Vol 13. Philadelphia: ISI Press, 1990:19-24.
- 12 Garfield E. How to use citation analysis for faculty evaluations, and when is it relevant? Parts 1 and 2. In: *Essays of an information scientist*. Vol 6. Philadelphia: ISI Press, 1984:354-72.

(Accepted 17 May 1996)

Lesson of the Week

Acute angle closure glaucoma masquerading as systemic illness

Margaret Dayan, Benjamin Turner, Charles McGhee

Acute angle closure glaucoma can present with systemic symptoms such as headache or gastrointestinal disturbance

Acute angle closure glaucoma is a comparatively rare, acute form of glaucoma that may be associated with various systemic symptoms; these can overshadow the primary ocular problem. In some cases this delay in diagnosis may result in blindness. We report on three women who presented to our department with acute angle closure glaucoma during one fortnight. Their initial symptoms were attributed to surgical and psychiatric problems rather than to their ocular disease.

Case reports

CASE 1

A 63 year old hypermetropic (long sighted) woman presented to her general practitioner with a 24 hour history of right sided headache followed, several hours later, by vomiting, diarrhoea, and non-specific abdominal pain. Her symptoms failed to settle and she was referred to hospital and admitted under a surgical team. No cause could be found for her gastrointestinal disturbance, but the following day her right eye was noted to be red; on direct questioning she admitted to a reduction in visual acuity in that eye, which had coincided with the onset of her symptoms. She was referred to the eye accident and emergency department.

The visual acuity in her right eye was reduced to hand movements and associated with a raised intraocular pressure of 44 mm Hg (normal range 10-21 mm Hg), a fixed, mid-dilated pupil, substantial corneal oedema, and a very shallow anterior chamber. Angle closure glaucoma was diagnosed and the intraocular pressure lowered medically. Her systemic symptoms resolved completely. Bilateral laser peripheral iridotomies were

performed on the next day. Her visual acuity improved to 6/12 in the right eye compared with 6/6 in the left. A visual field examination gave normal results six weeks later.

CASE 2

A 63 year old hypermetropic woman was given amitriptyline 25 mg once daily by her general practitioner for symptoms of depression, including fatigue and insomnia. Soon afterwards she developed recurrent evening headaches accompanied by mild blurring of vision, which, she later admitted, were also associated with seeing coloured haloes around bright lights. These symptoms entirely resolved with sleep and were attributed to her depression. Consequently, the dose of amitriptyline was increased to 100 mg at night, with a concurrent increase in the frequency of her symptoms. A month later she developed persistent headache and blurring of vision in the right eye, for which she attended the eye accident and emergency department.

On examination the visual acuity in her right eye was reduced to 6/9 (compared with 6/5 on the left). This was associated with a substantially raised intraocular pressure of 60 mm Hg, a fixed, mid-dilated pupil, moderate corneal oedema, and a very shallow anterior chamber. Acute on intermittent angle closure glaucoma, precipitated by amitriptyline, was diagnosed. The intraocular pressure in the right eye was lowered medically. Bilateral laser peripheral iridotomies were performed on the same day. The visual acuity in her right eye remained at 6/9 and visual field examination showed that the visual field was constricted on the right.

Sunderland Eye Infirmary, Sunderland SR2 9HP
Margaret Dayan, registrar in ophthalmology
Benjamin Turner, senior house officer in ophthalmology

University of Sunderland School of Health Sciences, Sunderland SR2 7EE
Charles McGhee, consultant ophthalmologist

Correspondence to:
Miss M Dayan, Department of Ophthalmology, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP.

BMJ 1996;313:413-5