(N - methyl - 1 - (3,4 - methylenedioxyphenyl) - 2 - butanamine) instead of ecstasy. Mixtures of these substances with ecstasy as well as with unrelated drugs such as amphetamine, caffeine, and l-phenylethylamine are also encountered. In addition, tablets have been found to contain other, less common, ring substituted amphetamines instead of ecstasy. These analogues may well differ in their neurotoxic potential. Finally, there have been reports of mixtures of ketamine and ephedrine, as well as other substitutes such as mixtures of lysergide (LSD) and amphetamine or of lysergide and ephedrine, being sold as ecstasy.

This diverse pharmacological content of tablets may go some way to explaining the variety of symptoms reported among and within users. Not only does it mean that users are exposing themselves to unknown long term effects but it leaves researchers with a problem. Neither the user nor the researcher can be certain what the ingested tablet contained until later analysis. Adverse symptoms, both short and long term, will depend on the interaction between the drug and the individual, which cannot be safely predicted.

ADAM R WINSTOCK
Registrar in psychiatry

Maudsley Hospital,
London SE5 9RS

LESLIE A KING
Head

Drugs Intelligence Laboratory,
Forensic Science Service,
Aldermaston,
Reading RG7 4PN

- 1 Green AR, Goodwin GM. Ecstasy and neurodegeneration. *BMJ* 1996;312:1493-4. (15 June.)
- 2 McGuire PK, Cope H, Fahy TA. Diversity of psychopathology associated with use of 3,4-methylenedioxymethamphetamine ("ecstasy"). *Br J Psychiatry* 1994;165:391-5.
- 3 McCann UD, Ricaurte GA. Lasting neuropsychiatric sequelae of (±) methylenedioxymethamphetamine ("ecstasy") in recreational users. *J Clin Psychopharmacol* 1991;11:302-5.
- 4 Colado MI, William JL, Green AR. The hyperthermic and neurotoxic effects of "ecstasy" (MDMA) and 3,4-methylenedioxymphetamine (MDA) in the dard agouti (DA) rat, a model of the CYP2D6 poor metaboliser phenotype. *Br J Pharmacol* 1995;115:1281-90.

...such as ketamine

EDITOR,—There is increasing concern over the toxic¹ and neurodegenerative² effects of ecstasy, and further research is clearly required. Potential hazards can also, however, arise through use of tablets sold as ecstasy that contain substances other than ecstasy, such as ketamine. Ketamine is an anaesthetic agent with a rapid action that has medical and veterinary applications. Drug users who have taken ketamine illicitly say that it induces a profound change in consciousness, involving a feeling of dissociation from the body, and invokes visual hallucinations. These users have reported the need for precautions when taking the drug, emphasising that the intensity of its effects could lead to problems such as psychological distress or accidents due to lack of physical control, particularly if it was taken in a public setting (such as a club) and if used unwittingly or by novices.³ There have been no reports of any serious toxicological or neurodegenerative problems arising from the illicit use of ketamine.

Until a few years ago illicit use of ketamine mostly entailed small scale diversion of injection ampoules from clinical supplies. Since 1993 in Britain, police seizures of ketamine notified to the Drugs Intelligence Laboratory at Aldermaston have increased considerably, from 21 in 1993 to 183 in 1995. In nearly all cases the ketamine was in tablet form, often resembling tablets commonly sold as ecstasy. The ketamine in these illicit tablets had often been mixed with ephedrine, but some tablets also contained selegiline or procaine. Recently a single police

seizure yielded almost 100 000 tablets containing ketamine and ephedrine, about one third of which carried an impression of a bird ("white dove"). Ecstasy tablets bearing this design have been common for some years.⁴

The recent trend in seizures of ketamine indicates an appreciable increase in the availability of tablets that are intended to mimic the effects of ecstasy and the appearance of ecstasy tablets. While it is important to note that ecstasy has never been found mixed with ketamine, illicit manufacturers seem likely to be targeting a market in which ecstasy is widely available; the potential for the problems described above is therefore considerable. The availability of tablets of ketamine mixed with ephedrine will continue to be monitored, but systematic research on the interactive effects of taking ketamine in conjunction with ephedrine may also be needed.

DAVID SHEWAN
Lecturer in psychology
PHIL DALGARNO
Research associate

Department of Psychology,
Glasgow Caledonian University,
Glasgow G4 0BA

LESLIE A KING
Head

Drugs Intelligence Laboratory,
Forensic Science Service,
Aldermaston,
Reading RG7 4PN

- 1 Green AR, Goodwin GM. Ecstasy and neurodegeneration. *BMJ* 1996;312:1493-4. (15 June.)
- 2 Green AR, Cross AJ, Goodwin GM. Review of the pharmacology of 3,4-methylenedioxymethamphetamine (MDMA or "ecstasy"). *Psychopharmacology* 1995;119:247-60.
- 3 Dalgarno PJ, Shewan D. Illicit use of ketamine in Scotland. *J Psychoactive Drugs* 1996;28:191-9.
- 4 Forsyth AJM. Ecstasy and illegal drug design: a new concept in drug use. *International Journal on Drug Policy* 1995;6:193-209.

γ-Hydroxybutyrate is a new recreational drug that may lead to loss of consciousness

EDITOR,—We were interested in A Richard Green and Guy M Goodwin's editorial on ecstasy and neurodegeneration.¹ We wish to draw attention to another recreational drug, which is taken in similar circumstances to ecstasy, whose use can lead to acute neurological effects.

γ-Hydroxybutyrate—also known as GHB, sodium oxybate, GBH, and liquid ecstasy—has been available in Britain since 1994, mostly in night clubs around London. It has recently become more readily available, and the National Poisons Information Service (London) has received calls regularly since 1994. It produces a state of relaxation and mild euphoria, which may be followed by drowsiness, epileptiform seizures, and coma. We have recently seen several cases of symptomatic toxicity; four patients presented in coma.

γ-Hydroxybutyrate was first reported as a drug of misuse in North America in 1990. That year the Food and Drug Administration banned over the counter sales of it, and in 1991 the Centers for Disease Control issued a warning about its side effects.² In Britain it is not controlled under the Misuse of Drugs Act, and so possession is not considered to be illegal. There have, however, been a few prosecutions under the Medicines Act, which forbids unauthorised manufacture and distribution. The police currently have little power to control the spread of its use, and it will probably become more widely available in Britain.

All accident and emergency staff need to be aware of the use of γ-hydroxybutyrate and its effects and modes of presentation. The drug is commonly sold as a clear, salty liquid in a bottle similar in size to the bottles used for type correction fluid, and the usual dose is one or two cap-

fuls. Side effects may include mild headache, vomiting, and drowsiness, which can be followed by seizures, loss of consciousness, and hyperventilation. Most patients maintain their airway well, and consciousness is usually regained within a few hours, although cases requiring ventilation have been reported. The effects may be worsened by coingestion of alcohol and other psychoactive drugs. The drug is not detectable with routine toxicological tests, and there are no clinically effective antidotes. Management is largely supportive.

Collapse after ingestion of γ-hydroxybutyrate should be added to the list of differential diagnoses for seizures and coma in patients presenting to accident and emergency departments.

IAN M STELL
Lecturer

Accident and Emergency Department,
Guy's Hospital,
London SE1 9RT

JOHN M RYAN
Consultant

Accident and Emergency Department,
Royal Sussex County Hospital,
Brighton BN2 5BE

- 1 Green AR, Goodwin GM. Ecstasy and neurodegeneration. *BMJ* 1996;312:1493-4. (15 June.)
- 2 Centers for Disease Control. Multistate outbreak of poisonings associated with illicit use of gamma hydroxybutyrate. *JAMA* 1991;265:447-8.

Recommendation to avoid all drugs during first trimester is unrealistic

EDITOR,—The message of Patrick Vallance's editorial on drugs and the fetus is rather academic and far from practical experience¹; exaggeration of the teratogenicity of drugs causes anxiety in women of childbearing age and increases the likelihood that doctors will practise defensive medicine.

Drugs are probably regarded as the most important of the supposed hazards during pregnancy. Of about 3000 drugs used in Hungary, only 20 are proved teratogens in humans.² Contrary to supposition, only a small proportion (about 1%) of congenital abnormalities is traceable to teratogenic drugs in humans.³ In 102 women who attempted suicide with extremely large doses of drugs in the first trimester of pregnancy the observed rate of congenital abnormalities was no higher than the expected population figure after women who were dependent on alcohol were excluded.⁴ In a Swedish study no congenital abnormality was found in 70 infants born to mothers who had taken an overdose of psychoactive drugs during the period of fetal development in which the organs form.⁵

In Hungary 91% of pregnant women use drugs during pregnancy³ and 8% need long term drug treatment for chronic disorders. Thus Vallance's recommendation that "all drugs should be avoided if possible during the first trimester" is unrealistic because drug treatment is necessary for many women. Because of inflexible application of this recommendation, however, necessary drug treatment is sometimes not given during pregnancy; this exacerbates disorders such as epilepsy, hypertension, and genitourinary infections. In the early 1980s the number of induced abortions in Hungary was about 80 000, of which about 16% were for medical indications. Analysis indicated, however, that about 5000 pregnancies were terminated for no truly appropriate cause—most were terminated because of the supposed teratogenic effect of drugs and defensive practice by physicians. Vallance mentions griseofulvin as an example of a teratogenic drug taken by men. The dataset of