Ecstasy and neurodegeneration

Advice is that "less is more"

EDITOR,—A Richard Green and Guy M Goodwin rightly point out that the risk of long term consequences of neurodegeneration associated with the use of ecstasy (3,4-methylenedioxymethamphetamine) may be greater than the risk of death from acute toxicity.¹ Having reviewed the compelling evidence linking ecstasy with neurotoxicity in animals, they omit what is perhaps the strongest indication of neurotoxicity in humans.

A universal finding of descriptive studies of users of ecstasy is that the first ecstasy tablet is the best and that with subsequent use the desired effects decline and side effects increase.^{2,4} As ecstasy is pharmacologically active for only a few hours and even frequent users usually limit their use to weekends, neurotoxicity seems a more likely explanation than tolerance through neuroadaptation. Even newsgroups on the Internet that promote the use of ecstasy are now debating not whether it causes serotonergic nerve cell damage but whether the result is detrimental, inconsequential, or even beneficial, with general advice to users being that "less is more" and "more than one tablet once a month is an overdose."

In Britain, use of ecstasy is inextricably related to dance culture and "raves." Perceiving a diminished effect as being due to poor quality, ecstasy "ravers" often "stack up," taking ecstasy in higher doses and more frequently than reported elsewhere. Whereas only a fifth of the Stanford cohort² and a third of the Sydney cohort³ had

taken ecstasy on more than 10 occasions, over half of a sample in London had used it more than 20 times.⁵

Media campaigns focusing on deaths from use of ecstasy are worse than useless: they no more deter young people from taking ecstasy than reports of deaths in an avalanche stop me from taking skiing holidays. Exaggerated, sensationalist drug prevention campaigns are recognised as such by young people and decrease the chance of prudent advice being taken seriously.

JOHN MERRILL
Consultant in drug dependence

Drugs North West,
Manchester M25 7BL

- 1 Green AR, Goodwin GM. Ecstasy and neurodegeneration. *BMJ* 1996;312:1493-4. (15 June.)
- 2 Peroutka SJ, Newman H, Harris H. Subjective effects of 3,4-methylenedioxymethamphetamine in recreational users. *Neuropsychopharmacology* 1988;1:273-7.
- 3 Solowij N, Hall W, Lee N. Recreational use of MDMA in Sydney: a profile of ecstasy users and their experiences with the drug. *Br J Addict* 1992;87:1161-72.
- 4 Greer J, Tolbert R. Subjective reports of the effects of MDMA in a clinical setting. *J Psychoactive Drugs* 1986;18:319-28.
- 5 Winstock AR. Chronic paranoid psychosis after misuse of MDMA. *BMJ* 1991;302:1150-1.

No evidence of neurotoxicity exists

EDITOR,—In their editorial on ecstasy (3,4-methylenedioxymethamphetamine) and neurodegeneration A Richard Green and Guy M Goodwin state that young people who misuse ecstasy should be fully informed of the risks, which they claim are considerable in the long term.¹ The authors cite animal studies indicating that recreational use of ecstasy can cause neurotoxicity to the serotonergic systems of the brain. The fact that the United States Food and Drug Administration recently (in May) approved dexfenfluramine for daily long term use could, however, imply that ecstasy is similarly safe for long term use. This is because much evidence of neurotoxicity, based on markers in animal studies, applies to research on both drugs. In this context G A Ricaurte (whose evidence is cited in the editorial) believes that neurotoxicity produced by dexfenfluramine is identical with that produced by ecstasy (personal communication, October 1995). Indeed, Green himself, with colleagues, has implicitly equated the neurotoxic effects of ecstasy with those produced by fenfluramine.²

Recent research casts further doubt on the methods used to provide evidence of neurotoxicity in humans. Postmortem examination of long term users of methamphetamine showed reduced levels of dopamine nerve terminal markers similar to those seen after use of ecstasy and fenfluramine. But the study went on to conclude that "levels of DOPA decarboxylase and the vesicular monoamine transporter, known to be reduced in Parkinson's disease, were normal. This suggests that chronic exposure to methamphetamine does not cause permanent degeneration of striatal dopamine nerve terminals at the doses used by the young subjects in our study."³

The editorial suggests that use of ecstasy may eventually produce effects similar to Parkinson's disease, as was the case with MPTP (*N*-methyl-4-phenyl-1,2,3,6-tetrahydropyridone). This is unlikely because the effects of MPTP were seen immediately. The analogy with MPTP—used by

those who lobby for the prohibition of ecstasy in the United States—was long ago shown to be unfounded.⁴ The strongest evidence against such damage being a normal consequence of ingestion of ecstasy must surely be the small number of casualties among the large number of users over the past two decades. Psychiatric consequences are generally restricted to people with considerable premorbid psychopathology, poly-substance misuse, and histories of taking excessive amounts of ecstasy.⁵

There is no evidence to show that ecstasy is neurotoxic in humans. Since the possibility of neurotoxicity is a matter of great public concern, however, it should be urgently investigated by research on human volunteers.

NICHOLAS SAUNDERS
Independent researcher

14 Neal's Yard,
London WC2H 9DP
<http://ecstasy.org/>

- 1 Green AR, Goodwin GM. Ecstasy and neurodegeneration. *BMJ* 1996;312:1493-4. (15 June.)
- 2 Colado MI, Murray TK, Green AR. 5-HT loss in rat brain following 3, 4-methylenedioxymethamphetamine (MDMA), p-chloroamphetamine and fenfluramine administration and effects of chloromethiazole and dizocilpine. *Br J Pharmacol* 1993;108:583-9.
- 3 Wilson JM, Kalasinsky KS, Lavery AI, Bergeron C, Reibel G, Anthony RM, et al. Striatal dopamine nerve terminal markers in human, chronic methamphetamine users. *Nature Medicine* 1996;2:699-703.
- 4 Beck J, Morgan P. Designer drug confusion: a focus on MDMA. *J Drug Educ* 1986;16: 267-82.
- 5 McGuire PK, Cope H, Fahy TA. Diversity of psychopathology associated with use of 3,4-methylenedioxymethamphetamine ("Ecstasy"). *Br J Psychiatry* 1994;165:391-5.

Tablets often contain substances in addition to, or instead of, ecstasy...

EDITOR,—A Richard Green and Guy M Goodwin emphasise the potential long term neurotoxicity associated with the use of ecstasy (3,4-methylenedioxymethamphetamine).¹ Small studies of clinical populations have already given a clue to the possible functional impact of this drug on the human nervous system by showing a wide range of psychopathology associated with its short term ingestion.^{2,3} They also point out that people who metabolise the drug quickly, although perhaps protected from acute toxicity (some of which will no doubt be idiosyncratic in nature), may be more at risk of longer term neurodegeneration than other people.

Those who metabolise the drug quickly may possibly also experience "desirable" effects of the drug for a shorter time than other people. This might lead them to take increasing doses to sustain the wanted effect, with this compounding any potential risk. In a study at the Maudsley Hospital many subjects have reported taking over 10 tablets in an evening; some have reported lifetime use exceeding 1000 tablets. With toxicity in animal studies being associated with a single dose,⁴ such lifetime consumption makes the likelihood of latent functional psychological problems even more likely.

Data from the forensic analysis of drugs seized by the police and customs show that tablets that are claimed to be ecstasy tablets may contain a variety of related substances. Tablets are commonly found to contain MDEA (3,4-methylenedioxyethylamphetamine), MDA (3,4-methylenedioxyamphetamine), or MBDB

Advice to authors

We receive more letters than we can publish: we can currently accept only about one third. We prefer short letters that relate to articles published within the past four weeks. Letters received after this deadline stand less chance of acceptance. We also publish some "out of the blue" letters, which usually relate to matters of public policy.

When deciding which letters to publish we favour originality, assertions supported by data or by citation, and a clear prose style. Wit, passion, and personal experience also have their place.

Letters should have fewer than 400 words (please give a word count) and no more than five references (including one to the *BMJ* article to which they relate); references should be in the Vancouver style. We welcome pictures.

Letters should be typed and signed by each author, and each author's current appointment and address should be stated. We encourage you to declare any conflict of interest.

Please enclose a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.