Key messages

- Cosmetic surgery is one of the few areas of health care to have been explicitly rationed in the NHS
- The predominant reason for a breast reduction referral was physical, although social, psychological, and practical reasons also had a role
- The health status of breast reduction patients differed significantly from that of the general female population before and after treatment
- The patients who underwent surgery experienced substantial improvements in physical, social, and psychological functioning
- The study shows the need for and benefits of breast reduction surgery and provides supporting evidence for including this form of plastic surgery in purchasing contracts

the procedure. Many women report feeling embarrassed, unattractive, and self conscious owing to the inordinate size of their breasts and the unwanted attention the size attracts. Surgery also addressed this aesthetic aspect of the problem, resulting in moderate change in social and psychological function and large change in self esteem.

The medical community now considers patients' subjective experience of treatment to be of primary importance when assessing outcomes.¹⁹ When asked directly, patients said that they saw a great deal of change in their appearance, reported that the change was what they wanted, and assessed the surgical result as being either excellent or very good.

It is increasingly argued that health authorities should decide which services to include and exclude from contracts informed by evidence of health outcomes, on the basis of quality of life as well as survival. The services most frequently excluded from purchasing contracts include plastic surgery undertaken for cosmetic purposes, removal of tattoos, reversal of sterilisation, and certain infertility treatments. The savings that these exclusions represent are of marginal significance in terms of their impact on overall NHS resources.¹³ This study has shown substantial improvements in the wellbeing of breast reduction patients. Such evidence challenges the basis for resource

allocation decisions with regard to breast reduction surgery, and perhaps therefore other cosmetic surgery.

We are very grateful for the help and support given to us by the plastic surgeons, their patients, and the plastic surgery staff at Wexham Park Hospital in Slough, the Radcliffe Infirmary in Oxford, and Stoke Mandeville Hospital in Aylesbury.

Funding: Anglia and Oxford Regional Health Authority.

Conflict of interest: None.

- 1 Department of Health. *Hospital Episode Statistics*. London: HMSO, 1994.
- 2 Soderlund N, Raftery J, Milne R, Gray A. *Cost weights and casemix index construction for healthcare related groups*. Winchester: National Casemix Office, NHS Executive, 1995.
- 3 Klein R, Redmayne S. *Patterns of priorities*. Birmingham: National Association of Hospitals and Trusts, 1992. (Research paper No 7.)
- 4 Bowling A, Jacobson B, Southgate L. Explorations in consultations of the public and health professionals on priority setting in an inner London health district. *Soc Sci Med* 1993;37:851-7.
- 5 Cole RP, Shakespeare V, Shakespeare P, Hobby JA. Measuring outcome in low-priority plastic surgery patients using Quality of Life indices. *Br J Plast Surg* 1994;47:117-21.
- 6 Jenkinson C, Wright L, Coulter A. *Quality of life measurement in health care. A review of measures and population norms for the UK SF-36*. Oxford: Health Services Research Unit, 1993.
- 7 Jenkinson C, Coulter A, Wright L. Short form 36 (SF 36) health survey questionnaire: normative data for adults of working age. *BMJ* 1993;306:1437-40.
- 8 Goldberg DP, Hillier VF. A scaled version of the general health questionnaire. *Psych Med* 1979;9:139-45.
- 9 Rosenberg M. *Society and the adolescent self image*. Princeton, NJ: Princeton University Press, 1965.
- 10 Wright L, Harwood D, Coulter A. *Health and lifestyles in the Oxford Region*. Oxford: Health Services Research Unit, 1992.
- 11 Kazis LE, Anderson JJ, Meenan RF. Effect sizes for interpreting change in health status. *Med Care* 1989;27(suppl):S178-89.
- 12 Cohen J. *Statistical power analysis for the behavioural sciences*. New York: Academic Press, 1977.
- 13 House of Commons Health Committee. *Priority setting in the NHS: purchasing. First report sessions 1994-5*. London: HMSO, 1995. (HC 134-1.)
- 14 Deyo R, Patrick D. The significance of treatment effects: the clinical perspective. *Med Care* 1995;33:AS286-91.
- 15 Garratt AM, Ruta DA, Abdalla MI, Buckingham JK, Russell IT. The SF36 health survey questionnaire: an outcome measure suitable for routine use in the NHS? *BMJ* 1993;306:1440-4.
- 16 Fitzpatrick R, Ziebland S, Jenkinson C, Mowat A, Mowat A. Importance of sensitivity to change as a criterion for selecting health status measures. *Quality in Health Care* 1992;1:89-93.
- 17 Bardsley M, Venables C, Watson J. Evidence for validity of a health status measure in assessing short term outcomes of cholecystectomy. *Quality in Health Care* 1992;1:10-4.
- 18 Jenkinson C, Lawrence K, McWhinnie D, Gordon J. Sensitivity to change of health status measures in a randomised controlled trial: comparison of the COOP Charts and the SF-36. *Quality of Life Research* 1995;4:47-52.
- 19 Ellwood PM. The Shattuck lecture. Outcome management: a technology of patient experience. *N Engl J Med* 1988;318:1549-56.

(Accepted 11 June)

Incidence of myocardial infarction in elderly men being treated with antihypertensive drugs: population based cohort study

Juan Merlo, Jonas Ranstam, Hans Liedholm, Bo Hedblad, Gunnar Lindberg, Ulf Lindblad, Sven-Olof Isacson, Arne Melander, Lennart Råstam

Abstract

Objective—To analyse the association between use of antihypertensive treatment, diastolic blood pressure, and long term incidence of ischaemic cardiac events in elderly men.

Design—Population based cohort study. Baseline examination in 1982-3 and follow up for up to 10 years.

Setting—Malmö, Sweden.

Subjects—484 randomly selected men born in 1914 and living in Malmö during 1982.

Main outcome measures—Observational comparisons of incidence rates and rate and hazard ratios of ischaemic cardiac events (myocardial infarction or death due to chronic ischaemic cardiac disease).

Results—The crude incidence rate of ischaemic cardiac events was higher in those subjects who were taking antihypertensive drugs than in those who were not (rate ratio 2.6 (95% confidence interval 1.7 to 3.9)). After adjustment for potential confounders (differences in baseline smoking

habits, blood pressure, time since diagnosis of hypertension, ischaemic or other cardiovascular disease, hypercholesterolaemia, hypertriglyceridaemia, diabetes mellitus, obesity, and raised serum creatinine concentration) this rate was reduced but still raised (hazard ratio 1.9 (1.0 to 3.7)). In men with diastolic blood pressure >90 mm Hg, antihypertensive treatment was associated with a twofold increase in the incidence of ischaemic cardiac events (rate ratio 2.0 (1.1 to 3.6)), which vanished after adjustment for potential confounders (hazard ratio 1.1 (0.5 to 2.6)). In those subjects with diastolic blood pressure ≤90 mm Hg, antihypertensive treatment was associated with fourfold increase in incidence (rate ratio 3.9 (2.1 to 7.1)), which remained after adjustment for potential confounders (hazard ratio 3.8 (1.3 to 11.0)).

Conclusion—Antihypertensive treatment may increase the risk of myocardial infarction in elderly men with treated diastolic blood pressures ≤90 mm Hg.

Correspondence to:
Dr Merlo.

BMJ 1996;313:457-61

**Department of
Community Medicine,
Lund University, Malmö
University Hospital,
S-205 02 Malmö, Sweden**
Hans Liedholm, *senior
registrar*
Bo Hedblad, *epidemiologist*
Sven-Olof Isacson, *professor*
Lennart Råstam, *professor*

**NEPI Foundation
(Swedish Network for
Pharmacoepidemiology),
Malmö and Stockholm,
Sweden**
Jonas Rånstam, *biostatistician*
Gunnar Lindberg, *clinical
epidemiologist*
Arne Melander, *professor*

**Skaraborg Institute,
Skövde, Sweden**
Ulf Lindblad, *research fellow*

Introduction

A major aim of antihypertensive treatment is to reduce the risk of myocardial infarction, but its ability to achieve this in clinical trials is modest,¹⁻³ and its effectiveness in reducing the incidence of myocardial infarction in the general population is questionable.⁴⁻⁶ Indeed, several investigations indicate that antihypertensive treatment may increase the risk of myocardial infarction,⁷⁻⁸ at least in subjects whose myocardium is hypertrophic or ischaemic.⁹ This may relate to critical reduction of diastolic blood pressure.⁷⁻⁹ Recent studies have also shown increased cardiovascular mortality in elderly men taking antihypertensive drugs who have reduced diastolic blood pressure.¹⁰ For these reasons, we investigated the long term incidence of ischaemic cardiac events in relation to antihypertensive treatment and diastolic blood pressure in a population based cohort of elderly men.

Subjects and methods

STUDY POPULATION

"Men born in 1914" is a prospective population study with the aim of investigating determinant factors for cardiovascular and respiratory diseases. It was carried out in the city of Malmö, Sweden, and a first examination was performed during 1969-70.¹¹ The cohort was redefined in 1982-3, when the men were 68 years old, and a baseline cross sectional study was carried out. Five hundred of the 621 men (80.5%) born in the even months of 1914 who were living in Malmö during 1982-3 participated in this examination. Information on both blood pressure and antihypertensive treatment was complete in 484 of the participants, and these men constituted the current study population.

EXAMINATION

The health examination began with a structured interview at home that included questions about medical history and drug treatment. A comprehensive evaluation of cardiovascular history and status was made at the department of community medicine and at the laboratory of clinical physiology at Malmö University Hospital, a week later. The procedure has been described in detail elsewhere.¹²

Casual systolic and diastolic (phase V) blood pressures were recorded to the nearest 5 mm Hg. The subjects were asked to abstain from smoking on the morning of the test day, and the measurements were made on the right arm, after 15 minutes of supine rest, with a mercury manometer and a rubber cuff (12 × 35 cm). The presence of angina and intermittent claudication was assessed by questionnaire,¹³ and episodes of ST segment depression were assessed by 24 hour electrocardiographic ambulatory registration (Holter monitoring).¹⁴ Ischaemic heart disease was defined as presence of angina or occurrences of ischaemic type ST segment depression.

Blood samples were drawn after a minimum fasting period of 8 hours, and concentrations of serum total cholesterol, serum triglyceride, serum creatinine, and blood glucose were determined at the department of clinical chemistry at Malmö University Hospital. Hypercholesterolaemia was defined as a serum cholesterol concentration ≥ 6.5 mmol/l, and hypertriglyceridaemia as a serum triglyceride concentration ≥ 2.3 mmol/l.¹⁵ Creatinine concentration was classed as high when it was equal to or higher than the 95th centile of the distribution of the participants.

Body mass index was calculated as weight (kg)/(height (m))². Obesity was defined as a body mass index >28 .¹⁶ Diabetes was diagnosed when subjects were receiving drug treatment for the disease or had a fasting blood glucose concentration ≥ 7.0 mmol/l.

INTERVIEW

Data on current drug treatment, years since diagnosis of hypertension, and smoking were assessed by a structured questionnaire, which was filled in during the home interview. Subjects reporting continuous use of antihypertensive drugs were asked to show the packages of prescribed drugs to the research assistant, who recorded the trade names. Reported duration of hypertension was arbitrarily dichotomised into <5 years and ≥ 5 years. The men were classed as current smokers, former smokers (those who had stopped smoking at least one month before investigation), and those who had never smoked. Survey data on previous admission to hospital for myocardial infarction or cerebrovascular events (transient ischaemic attacks or stroke) were validated against medical records.

FOLLOW UP

The men were followed up from baseline in 1982-3 until 31 December 1992, or until death or the occurrence of an ischaemic cardiac event. Fatal and non-fatal ischaemic cardiac events were identified according to the procedure used for the Malmö heart infarction register.¹⁷ All events were validated against medical records or necropsy reports. An ischaemic cardiac event was defined as either a myocardial infarction (ICD-8 code 410 (international classification of diseases, eighth revision)) or death due to chronic ischaemic cardiac disease (ICD-8 code 412). In men with more than one ischaemic cardiac event only the first event was used for analysis. Information on deaths outside hospital was obtained from the national mortality register.

STATISTICAL METHODS

The crude incidence, the rate ratio, and Kaplan-Meier estimates were calculated to compare the incidence of ischaemic cardiac events between the men who took antihypertensive drugs and those who did not. Proportional hazards models were used to estimate the hazard ratio of ischaemic cardiac events among users and non-users of antihypertensive drugs after adjustment for potential confounders—systolic blood pressure, diastolic blood pressure (except when the analysis included stratification on diastolic blood pressure), presence or absence of previous myocardial infarction, other ischaemic heart disease, intermittent claudication, cerebrovascular events, hypercholesterolaemia, hypertriglyceridaemia, elevated serum creatinine concentration, diabetes mellitus, obesity, a history of hypertension of ≥ 5 years, use of cardiac glycosides, and history of smoking. All confidence intervals were calculated at the 95% level.

Results

Of the 484 men in the present study, 395 (82%) had also participated in the initial examination in 1969-70.¹¹ The prevalence of antihypertensive treatment was then only 3%. Mean blood pressures in 1969-70 were 156/93 mm Hg in those who were taking antihypertensive drugs in 1982-3 and 131/81 mm Hg in those who were not. Men taking antihypertensive drugs in 1982-3 had almost identical mean blood pressures if they had had cardiovascular disease before 1982-3 (157/93 mm Hg) or if they had not (156/92 mm Hg). Cardiovascular disease was uncommon in 1969-70 but seemed to be more common among those who took antihypertensive drugs in 1982-3 (11%) than among those who did not (6%). However, of those men who had had cardiovascular disease in or before 1969-70 and who were taking antihypertensive drugs in 1982-3, only two experienced an ischaemic cardiac event during follow up.

Table 1 shows the baseline characteristics of the 484 men in this study. Of these, 113 (23%) were taking anti-

Table 1—Characteristics of 484 elderly men at baseline by use of antihypertensive drugs. Values are numbers (percentages)

	All subjects (n = 484)	Use of antihypertensive drugs	
		Yes	No
Previous myocardial infarction	40 (8)	13 (12)	27 (7)
Ischaemic heart disease*	149 (31)	46 (41)	103 (28)
Intermittent claudication	17 (4)	5 (5)	12 (3)
Previous cerebrovascular event*	29 (6)	16 (14)	13 (6)
Moderate or severe hypertension†	106 (22)	35 (31)	71 (19)
Duration of hypertension ≥5 years	115 (24)	83 (74)	32 (9)
Diabetes mellitus*	25 (5)	11 (10)	14 (4)
Obesity*	91 (19)	23 (20)	68 (18)
Hypercholesterolaemia*	140 (29)	36 (32)	104 (28)
Hypertriglyceridaemia*	55 (11)	18 (16)	37 (10)
Current smoker	167 (35)	26 (23)	141 (38)
Former smoker	221 (46)	63 (56)	158 (43)
Raised creatinine concentration‡	25 (5)	14 (13)	11 (3)

*See methods section for details.

†Systolic blood pressure ≥180 mm Hg or diastolic pressure ≥105 mm Hg.

‡Serum concentration ≥133 µmol/l.

Table 2—Crude incidence rates and rate ratios of ischaemic cardiac events among 484 elderly men in relation to potential risk factors

	No of events	Person years	Incidence rate*	Rate ratio (95% confidence interval)
Use of antihypertensive drugs:				
No	51	3069	16.6	1.0
Yes	33	778	42.4	2.6 (1.7 to 3.9)
Systolic blood pressure (mm Hg):				
<160	55	2675	20.6	1.0
≥160	29	1171	24.8	1.2 (0.8 to 1.9)
Diastolic blood pressure (mm Hg):				
≤90	44	2175	20.2	1.0
>90	40	1671	23.9	1.2 (0.7 to 1.8)
Serum cholesterol concentration (mmol/l):				
<6.5	56	2732	20.5	1.0
≥6.5	28	1115	25.1	1.2 (0.8 to 1.9)
Serum triglyceride concentration (mmol/l):				
<2.5	73	3403	21.5	1.0
≥2.5	10	441	22.7	1.1 (0.6 to 2.1)
Tobacco smoking:				
Never	9	757	11.9	1.0
Ever	75	3080	24.4	2.0 (1.0 to 4.0)
Body mass index (kg/m ²):				
<28	65	3144	20.7	1.0
≥28	18	688	26.2	1.3 (0.8 to 2.1)
Diabetes:				
No	75	3682	20.4	1.0
Yes	9	165	54.5	2.7 (1.3 to 5.2)
Previous myocardial infarction:				
No	66	3585	18.4	1.0
Yes	18	262	68.7	3.7 (2.3 to 6.1)
Previous stroke or transient ischaemic attacks:				
No	71	3687	19.3	1.0
Yes	13	159	81.8	4.2 (2.5 to 7.3)
Other risk factors‡:				
No	34	2434	14.0	1.0
Yes	49	1404	34.9	2.5 (1.6 to 3.8)

*No of events per 1000 person years.

†Angina pectoris, intermittent claudication, episodes of ST segment depression, or use of cardiac glycosides.

hypertensive drugs: 68 used one drug, 39 used two, and six used three drugs. The antihypertensive drugs used were diuretics (mainly thiazides, 72 men), β blockers (64 men), hydralazine (18), nifedipine (2), bethanidine (2), prazosin (2), methyldopa (2), and reserpine (1). The men taking antihypertensive drugs had a higher prevalence of hypertension, previous cerebrovascular and cardiovascular events, ischaemic heart disease, diabetes mellitus, hypertriglyceridaemia, raised serum creatinine concentrations, and former smoking. The total follow up time was 3842 person years, and the mean follow up was 8 person years. No subject was lost to follow up.

Table 2 shows the crude incidence rates and rate ratios of ischaemic cardiac events in relation to several potential risk factors. Figure 1 shows the occurrence of ischaemic cardiac events during follow up among men who were taking antihypertensive drugs and among those who were not.

Table 3 shows the rate ratios of ischaemic cardiac events and the hazard ratios after adjustment for differences in risk factors and with stratification by diastolic blood pressure. The crude incidence rate of ischaemic cardiac events was higher in the men taking antihypertensive drugs than in those who were not (rate ratio 2.6 (95% confidence interval 1.7 to 3.9)) and differed only marginally by type of drug taken (rate ratio for use of β blockers *v* non-use, 2.2 (1.3 to 3.8); for use of diuretics, 2.6 (1.6 to 4.3); for use of other antihypertensive drugs, 2.4 (1.1 to 5.2)). After adjustment for potential confounders (smoking habits, systolic and diastolic blood pressure, years since diagnosis of hypertension, and prevalence of ischaemic or other cardiovascular or cerebrovascular disease, hypercholesterolaemia, hypertriglyceridaemia, diabetes mellitus, obesity, and raised creatinine concentrations) the rate was reduced but still raised (hazard ratio 1.9 (1.0 to 3.7)).

This analysis was then repeated with stratification by diastolic blood pressure (table 3). Among men with diastolic blood pressure >90 mm Hg the use of antihypertensive drugs was associated with a twofold increase in incidence rate (rate ratio 2.0 (1.1 to 3.6)), but this increase vanished after adjustment for potential confounders (hazard ratio 1.1 (0.5 to 2.6)). Among men with a diastolic blood pressure ≤90 mm Hg the use of antihypertensive drugs was associated with a fourfold increase in the incidence rate (rate ratio 3.9 (2.1 to 7.1)), which remained after adjustment for potential confounders (hazard ratio 3.8 (1.3 to 11.0)).

Discussion

In this population based cohort study of elderly men the use of antihypertensive drugs was associated with an increased incidence of ischaemic cardiac events. However, after adjustment for confounders the twofold increase in risk among those with a diastolic pressure above 90 mm Hg disappeared, whereas the fourfold increased risk among those with a pressure of 90 mm Hg or below remained.

Fig 1—Event free survival of 113 men who were taking antihypertensive drugs and 371 men who were not

Table 3—Incidence of ischaemic cardiac events by use of antihypertensive drugs and diastolic blood pressure in 484 elderly men

Category	No of events	Person years	Incidence rate (95% confidence interval)*	Rate ratio (95% confidence interval)	Hazard ratio (95% confidence interval)†
All subjects					
Not taking antihypertensive drugs (n = 371)	51	3064	17 (9 to 25)	1.0	1.0
Taking antihypertensive drugs (n = 113)	33	778	42 (29 to 55)	2.6 (1.7 to 3.9)	1.9 (1.0 to 3.7)
Total (n = 484)	84	3842	22 (13 to 31)		
Diastolic blood pressure ≤90 mm Hg					
Not taking antihypertensive drugs (n = 231)	31	1957	16 (8 to 24)	1.0	1.0
Taking antihypertensive drugs (n = 36)	13	212	61 (45 to 76)	3.9 (2.1 to 7.1)	3.8 (1.3 to 11.0)
Total (n = 267)	44	2169	20 (11 to 29)		
Diastolic blood pressure >90 mm Hg					
Not taking antihypertensive drugs (n = 140)	20	1107	18 (10 to 26)	1.0	1.0
Taking antihypertensive drugs (n = 77)	20	556	35 (23 to 47)	2.0 (1.1 to 3.6)	1.1 (0.5 to 2.6)
Total (n = 217)	40	1663	24 (14 to 34)		

*No of events per 1000 person years

†Adjusted for differences in previous myocardial infarction, previous cerebrovascular disease, ischaemic heart disease, intermittent claudication, diabetes mellitus, systolic blood pressure, duration of hypertension, hypercholesterolaemia, hypertriglyceridaemia, raised serum creatinine concentration, obesity, use of cardiac glycosides, and history of smoking (and, for the analysis of all subjects, diastolic blood pressure). In the adjustment analyses seven men were excluded because of incomplete information.

RELIABILITY OF STUDY

Observational non-randomised studies usually have various weaknesses, and studies of patients treated for hypertension may have problems in assessing the occurrence of non-fatal events and the extensive follow up necessary for an appropriate evaluation. These arguments are particularly pertinent as our findings are at variance with common opinion on the benefit of antihypertensive treatment. On the other hand, results similar to ours have been obtained in other studies.⁴⁻¹⁰

As the study population constituted a random sample from the total population and as the participation rate was high, it is not likely that selection problems invalidate the findings. Furthermore, an investigation on the characteristics of non-participants has confirmed the representativeness of the study population.¹²

The study was performed in an urban area with only one general hospital, and this facilitated the assessment of occurred non-fatal events. Moreover, computerised systems for registration of myocardial infarctions and causes of death in the population have long been established and the number of necropsies performed during follow up was high. Hence, misclassification of cardiac events is unlikely.

All the men were personally interviewed about exposure to risk factors and drug treatments at baseline, and they were all examined by a physician. The reliability of this information was high when investigated in a separate study.¹⁷ Hence, major misclassification of risk factors is unlikely. Also, a large number of potential confounding factors have been accounted for in the analysis, including differences in smoking habits, systolic blood pressure, the duration of hypertension, and the prevalence of cardiovascular and cerebrovascular disease, diabetes mellitus, hyperlipidaemia, and raised creatinine concentrations.

In order to reduce the problems associated with subgroup analysis, the division of diastolic blood pressure at 90 mm Hg was based solely on the World Health Organisation's definition of hypertension.¹⁸

EXPLANATION OF RESULTS

The higher incidence of ischaemic cardiac events in men with diastolic blood pressure ≤90 mm Hg might reflect severity bias or confounding by indication—that is, that men with previous myocardial infarction or cerebrovascular disease and other high risk subjects were

treated more aggressively with antihypertensive drugs and consequently had a lower diastolic blood pressure. This seems unlikely, however, since our adjustments accounted for possible differences in indications for treatment at baseline (these adjustments seem credible as they eliminated the increased risk in men with diastolic blood pressure >90 mm Hg). In addition, the data from the physical examinations of the subjects 13-14 years earlier (in 1969-70) do not suggest that severity bias could explain the findings.¹¹

The main antihypertensive drugs used in this study were diuretics (mainly thiazides) and β blockers. It is often claimed that metabolic side effects may antagonise the cardioprotective purpose of these drugs, but metabolic side effects of thiazides are negligible at the low doses used in Sweden during the time of follow up.¹⁹ Furthermore, differences in hypercholesterolaemia, hypertriglyceridaemia, and diabetes mellitus at baseline were accounted for in the adjustments. As the incidence of ischaemic cardiac events was higher in those with lower blood pressure, poor compliance with treatment does not seem to be a reasonable explanation.

A plausible explanation for the increased incidence of ischaemic cardiac events in men taking antihypertensive drugs whose diastolic blood pressure was ≤90 mm Hg is that it was confined to a subgroup of subjects in whom lowering of diastolic blood pressure could be detrimental. Patients with compromised coronary flow reserve could be such a group.⁸ This would be compatible with the idea of a J shaped curve for diastolic blood pressure in relation to risk of myocardial infarction—that is, that lowering blood pressure below an optimal level increases the risk.⁹

This study was presented in part at the 11th congress of the International Society of Pharmacoepidemiology, Montreal, 27-30 August 1995, and the first congress of the European Association of Clinical Pharmacology and Therapeutics, Paris, 27-30 September 1995.

Funding: The study was supported by the NEPI Foundation; by grants from the National Corporation of Pharmacies' (Apoteksbolaget's) fund for research and studies in health economics and social pharmacy, the Bank of Sweden Tercentenary Foundation, the Swedish Heart and Lung Foundation, and the Medical Faculty of Lund University; and by the National Institute for Public Health.

Conflict of interest: None.

Key messages

- "The lower the better" is the predominant view in treating high blood pressure, but some studies have indicated that lowering diastolic pressure below a certain level may promote myocardial infarction
- In our 10 year follow up of 484 elderly men the risk of an ischaemic cardiac event was higher in men who were taking antihypertensive drugs than in those who were not
- Among men with diastolic blood pressure >90 mm Hg, the risk was increased twofold but disappeared when adjustments were made for other cardiovascular risk factors
- Among those with diastolic blood pressure ≤90 mm Hg, the risk associated with taking antihypertensive drugs was four times higher and remained after adjustment for other cardiovascular risk factors
- These findings support the concept of a J shaped curve for risk of myocardial infarction in relation to treated diastolic blood pressure

- 1 Cutler JA, MacMahon SW, Furberg CD. Controlled clinical trials of drug treatment for hypertension. *Hypertension* 1989;13(suppl 5):136-44.
- 2 Collins R, Peto R. Antihypertensive drug therapy: effects on stroke and coronary heart disease. In: Swales JD, ed. *Textbook of hypertension*. Oxford: Blackwell, 1994:1156-64.
- 3 Kaplan NM. Meta-analysis of hypertension treatment trials. *Lancet* 1990;335:1093.
- 4 Thürmmer HL, Lund-Larsen PG, Tverdal A. Is blood pressure treatment as effective in a population setting as in controlled trials? Results from a prospective study. *J Hypertens* 1994;12:481-90.
- 5 Clausen J, Jensen G. Blood pressure and mortality: an epidemiological survey with 10 years follow-up. *J Hum Hypertens* 1992;6:53-9.

- 6 Yano K, McGee D, Reed D. The impact of elevated blood pressure upon 10-year mortality among Japanese men in Hawaii: the Honolulu Heart Program. *J Chronic Dis* 1983;36:569-79.
- 7 Cruickshank JM, Thorpe JM, Zacharias FJ. Benefits and potential harm of lowering high blood pressure. *Lancet* 1987;i:581-4.
- 8 Farnett L, Mulrow CD, Linn WD, Lucey CR, Tuley MR. The J-curve phenomenon and the treatment of hypertension. Is there a point beyond which pressure reduction is dangerous? *JAMA* 1991;265:489-95.
- 9 Lindblad U, Råstam L, Rydén L, Ranstam J, Isacson SO, Berglund G. Control of blood pressure and risk of first acute myocardial infarction: Skaraborg hypertension project. *BMJ* 1994;308:681-6.
- 10 Langer RD, Criqui M, Barret-Connor E, Klauber M, Ganiats T. Blood pressure change and survival after age 74. *Hypertension* 1993;22:551-9.
- 11 Isacson S-O. Venous occlusion plethysmography in 55-year-old men. A population study in Malmö, Sweden [dissertation]. *Acta Med Scand* 1972;537(suppl).
- 12 Janzon L, Hanson BS, Isacson S-O, Lindell S-E, Steen B. Factors influencing participation in health surveys: results from the prospective study "Men born in 1914" in Malmö, Sweden. *J Epidemiol Community Health* 1986;40:174-7.
- 13 Rose GA, Blackburn H. *Cardiovascular survey methods*. Geneva: WHO, 1968.
- 14 Hedblad B, Janzon L. Hypertension and ST segment depression during ambulatory electrocardiograph recording. Results from the prospective study "Men born in 1914" from Malmö, Sweden. *Hypertension* 1992;20:32-7.
- 15 Treatment of hyperlipidaemia. *Information from the Medical Products Agency, Sweden* 1995;3:178-87. (In Swedish.)
- 16 Waaler HT. Hazard of obesity—the Norwegian experience. *Acta Med Scand* 1988;723:17-21.
- 17 Hanson BS, Östergren P-O. Different social network and social support characteristics, nervous problems and insomnia: theoretical and methodological aspects on some results from the population study "Men born in 1914, Malmö, Sweden." *Soc Sci Med* 1987;25:849-59.
- 18 Zanchetti A, Chalmers JP, Arakawa K, Gyarfas I, Hamer P, Hansson L, et al. The 1993 guidelines for the management of mild hypertension: memorandum from a WHO/ISH meeting. *Blood Pressure* 1993;2:86-100.
- 19 Cushman WC. Optimising diuretic therapy in elderly patients with hypertension. *Drugs Aging* 1995;7:88-96.

(Accepted 17 June 1996)

Prevalence and determinants of antibodies to hepatitis C virus and markers for hepatitis B virus infection in patients with HIV infection in Aquitaine

F Saillour, F Dabis, M Dupon, D Lacoste, P Trimoulet, P Rispal, E Monlun, J-M Ragnaud, P Morlat, J-L Pellegrin, H Fleury, P Couzigou for the Groupe d'Epidémiologie Clinique du SIDA en Aquitaine

Abstract

Objective—To evaluate the prevalence of antibodies to hepatitis C virus and serological markers for hepatitis B virus infection in patients with HIV.

Design—Cross sectional survey.

Setting—Aquitaine, southwestern France, 1991-94.

Subjects—1935 HIV positive patients seen at least once since June 1991.

Main outcome measures—Presence of antibodies to hepatitis C virus were detected by second or third generation enzyme linked immunosorbent assay (ELISA) and recombinant immunoblot assay (RIBA) and markers for hepatitis B virus detected by ELISA.

Results—The prevalence was 42.5% (823) for antibodies to hepatitis C virus, 56.4 (507) for antibodies to hepatitis B core antigen, 6.9% (133) for hepatitis B surface antigen, 30.2% (584) for antibodies to hepatitis B core and surface antigen with no detectable surface antigen, 26.2% (507) for antibodies to core antigen only, and 4.8% (92) for antibodies to surface antigen only. The prevalence of antibodies to hepatitis C virus was 86.1% (726/843) in subjects who had bloodborne HIV infection and 7.3% (66/899) in those with sexually acquired infection. The prevalence of markers for hepatitis B was higher among homosexuals than in the other groups of patients, except for antibodies to surface antigen alone. The relation between markers for hepatitis B and hepatitis C virus was negative among men but positive among women.

Conclusions—The results favour the hypothesis that hepatitis C virus is sexually transmitted much less commonly than either HIV or hepatitis B virus.

Introduction

Hepatitis C virus was quickly identified as the agent causing most cases of non-A non-B hepatitis after transfusion.¹ Second and third generation enzyme linked immunosorbent assays (ELISA) and recombinant immunoblot assays (RIBA) detecting hepatitis C virus antibody are reliable tools for diagnosing infection.² We provide an estimate of the prevalence of antibodies to hepatitis C virus and serological markers of hepatitis B virus infection in a representative sample of patients with HIV infection in Aquitaine, southwestern France.

Patients and methods

In 1987 the Groupe d'Epidémiologie Clinique du SIDA en Aquitaine started a surveillance system of HIV infection which allowed the creation of the Aquitaine cohort described elsewhere.³ Our study population comprised 2957 HIV positive patients from Aquitaine who were recruited or followed up at least once between June 1991 and September 1994. The start date was chosen because second generation tests for hepatitis C virus were introduced at this time.

We routinely used two ELISAs for detecting hepatitis C virus antibodies (Ortho Diagnostic and Diagnostics Pasteur). If the two results did not agree a RIBA test (Chiron RIBA HCV) was used. The criterion for posi-

Unité INSERM No 330,
Université de Bordeaux II,
33076 Bordeaux Cedex,
France
F Saillour, resident in public
health
F Dabis, professor of
epidemiology

Correspondence and
requests for reprints to:
Professor Dabis.

BMJ 1996;313:461-4