cancer units in district general hospitals.¹ It refers to the number of patients with common cancers whom surgeons need to treat to ensure adequate skill and suggests that figures regarded as adequate now will be revised upwards in the future.

At present no firm recommendations exist about the critical number of patients who are required to ensure adequate clinical skill, though 50-80 per specialist has been suggested by the British Association of Surgical Oncology and the British Breast Group. The only evidence currently available suggests that outcome is better when the annual caseload per consultant exceeds 30 compared with under 10.² Evidence from the same source showed a slightly worse (though not significant) outcome, with fewer patients receiving adjuvant treatments, when the caseload exceeded 50, but this has received no attention. The study concluded that improved survival could be explained by the introduction of a multidisciplinary approach to treatment and that the effect of patient numbers was probably a function of this and of clinical organisation.

A recent paper suggested that the prognosis of patients with breast cancer is improved by specialist care.³ A specialist interest was defined on the basis of four criteria: having a dedicated breast clinic; having a defined association with pathologists and oncologists, organising and facilitating clinical trials, and maintaining a separate record of all patients. The paper received some criticism (though general support for specialist care),^{4,5} but, interestingly, the number of patients treated was not a criterion.

It is nonsense to assume that the quality of a clinical service bears a linear relation to the number of patients treated. Just as a minimum number of patients is necessary to maintain skill, there is probably a maximum number (still to be determined) above which the service becomes overwhelmed. For every new patient seen with breast cancer, up to 15 are now referred with benign disease, implying a total of 1000-1500 new referrals in a practice seeing 60-100 new cancers a year; this is quite enough to ensure continuing skill in a district general hospital committed to subspecialisation.

The management of cancer is not simply a matter of number crunching: 200 patients can be treated just as badly as 20. The quality of a cancer unit should be measured chiefly by the commitment of its staff to subspecialisation; multidisciplinary consensus management; and participation in continuing medical education, audit, and clinical trials—and ultimately by its results.

ADRIAN BALL
Consultant surgeon

Breast Service,
Department of Surgery,
Crawley Horsham NHS Trust,
Crawley Hospital,
Crawley,
West Sussex RH11 7DH

- 1 Langlands A, Calman K. *A policy framework for commissioning cancer services*. Leeds: NHS Executive, 1996. (EL(96)15.)
- 2 Sainsbury R, Haward R, Rider L, Johnston C, Round C. Influence of clinician workload and patterns of treatment on survival from breast cancer. *Lancet* 1995;345:1265-70.
- 3 Gillis CR, Hole DJ. Survival outcome of care by specialist surgeons in breast cancer: a study of 3786 patients in the west of Scotland. *BMJ* 1996;312:145-8. (20 January.)
- 4 Baum M. Specialist surgeons and survival in breast cancer. *BMJ* 1996;312:1155. (4 May.)
- 5 Campbell I. Specialist surgeons and survival in breast cancer. *BMJ* 1996;312:1156. (4 May.)

Authors' reply

EDITOR,—We wish to respond to several points raised in the previously published letters¹ and the above letters about our paper.² None of Ian Campbell's points¹ change our results or their interpretation. We considered selective bias by

analysing our data on the basis of hospital catchment areas. Specialists and non-specialists worked in the same hospital serving the same catchment areas. There was a 4.6% advantage in five year survival for those catchment areas where the specialists saw 44% of patients compared with those where the specialists saw only 9% of patients. As both analyses showed an advantage for specialist care, selection bias is an unlikely explanation of the differences seen.

We relied heavily on tumour size as the main prognostic factor because of our stated concerns about the accuracy of nodal assessment. Using actual tumour size, or six rather than three categories, as a covariate in the survival analysis leaves the hazard ratio for specialists relative to non-specialists unaltered at 0.83.

Our data fit the proportional hazards model well. Analysis of the data in two separate periods of follow up (0-5 and 5-10 years) produced ratios of 0.81 (95% confidence interval 0.70 to 0.93) and 0.89 (0.71 to 1.11) respectively for specialists compared with non-specialists.

To say that our study was biased because we did not take all prognostic factors into account is to say that all non-randomised studies of disease are biased. While randomised controlled trials give an indication of efficacy in a selected population, observational studies show how effective findings are in the general population. Also, when there are difficulties in carrying out a randomised trial, observational studies are the only means of proceeding.

We agree with Michael Baum¹ that the maximum difference that could be attributed to the application of optimal adjuvant treatment is 4%, compared with the 11% advantage that we observed for specialists in patients aged 50-64. We present no evidence to explain this observation. We believe that the only way to test the validity of our findings is to extend the period of the study to the present, with the cooperation of the specialists. This is now being done.

We agree with Adrian Ball, in the letter above, that the number of patients treated should never be the sole criterion for designating a service as a specialist service. The ultimate assessment of survival will come from prospective audit. This will give precise meaning to the term "specialist."

CHARLES R GILLIS
Director

DAVID J HOLE
Principal epidemiologist

West of Scotland Cancer Surveillance Unit,
Ruchill Hospital,
Glasgow G20 9NB

- 1 Specialist surgeons and survival in breast cancer [letters]. *BMJ* 1996;312:1155-6. (4 May.)
- 2 Gillis CR, Hole DJ. Survival outcome of care by specialist surgeons in breast cancer: a study of 3786 patients in the west of Scotland. *BMJ* 1996;312:145-8. (20 January.)

Anaesthesia in the United States

Anaesthesia in Britain is as bad as in the United States

EDITOR,—I am a registrar and, like Peter Hambly¹, on an exchange—but from the Groote Schuur Hospital, Cape Town, to a similar post in Kent. In Britain, as in the United States, anaesthesiologists do not warrant the time of day. Surgeons do not inform us of when they are about to make the incision, dilate the cervix, or stretch the anus; nor do they keep us informed of their progression or imminent termination of the operative procedure. Surgeons do not inform us that they are going to inject adrenaline (into patients with heart disease) or pump the abdomen full of carbon dioxide. We are just there to deal with the consequences. Similarly, we are

expected somehow to know which antibiotics are favoured in particular situations by every surgeon.

Surgeons do not discuss with the anaesthetist planned major surgery, changes to their lists, or variations from the scheduled starting times (I can count on one hand the number of operating sessions that have started on time in the past year). They post their out of hours operating list on the board and expect the anaesthetists to assimilate it by osmosis or extrasensory perception. They would not think of telephoning me to discuss preoperative care and preparations; this results in money being wasted on unnecessary investigations, while patients most in need of specialised investigations do not have them, which results in delayed surgery.

Anaesthetists are seen as dispensable technicians, great for looking after central venous lines and for making frequent adjustments to the light but not to be consulted in greater matters of clinical state. I was also astounded to learn that in Britain anaesthetists seem to have a monopoly on the airway and that doctors who spend their entire working lives in accident and emergency medicine either cannot or will not intubate patients.

Surgeons also fail to realise their responsibilities in transferring patients from the bed to the operating table, since they are never in theatre when this takes place. A case heard in Windhoek (Namibia) Supreme Court found that the surgeon and anaesthetist had to be in the operating room when such a transfer was made.

The training that I received in Cape Town could not have been bettered in any way, either in the variety of clinical experiences or in the many skills imparted. The new, seamless modular training proposed by Calman will produce anaesthetists without this wide range of clinical experiences. Hambly misses British surgeons; I miss the Cape Town surgeons for their unfailing courtesy and their communication skills.

ERROL M CORNISH

Specialist registrar in anaesthetics and intensive care

11 Holmbury Park,
Bromley,
Kent

- 1 Hambly P. Visions of anaesthetic hell. *BMJ* 1996;312:1045. (20 April.)

Anaesthesia is an unpopular specialty in the United States

EDITOR,—Peter Hambly's visions of anaesthesia hell accurately describe the relationship between anaesthetists and surgeons in one of the more aggressive surgical institutions in the United States.¹ A more positive professional ambience is normal in private practice and in my own experience at the University of Virginia. The reasons that Hambly gives for this unfortunate relationship echo my own, but recently issued American training statistics for the specialty paint a dismal picture.

Over the past five years the number of residency positions available each year has fallen from over 1300 to 712 for July 1997. Of those 712 positions offered, only 211 (30%) have been filled. These positions were offered across the country by 118 training programmes, only six of which filled all their positions. Last year saw similarly dismal figures, with only 44% of positions being filled.

Hambly's article goes some way towards explaining why the number of people entering the specialty has fallen so dramatically. Discussions in coffee rooms here suggest that the availability of jobs at the end of training and financial reward have a more pivotal role in attracting new residents. It is now well recognised that hospital

maintenance organisations and capitation are here to stay, and large drops in clinical earnings are predicted. Partners earning high salaries are reducing their holidays and supervising more certified nurse anaesthetists rather than employing another partner and reducing income. It is rumoured that deans of medical schools are dissuading medical students from entering training in anaesthesia.

The staffing crisis in teaching hospitals is only just starting and has yet to be resolved. Academic anaesthesiologists, who are accustomed to spending most of their working week in the laboratory, may be changing their timetables to suit clinical needs. Increased demands on residents may make training even more unattractive to potential applicants.

There are lessons to be learnt for Britain. Structured training as proposed by Calman enables us to predict the number of fully trained staff for a given year but leaves no flexibility to employ specialist registrars beyond their "sell by" date. The ability of "old style" senior registrars to continue in post gives advantages to both them and their employer: trained senior registrars have the cushion of employment until they find a consultant position, while their employing hospital retains a highly skilled professional at a good price.

Although unemployment among anaesthetists seems unlikely in Britain in the present climate, it should not be allowed to occur in the future. It is a reality here in the United States.

COLIN BERRY

Visiting assistant professor of anaesthesiology

Department of Anaesthesiology,
PO Box 238,
University of Virginia,
Charlottesville, VA 22908,
USA

1 Hambly P. Visions of anaesthesia hell. *BMJ* 1996;312:1045. (20 April.)

Trainees have much to gain from working in the United States

EDITOR,—The Shock Trauma Center in Baltimore offers the highest level of trauma care in Maryland and admits 5300 patients annually, with a mortality of 6%. It is a recognised centre of excellence in managing severe trauma. The titles of visiting assistant professor and attending anaesthesiologist, which Peter Hambly holds,¹ indicate an appointment at the University of Maryland and as a specialist in anaesthesia in the trauma centre. The titles are appropriate to the level of training and experience and represent the nomenclature of a different system; they allow experienced British anaesthetists to work in the trauma centre and gain experience that is unavailable in Britain.

Hambly's suggestion that the critical care doctors have no skills in managing airways is wrong. Currently the "critical care attendings" include two internists, two anaesthesiologists, and one surgeon. Eight critical care fellows are surgeons and two are anaesthesiologists. As happens in many facilities around the world, out of normal working hours the resident doctors may not be as familiar with difficult airway problems as specialist anaesthetists are. An experienced anaesthesiologist is always, however, available. It is insulting to suggest that anaesthetists are the only doctors capable of providing critical care when so many physicians and surgeons contribute to this multidisciplinary specialty.

At the centre we were also referred to as "anaesthesia" at times but did not take this as an insult. We have been called "gas man" and occasionally far worse, but not in the United States. Again this reflects different cultures.

There is an element of "you have to prove yourself," and the American system tends to produce more assertive individuals than the British system does. Confrontation on a personal basis is more common in the United States, again reflecting a difference in the system and culture. Any serious confrontations were dealt with by the system in a manner that we can only hope for in Britain and served to improve the communication between the individuals concerned.

It is a shame that Hambly's criticisms were not presented while he was working in the trauma centre. As an attending he was in a position to address problems and improve anaesthesiologists' relationships with surgeons. Any concerns about patients' management could have been referred to the quality assurance programme.

Every system has deficiencies. The way to improvement is through reasoned discussion and good communication. We hope that Hambly's view will not discourage trainees from seeking experience in the United States as there is much to be gained for those willing to take up the challenge.

MICHAEL PARR (consultant in anaesthesia and intensive care),
DAVID GABBOTT (senior registrar in anaesthesia),
PETER BASKETT (consultant anaesthetist), Frenchay Hospital,
Bristol BS16 1LE; JOHN CRISWELL (consultant anaesthetist),
Royal Hampshire Hospital, Winchester SO22 5DG;
ALASDAIR DOW (consultant in anaesthesia and intensive care),
Royal Devon and Exeter Hospital, Exeter EX2 5DW;
ALEXANDER HUNTER (senior registrar in anaesthesia),
PAUL MARTIN (consultant in anaesthesia and intensive care),
RONA PATEY (consultant in anaesthesia and intensive care),
Aberdeen Royal Infirmary, Aberdeen AB9 2ZB;
JERRY NOLAN (consultant in anaesthesia and intensive care),
Royal United Hospital, Bath BA1 3NG; BRYCE RANDALLS
(consultant in anaesthesia and intensive care), Torbay Hospital,
Torbay TQ2 7AA; FERGUS MILLAR (consultant in anaesthesia),
Ninewells Hospital, Dundee DD2 1UB;
PETER DONAHUE (senior registrar in anaesthesia), Royal
Prince Henry Hospital, Sydney, Australia; FRANCES FORREST
(consultant in anaesthesia), IAN RYDER (senior registrar in anaesthesia),
Bristol Royal Infirmary, Bristol BS2 8HW;
GORDON CRAIG (senior registrar in anaesthesia), Royal Perth
Hospital, Perth, Australia

1 Hambly P. Visions of anaesthesia hell. *BMJ* 1996;312:1045. (20 April.)

Educational experience for trainees is better than in Britain

EDITOR,—We wish to address some of the comments in Peter Hambly's personal view of anaesthesia in the United States and to identify some of the benefits of American training.¹ Unlike in Britain, university (teaching) hospitals in the United States frequently have all the surgical specialties (neurosurgery, cardiac surgery, paediatric surgery, obstetrics, etc) under one roof. Trainees are therefore evaluated by a consistent group of faculty (that is, consultants), who communicate daily and regularly review the trainees' experience and their evaluations. In our opinion, this continued and consistent review is superior to the sketchy evaluations that occur in remote hospitals to which trainees rotate in Britain to obtain specialist training.

If Hambly believes that his only role in the anaesthesia that he supervises in Baltimore is to sign the anaesthesia chart then he does a disservice to the nurse anaesthetists and residents (senior house officers and registrars) he supervises and to the patients. The intraoperative consultant is available to assist, teach, and evaluate the person being supervised. Nurses are involved in 90% of cases because there is a large service role and a variable and unpredictable caseload in the trauma centre. Training of residents is more flexible than in Britain, with a faculty anaesthesiologist and an experienced nurse anaesthetist being available 24 hours a day throughout the year to assist and to allow a resident to attend educational sessions during the

working day or be involved in research projects. At the trauma centre in Baltimore the service role does not depend on the participation of residents. This, we believe, adds up to a better educational experience for trainees.

Hambly is wrong about anaesthesiologists' involvement in perioperative medicine. In Baltimore every patient with trauma is seen by the anaesthesia care team on admission (no other medical or surgical service does this); many patients require this attention at the heliport (anaesthesia is the only medical service providing this care). All preoperative evaluations for surgery, intraoperative care, postoperative recovery, and acute pain management are controlled by the anaesthesiologist working as a member of the trauma team. Two of the five critical care consultants are anaesthesiologists. It is not the land of the surgeon intensivist. There is only one consultant surgeon doing critical care. Lastly, the title visiting assistant professor is a courtesy (not a preposterous) title.

COLIN F MACKENZIE
Professor

Department of Anaesthesiology,
University of Maryland School of Medicine,
Baltimore, MD 21201,
USA

1 Hambly P. Visions of anaesthesia hell. *BMJ* 1996;312:1045. (20 April.)

Changes in specialty's name reflect greater involvement in preoperative and postoperative care

EDITOR,—Peter Hambly's visions of anaesthesia in the United States are at considerable variance with my experience.¹ Physicians in the United States who conduct themselves as professionals and show clinical competence and genuine concern for their patients enjoy collegial relationships with other physicians in all specialties, including anaesthesia. The fact that Hambly has not been able to establish this relationship is unfortunate; clearly, a language barrier does not exist (even though the British and American peoples have been characterised as being separated by the Atlantic Ocean and a common language).

Hambly perceives "a loss of control of all three areas of patient care" at a time when several academic departments have changed their names to departments of anaesthesia and perioperative medicine or departments of anaesthesia and pain management. These changes reflect a greater involvement in preoperative and postoperative care. Thirty years ago many departments of anaesthesia added "critical care" to their name, to reflect the extension of intensive care and skills provided by anaesthesiologists in the operating room and recovery room. This care and these skills can now be provided throughout the perioperative period and to patients who do not require surgery. Saidman recently advocated a complete change of name for the specialty, to perioperative medicine and pain management.²

Finally, I am glad that Hambly realises that his opinions are based on a limited view of American anaesthesia and that, despite his "hellish" experience, he has "learnt an enormous amount."

NORIG ELLISON
President

American Society of Anesthesiologists,
520 N Northwest Highway,
Park Ridge, IL 60068-2573,
USA

1 Hambly P. Visions of anaesthesia hell. *BMJ* 1996;312:1045. (20 April.)

2 Saidman LJ. What I have learned from 9 years and 9000 papers. *Anesthesiology* 1995;83:191-7.