- 3 Cochrane AL, Holland WW. Validation of screening procedures *Br Med Bull* 1971;27:3-8.
- 4 Cummings SR, Nevitt MC, Browner SW, Stone K, Fox KM, Ensrud KE, et al. Risk factors for hip fracture in white women. *N Engl J Med* 1995;332:767-73.

Ureteric obstruction secondary to endometriosis

Ultrasonography is best method of detection

EDITOR,—R J Brough and K O'Flynn report a case of bilateral ureteric obstruction secondary to recurrent endometriosis in a patient receiving hormone replacement therapy.¹ This is a rare complication; even the 1.2% rate of ureteric obstruction that the authors quote is probably an overestimate, as a Medline search for 1966-96 shows fewer than 200 reported cases worldwide, with the largest modern series being just eight cases from the combined records of two large teaching hospitals.²

Ureteric obstruction can result in irretrievable renal loss, and early recognition is vital, as Brough and O'Flynn emphasise. Nevertheless, we believe that the authors' suggestion that "routine imaging, either intravenous urography or preferably isotope renography," should be used to assess any patient with endometriosis³ is misleading; more importantly, we believe that this would be unsafe practice as it would expose women, many of whom will be of reproductive age, to harmful irradiation. Ultrasonography is a more appropriate investigation in assessing renal obstruction because it is sensitive in chronic obstruction, relatively inexpensive, and safe.

Ultrasonography will not identify non-obstructive ureteric endometriosis, but neither does urography lead to a confident diagnosis of ureteric involvement.³ This is not to say that we advocate routine use of ultrasonography in all patients with endometriosis. Our view is that patients with loin pain, as in this case, merit renal assessment by ultrasonography. In women with a history of endometriosis (particularly if they have recently started hormone replacement therapy) or with a cyclical history of pain or haematuria, endometriosis should be suspected, looked for, and excluded.

N D P MARCHBANK
Senior registrar in radiology
U PATEL
Consultant radiologist
A E A JOSEPH
Consultant radiologist

Department of Diagnostic Radiology,
St George's Hospital,
London SW17 0QT

- 1 Brough RJ, O'Flynn K. Recurrent pelvic endometriosis and bilateral ureteric obstruction with hormone replacement therapy. *BMJ* 1996;312:1221-2. (11 May.)
- 2 Kane C, Drouin P. Obstructive uropathy associated with endometriosis. *Am J Obstet Gynecol* 1985;151:207-11.
- 3 Maxson WS, Hill GA, Herbert CM, Kaufman AJ, Pittaway DE, Daniell JF, et al. Ureteral abnormalities in women with endometriosis. *Fertil Steril* 1986;46:1159-61.

Authors' reply

EDITOR,—We disagree with several of N D P Marchbank and colleagues' statements. Firstly, although the number of reported cases in the literature is small, the number of patients at risk of ureteric obstruction is likely to increase considerably as hormone replacement therapy becomes more common. Secondly, although the use of ultrasonography to assess the dilated pelvic-iceal system provides static anatomical information, it is unreliable in diagnosing obstruction

and has largely been replaced by diuresis renography,^{1,2} which gives dynamic information on kidney drainage. Isotope renography exposes the patient to a small radiation dose of 0.5 mSv, which is smaller than that resulting from plain abdominal radiography; in these symptomatic postmenopausal women such a dose represents a minimal risk in comparison with the dangers of renal failure in patients in whom the obstruction remains undiagnosed.

R J BROUGH
Registrar
K J O'FLYNN
Consultant

Department of Urology,
Hope Hospital,
Salford M6 8HD

- 1 O'Reilly PH, Lawson RS, Shields RA, Testa HJ. Idiopathic hydronephrosis. The diuresis renogram: a new non-invasive method of assessing equivocal pelviureteric junction obstruction. *J Urol* 1979;121:153-5.
- 2 Koff SA, Thrall JH, Keyes JW. Assessment of hydronephrosis in children using diuretic radionuclide urography. *J Urol* 1980;123:531-4.

Rate of Creutzfeldt-Jakob disease in farmers is not significant

EDITOR,—After the death of four farmers from sporadic Creutzfeldt-Jakob disease Gore concluded that the probability of this occurring by chance was less than 1 in 10 000.¹ Recently Will *et al* reported 10 cases of a new variant of Creutzfeldt-Jakob disease (two of the patients were still alive).² While bovine spongiform encephalopathy could not be proved to be responsible for these cases, it was considered to be the most plausible explanation. It might therefore be concluded that bovine spongiform encephalopathy has been responsible for two new diseases (sporadic Creutzfeldt-Jakob disease and the new variant)—a surprising result. I have checked the robustness of Gore's calculation of probability.

The existence of a new variant of Creutzfeldt-Jakob disease, different from sporadic Creutzfeldt-Jakob disease, seems established because of the unusual neuropathological profile and the young age distribution of the patients. Seven of the eight patients who died were aged 15-34, and I have calculated that the probability of this occurring by chance is 1 in 100 000 000. To check the robustness of this calculation, I increased the expected rate by 2 SD, when a probability of 1 in 1 000 000 was found. The result was thus robust.

My main worry about Gore's calculation of probability is the size chosen for the population at risk. This was taken to be 46 000 farmers (9% of the total of 500 000 farmers). These were only male, full time, lifelong farmers. There was an implied assumption that small and large herds of cattle are looked after by the same numbers of farmers and contain equal numbers of cows with bovine spongiform encephalopathy. If the number of farmers at risk had been regarded as 100 000 or 200 000 (20% and 40%, respectively, of the total number of farmworkers) then the probability of four farmers developing sporadic Creutzfeldt-Jakob disease over five years would be 1 in 500 or 1 in 30 respectively. The first figure is significant at the 3 SD level, but the second is not significant.

It has been assumed that the rate of sporadic Creutzfeldt-Jakob disease is one per million population per year. As recording improves, however, the rate is increasing with time in most countries—except in the United States, where Holman *et al* found a rate of 0.9 (range 0.8-1.1)

cases during 1979-90 (total number of cases, 2614).³ In Austria, where many necropsies are performed, the rate was 1.25 in 1995 (H Budka, personal communication).

I conclude that Gore's calculation is not robust and that bovine spongiform encephalopathy has plausibly caused only one disease, the new variant of Creutzfeldt-Jakob disease.

I thank Dr F James for helpful discussion.

DOUGLAS R O MORRISON
Research scientist

CERN,
European Laboratory for Particle Physics,
CH-1211 Geneva,
Switzerland

- 1 Gore SM. More than happenstance: Creutzfeldt-Jakob disease in farmers and young adults. *BMJ* 1995;311:1416-8.
- 2 Will RG, Ironside JW, Zeidler M, Cousens SN, Estibeiro K, Alperovitch A, et al. A new variant of Creutzfeldt-Jakob disease in the UK. *Lancet* 1996;347:921-5.
- 3 Holman RC, Khan AS, Kent J, Strine TW, Schonberger LB. Epidemiology of Creutzfeldt-Jakob disease in the United States, 1979-1990: analysis of national mortality data. *Neuroepidemiology* 1995;14:174-181.

Performance procedures for seriously deficient professional performance are flawed

EDITOR,—The nuances of Stephen Brearley's editorial on seriously deficient professional performance show the flaw in the performance procedures that have been introduced, at least in so far as patients are concerned.¹ Patients who are on the receiving end of poor performance are more concerned with the accountability of those who were responsible than with the definition of seriously deficient performance. If a patient has suffered injury as a result of treatment by a doctor, or as a result of a failure to treat, then surely he or she must have been at risk. If the patient was at risk then there must be a danger that others will be at risk in the future.

The problem is that the new procedures are aimed solely at a pattern of behaviour. How many ureters must a surgeon divide before a pattern is established? How many babies must suffer asphyxia? What patients want to know is whom doctors will be held accountable to after the first occasion that they cause such a serious injury.

ARNOLD SIMANOWITZ
Chief executive

Action for Victims of Medical Accidents,
1 London Road,
London SE23 3TP

- 1 Brearley S. Seriously deficient professional performance. *BMJ* 1996;312:1180-1. (11 May.)

Tuberculosis is important problem in children with HIV infection in sub-Saharan Africa

EDITOR,—The suggestion by Sebastian B Lucas and colleagues¹ and Charles Gilks² that tuberculosis is not an important problem in children with HIV infection in sub-Saharan Africa deserves further comment. In Lucas and colleagues' necropsy study of children with and without HIV infection in Abidjan, Côte d'Ivoire, only one of the 78 children with the infection had tuberculosis.¹ The suggested explanation—that "with an annual risk of tuberculosis infection of about 2% in many African countries, most children infected with HIV will die without having been exposed to *Mycobacterium tuberculosis*"—may not be valid. Firstly, the risk of