ent from the international non-proprietary name. On the other hand, the World Health Organisation cannot impose on these countries a name that they consider to be unacceptable. The most logical solution would be to set up bilateral discussions between the organisation's experts and the national nomenclature committees with a view to the adoption of a single non-proprietary name for those drugs for which discrepancies exist.

FERNANDO A NAVARRO
Clinical pharmacologist

F Hoffmann-La Roche,
CH-4070 Basle, Switzerland

- 1 George CF. Naming of drugs: pass the epinephrine, please. *BMJ* 1996;312:1315-6. (25 May.)
- 2 Navarro FA. La nomenclatura de los fármacos. II. Las denominaciones comunes internacionales en España. *Med Clin* 1995;105:382-8.
- 3 British Pharmacopoeia Commission. *British approved names*. London: HMSO, 1990.
- 4 World Health Organisation. *Dénominations communes internationales (DCI) pour les substances pharmaceutiques. International nonproprietary names (INN) for pharmaceutical substances*. Geneva: WHO, 1992. (Cumulative list No 8.)
- 5 Fleeger CA, ed. *USAN and the USP dictionary of drug names*. 31st ed. Rockville: USP, 1993.

BMJ should use "paracetamol" instead of "acetaminophen" in its index

EDITOR,—One drug is notably absent from C F George's editorial on the naming of drugs.¹ Paracetamol is widely used for pain and fever from cradle to grave. It is also the substance most commonly taken in deliberate self poisoning,² causing 150 deaths a year in Britain.³ It is a medicine that people travelling abroad take with them or buy when they get to their destination, so everyone should know what to call it. Yet it still has two names: paracetamol and acetaminophen. This creates two problems, one for the general public and one for doctors and medical editors.

Acetaminophen is the United States adopted name,⁴ and in the United States the substance is always and only called acetaminophen. Paracetamol is the recommended international non-proprietary name,⁵ the British approved name,⁴ and the name used for the substance throughout the world outside the United States. Two names for such a common over the counter drug must create confusion and risk for the public travelling to and from the United States. Ideally, the United States would abandon the name acetaminophen and adopt paracetamol. A pragmatic first step would be to label preparations as containing the substance "acetaminophen (paracetamol)" in the United States and "paracetamol (acetaminophen)" in Britain and other countries.

Doctors in Britain who do not know that paracetamol has another name could come unstuck clinically and in their reading. Here the *BMJ* has a role. Since 1992 (volume 305) the index for the *BMJ* has said "see acetaminophen" beside the entry for paracetamol. It then indexes all entries about paracetamol under the American name, even though the articles use the name paracetamol. I wonder why the *BMJ* does this, because it is not the usual or proper practice in Britain. The name paracetamol appears, without reference to acetaminophen, in the index and text of the March 1996 edition of the *British National Formulary*, the 1995 indexes of the *Drug and Therapeutics Bulletin* and the *Lancet*, and the 1995 edition of *A Textbook of Pharmacology*.⁵ It makes no sense for the *BMJ* alone to use acetaminophen as the reference name for this drug.

NICK BRADLEY
General practitioner

Ide Lane Surgery,
Alphington,
Exeter EX2 8UP

- 1 George CF. Naming of drugs: pass the epinephrine, please. *BMJ* 1996;312:1315-6. (25 May.)
- 2 Hawton K, Fagg J. Trends in deliberate self poisoning and self injury in Oxford, 1976-90. *BMJ* 1992;304:1409-11.
- 3 Spooner JB, Harvey JG. Paracetamol overdosage—facts not misconceptions. *Pharm J* 1993;250:706-7.
- 4 Reynolds JEF, ed. *Martindale: the extra pharmacopoeia*, 31st ed. London: Pharmaceutical Press, 1996.
- 5 Ritter JM, Lewis LD, Mant TGK. *A textbook of clinical pharmacology*. 3rd ed. London: Edward Arnold, 1995.

*The *BMJ*'s index uses the medical subject headings published by the United States National Library of Medicine. These headings are used in *Index Medicus* and *Medline* and are an international standard; it is appropriate to use an international standard because the *BMJ* is read worldwide. As Nick Bradley points out, however, the headings do not always reflect current usage, and so at the beginning of this year we decided to make some changes; thus, for instance, the index now refers to cars instead of automobiles and, indeed, to paracetamol instead of acetaminophen.—EDITOR

Cerebral oedema after MDMA ("ecstasy") and unrestricted water intake

Values for plasma osmolality may have been wrong

EDITOR,—S M Matthai and colleagues report two cases of cerebral oedema associated with the syndrome of inappropriate secretion of antidiuretic hormone after ingestion of 3, 4-methylenedioxymethamphetamine (MDMA or "ecstasy").¹ Their letter is, however, somewhat misleading.

In case 1 the patient was reported to have a plasma sodium concentration of 119 mmol/l and a plasma osmolality of 156 mmol/kg. Some time later the urine sodium concentration was 6 mmol/l and urine osmolality 655 mmol/kg. In case 2 the patient was reported to have a plasma sodium concentration of 112 mmol/l and a plasma osmolality of 142 mmol/kg. Urinary sodium concentration was 99 mmol/l and urine osmolality 184 mmol/kg. They do not state whether the osmolalities were measured or calculated.

Plasma osmolality can be calculated from the equation: plasma osmolality = (2 × plasma sodium) + plasma urea concentration + plasma glucose concentration. Some people chose to include potassium concentration in the calculation. To have a plasma osmolality of 156 mmol/kg with a plasma sodium concentration of 119 mmol/l or an osmolality of 142 mmol/kg with a plasma sodium concentration of 112 mmol/l is extremely unlikely. In both instances there would need to be a drastic deficiency in anions and these values are probably not compatible with life. Both values may have been typographical errors and osmolalities in case 1 and 2 were 256 and 242 mmol/kg, respectively.

The features of the syndrome of inappropriate secretion of antidiuretic hormone are, however, a low plasma osmolality and low plasma sodium concentration with an inappropriately concentrated urine. There is usually a high urinary sodium concentration due to low urine volume in association with low circulating aldosterone concentrations. In case 1 urine osmolality was high and urinary sodium concentration low; this could be produced by severe dehydration (causing appropriately raised concentrations of antidiuretic hormone) in which rehydration fluids were deficient in sodium (such as from copious drinking of water). Severe hyponatraemia may also occur in acute alcohol intoxication. In case 2 the urine osmolality was not high (particularly if the plasma osmolality was incorrectly

quoted) and the values do not seem to be compatible with the syndrome of inappropriate secretion of antidiuretic hormone. Instead, they suggest more that simple (severe) water intoxication was the cause of cerebral oedema.

T M COOK
Locum senior registrar in anaesthesia

Southmead Hospital,
Bristol BS10 5NB

- 1 Matthai SM, Sills JA, Davidson DC, Alexandrou D. Cerebral oedema after ingestion of MDMA ("ecstasy") and unrestricted intake of water. *BMJ* 1996;312:1359. (25 May.)

Hyponatraemia must be treated with low water input

EDITOR,—S M Matthai and colleagues have rightly brought our attention again to hyponatraemia as a complication of ingesting 3, 4-methylenedioxymethamphetamine (MDMA or "ecstasy"), adding two cases¹ to at least four previously described. The syndrome of inappropriate secretion of antidiuretic hormone has been proposed as the cause in five of these six cases.¹

I am unconvinced that the clinical and biochemical details presented in these cases point indisputably to the syndrome of inappropriate secretion of antidiuretic hormone. In this syndrome the inappropriateness is not that the urine is concentrated in the presence of dilute plasma, because this can occur in other diseases such as hyponatraemic dehydration complicating diarrhoeal illness, but rather that water is inappropriately conserved by the kidney in the presence of total body water overload. Overhydration explains the natriuresis and high urine sodium concentration that usually accompanies the true syndrome of inappropriate secretion of antidiuretic hormone, despite the hyponatraemia. Natriuresis was not present in case 1,¹ the urine sodium concentration of 6 mmol/l suggesting avid renal sodium conservation; it was also not documented in three of the four previous cases mentioned.¹ In case 2 the stated urine osmolality (184 mmol/kg) is accounted for entirely by the sodium concentration (99 mmol/l) and its attendant anion(s); the urinary urea concentration cannot have been high so this patient cannot have been avidly retaining water at that time. I assume that the stated plasma osmolalities of 156 and 147 mmol/kg should have read 256 and 242 mmol/kg. Urine osmolality was also far from maximal (365 mmol/kg) in one of the other cases alleged to be the syndrome of inappropriate secretion of antidiuretic hormone mentioned by Matthai and colleagues.¹

Severe dehydration, and presumably sodium depletion, caused by hyperthermia and sweating as part of the MDMA intoxication syndrome is well known and supported by experimental evidence in animals.^{2,3} Dehydration would explain why serum antidiuretic hormone concentrations might be appropriately high, even had they been measured. I think that the hyponatraemia was more likely to have been caused by a combination of sodium depletion from severe sweating or vomiting, or both, and excessive water intake without sodium repletion, as suggested by the title of the letter by Matthai and colleagues.¹

Though I disagree with the postulated pathophysiology, I agree that unrestricted water intake is to be discouraged; sports drinks containing salt might be more appropriate for supply at "rave" parties.

Cerebral oedema in cases 1 and 2¹ is not necessarily indicative of total body water overload (the person may still be dehydrated despite ingesting large amounts of water). It has been postulated to be caused by acute water ingestion (or administration), leading to a sudden lowering of plasma osmolality with intracellular water shift.^{4,5}