Secretary of state for health declines to promote free distribution of smoke alarms

EDITOR,—A recent paper in the *New England Journal of Medicine*¹ on a topic of direct relevance to the Health of the Nation's targets for reducing accidents² stimulated a parliamentary question that received a rather disappointing response from the Department of Health.

Residential fires are the second commonest cause of accidental death of children in Britain: each year about 70 children die as a result of residential fires in England and Wales.³ Observational epidemiological studies indicate that smoke alarms are effective in preventing deaths due to fires.⁴ The most compelling evidence for the effectiveness of smoke alarms, however, was provided by a recent report of a controlled intervention trial in which smoke alarms were given away in Oklahoma City.¹

Surveillance data were used to identify an area of the city with a high rate of injury due to fires. This area was then targeted in a programme in which smoke alarms were given away. The annual rate of injuries due to fires per 100 000 population fell by 80% in the target area during the four years after the intervention, but little change occurred in the rest of the city. The programme prevented an estimated 39 deaths or serious injuries. In Britain the rate of deaths due to fires among children in social class V is over 14 times that among children in social class I. Data from the Office for National Statistics show that poor families are least likely to have smoke alarms. Taken together, these facts suggest that families most likely to benefit from smoke alarms

are least likely to be able to afford them. Thus the free distribution of smoke alarms by health visitors and community care organisations seems to be worthy of consideration.

A Labour MP, John Fraser, also thought that this approach was worth further consideration. The parliamentary question he posed and the response provided were as follows:

"Mr Fraser: To ask the Secretary of State for Health if he will encourage the free distribution of smoke alarms by health visitors and community care organisations.

"Mr Malone: Safety in the home is something which all agencies concerned with community care seek to promote. Fire precautions specifically, however, are a matter for my right honourable and learned Friend the Secretary of State for the Home Department."⁵

Has the secretary of state for health lost interest in achieving the targets in the Health of the Nation or is this response simply an attempt to pass on work to another government department? Either way, British children deserve better.

IAN ROBERTS
Director

Child Health Monitoring Unit,
Institute of Child Health,
London WC1N 1EH

- 1 Mallonee S, Istre GR, Rosenberg M, Reddish-Douglas M, Jordan F, Silverstein P, et al. Surveillance and the prevention of residential fire injuries. *N Engl J Med* 1996;335:27-31.
- 2 Department of Health. *The health of the nation: key area handbook. Accidents*. London, DoH, 1993.
- 3 Roberts I. Childhood deaths from residential fires. *BMJ* 1995;311:1381-2.
- 4 Runyan CW, Bangdiwala SI, Linzer M, Sacks JJ, Butts J. Risk factors for fatal residential fires. *N Engl J Med* 1992;327:859-63.
- 5 Written answer. *House of Commons official report (Hansard)* 1996 Jul 24;282:col 50. (No 142.)

Incidence and prognosis of asthma to age 33

Bleeding in first 27 weeks of pregnancy had protective effect

EDITOR,—As an obstetrician with mild asthma, I was interested to read David P Strachan and colleagues' paper on their long term cohort study of the incidence and prognosis of asthma.¹ Several perinatal factors showed an independent association with asthma in the study population: the odds ratio was 2.98 (95% confidence interval 1.33 to 6.66) for antepartum haemorrhage in the third trimester and asthma at age 8-16; 1.63 (1.14 to 2.34) for albuminuria during pregnancy and asthma at age 17-33; and 1.72 (1.11 to 2.67) for maternal smoking (in pregnancy only) and asthma at age 0-7.

The authors do not comment on a significant protective association between bleeding at <28 weeks (that is, threatened miscarriage) and the risk of asthma at age 0-7, for which the odds ratio was 0.40 (0.19 to 0.85). Roughly 10-20% of pregnancies miscarry in the first 12 weeks of pregnancy,² but 30-40% of all women have some bleeding during this period. Bleeding is much less common after 12 weeks, and it would be interesting to know whether the beneficial association in terms of asthma is related to bleeding in the first or the second trimester.

Barker *et al* have suggested that the antecedents of adult cardiovascular disease may be

programmed by events in utero.³ Perhaps the perinatal associations with asthma in childhood and adulthood described by Strachan and colleagues are the first pointers to a causative effect during fetal development.

KIM HINSHAW

Consultant obstetrician and gynaecologist

District General Hospital,
Sunderland SR4 7TP

- 1 Strachan DP, Butland BK, Anderson HR. Incidence and prognosis of asthma and wheezing illness from early childhood to age 33 in a national British cohort. *BMJ* 1996;312:1195-9. (11 May.)
- 2 Alberman E. Spontaneous abortions: epidemiology. In: Stabile I, Grudzinskas G, Chard T, eds. *Spontaneous abortion—diagnosis and treatment*. London: Springer, 1992.
- 3 Barker DJP, Gluckman PD, Godfrey KM, Harding JE, Owens JA, Robinson JS. Fetal nutrition and cardiovascular disease in adult life. *Lancet* 1993;341:938-41.

Respiratory sections of questionnaires are imprecise

EDITOR,—The national child development study is a source of abundant but often inaccurate data that is all too tempting for paper generators. Unfortunately, the respiratory sections of the questionnaires used in the study are so imprecise as to be almost worthless. Diagnoses rather than symptoms predominate and are thus open to error. Questions have differed considerably with each administration of the questionnaires. The use of the phrase "wheeze or whistling," as in too many questionnaires about asthma, reinforces by implication the commonly held public perception that wheeze is something other than whistling—for example, bubbling.

Nearly all of David P Strachan and colleagues' interpretations of the data are based on memory of symptoms experienced from birth or since the previous follow up seven to 10 years previously.¹ The accuracy of questionnaire data for events that occurred before the previous 12 months is known to be extremely poor—a point made well by the authors at the beginning of their discussion but then apparently ignored. In these circumstances, it seems highly probable that the same question asked five times over 25 years in five different ways will generate a positive response in at least half of cases.

This paper shows that an inadequate measure may confirm strong associations, lose weaker relations, and generate false ones. For instance, the association between bleeding in pregnancy and asthma was significant for only one out of six confidence intervals and is therefore likely to be spurious.

The phenomenon of recurrence of wheezing in later adult life after a period free of symptoms is not widely known and its replication is thus useful, but it is not a unique observation: it was reported by Blaire in 1977² and later in a study in Melbourne.³

ROLLO CLIFFORD
Consultant

Department of Paediatrics,
West Dorset Hospital,
Dorchester DT1 2JY

- 1 Strachan DP, Butland BK, Anderson HR. Incidence and prognosis of asthma and wheezing illness from childhood to age 33 in a national British cohort. *BMJ* 1996;312:1195-9. (11 May.)
- 2 Blaire H. Natural history of childhood asthma: 20 year follow up. *Arch Dis Child* 1977;52:613-9.
- 3 Kelly WJW, Hudson I, Phelan PD, Pain MCF, Olinsky A. Childhood asthma in adult life: a further study at 28 years of age. *BMJ* 1987;294:1059-62.

Advice to authors

We receive more letters than we can publish: we can currently accept only about one third. We prefer short letters that relate to articles published within the past four weeks. Letters received after this deadline stand less chance of acceptance. We also publish some "out of the blue" letters, which usually relate to matters of public policy.

When deciding which letters to publish we favour originality, assertions supported by data or by citation, and a clear prose style. Wit, passion, and personal experience also have their place.

Letters should have fewer than 400 words (please give a word count) and no more than five references (including one to the *BMJ* article to which they relate); references should be in the Vancouver style. We welcome pictures.

Letters should be typed and signed by each author, and each author's current appointment and address should be stated. We encourage you to declare any conflict of interest.

Please enclose a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.